

# *Vaccinium bangliangense*, a new species of Ericaceae from limestone areas in Guangxi, China

Yu-Song Huang<sup>1</sup>, Shi-Yue Nong<sup>1,2</sup>, Xing-Kang Li<sup>3</sup>, Gao Xie<sup>1,4</sup>, Yi-Hua Tong<sup>5,6</sup>

**1** Guangxi Key Laboratory of Plant Functional Phytochemicals and Sustainable Utilization, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and the Chinese Academy of Sciences, Guilin, Guangxi, 541006, China **2** College of Life Sciences, Guangxi Normal University, Guilin, Guangxi, 541004, China **3** Guangxi Bangliang Gibbon National Nature Reserve Administration, Baise, Guangxi, 533800, China **4** College of Tourism & Landscape Architecture, Guilin University of Technology, Guilin, Guangxi, 541006, China **5** Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, Guangdong, 510650, China **6** Center of Conservation Biology, Core Botanical Gardens, Chinese Academy of Sciences, Guangzhou, Guangdong, 510650, China

Corresponding author: Yi-Hua Tong ([yh-tong@scbg.ac.cn](mailto:yh-tong@scbg.ac.cn))

Academic editor: Doug Soltis | Received 22 January 2022 | Accepted 24 March 2022 | Published 11 April 2022

**Citation:** Huang Y-S, Nong S-Y, Li X-K, Xie G, Tong Y-H (2022) *Vaccinium bangliangense*, a new species of Ericaceae from limestone areas in Guangxi, China. *PhytoKeys* 194: 23–31. <https://doi.org/10.3897/phytokeys.194.81018>

## Abstract

*Vaccinium bangliangense*, a new species from limestone areas in Guangxi, China, is described and illustrated. It is morphologically most similar to *V. pseudotonkinense* and *V. sciaphilum* in having small and dense obovate leaf blades with a retuse apex, hairy young branches and calyx and campanulate corollas, but can be distinguished from them by the distance of basal gland from petiole, the length of peduncle, pedicel and filaments, the indumentum of calyx tube and corolla and the existence of apical glands on calyx lobes. A table to distinguish the new species from other morphologically similar *Vaccinium* species, as well as colour plates of comparison of key characters, is also provided.

## Keywords

limestone flora, morphology, new species, south-western Guangxi, *Vaccinium*

## Introduction

The genus *Vaccinium* L. contains 450–500 species distributed worldwide (Fang 1991; Fang and Stevens 2005; Vander Kloet and Dickinson 2009). Currently, there are 98 species of *Vaccinium* known from China (Fang 1986; Fang and Stevens 2005; Tong and

Xia 2015; Tong et al. 2018, 2020, 2021a, b, 2022). As one of the most biodiverse regions of China, Guangxi has 27 species and two varieties of the genus *Vaccinium*, including four endemic species, viz. *V. damingshanense* Y.H. Tong & N.H. Xia, *V. napoense* Y.H. Tong & N.H. Xia, *V. crassivenium* Sleumer and *V. cuspidifolium* C.Y. Wu & R.C. Fang (Qin and Liu 2010; Tang 2011; Huang et al. 2015; Tong and Xia 2015; Tong et al. 2018, 2020).

During fieldwork in Bangliang Gibbon National Nature Reserve of Guangxi in June 2021, we discovered a special flowering plant of *Vaccinium* never recorded from Guangxi with the characteristics of inflorescence being shortly racemose, axillary or borne on leafless old stems, peduncle being very short or 4–5 mm long and corolla being broadly campanulate, yellowish-green or tinged reddish. After consulting Flora of China (Fang and Stevens 2005) and other relevant literature (Dop 1930; Pham 1999; Nguyen 2005; Newman et al. 2007; Qin and Liu 2010; Tang 2011; Tong and Xia 2015; Watthana 2015; Tong et al. 2020), as well as comparisons amongst this unknown species and its morphologically most similar species, based on herbarium specimens including types, we confirmed that this species is new to science, which is described and illustrated below.

## Materials and methods

Field surveys have been conducted in flowering and fruiting phases at the type locality. Measurements and assessments of morphological characters were based on the living plants in the wild and the specimens gathered from the type locality. Type specimens were deposited in the herbaria of South China Botanical Garden (IBSC) and Guangxi Institute of Botany (IBK). The comparisons amongst this unknown species, *V. sciaphilum* C.Y. Wu and *V. pseudotonkinense* Sleumer were based on the descriptions from protologues and the examination of herbarium specimens or photos of specimens (including types) at IBK, IBSC, KUN and P (Sleumer 1941; Fang and Wu 1987). The habitat information and threatened factors were recorded during field surveys. The assessment of threatened status of the new species is based on the IUCN Red List of Threatened Species Categories and Criteria and Guidelines for using the IUCN Red List Categories and Criteria (IUCN 2012; IUCN Standards and Petitions Committee 2022).

## Taxonomic treatment

*Vaccinium bangliangense* Y.S. Huang & Y.H. Tong, sp. nov.

urn:lsid:ipni.org:names:77296991-1

Figs 1, 2A–C, 3A, B, 4

**Diagnosis.** *Vaccinium bangliangense* Y.S. Huang & Y.H. Tong belongs to *V.* section *Conchophyllum* Sleumer (1941) and is morphologically similar to *V. pseudotonkinense* Sleumer and *V. sciaphilum* C.Y. Wu in having small and dense obovate leaf blades with a retuse apex, hairy young branches and calyx and campanulate corollas, but can be distinguished from the former by basal glands on leaf blade margin at 0.3–0.8 mm (vs.


**Table 1.** A morphological comparison of key characters of *Vaccinium bangliangense*, *V. sciaphilum* and *V. pseudotonkinense*.

Character	<i>V. bangliangense</i>	<i>V. pseudotonkinense</i>	<i>V. sciaphilum</i>
Distance of basal gland from petiole	0.3–0.8 mm	2.6–4 mm	0.8–1.7 mm
Inflorescence	2-flowered or solitary, peduncle very short or 3–5 mm long	solitary, sometimes 2-flowered, peduncle very short	solitary, sometimes 2-flowered, peduncle very short
Pedicel	5–7 mm long, glabrous	ca. 4 mm long, glabrous	ca. 3 mm long, densely pubescent
Calyx tube	glabrous or sparsely villous	glabrous or sparsely pubescent	densely hispid
Calyx lobes	glabrous or sparsely pubescent abaxially, margin ciliate, apex with a gland	glabrous or sparsely pubescent abaxially, margin ciliate and glandular, apex without a gland	densely hispid abaxially, margin ciliate, apex without a gland
Corolla lobes	pubescent abaxially at apex	glabrous	glabrous
Filaments	densely villous, ca. 2 mm long	sparsely pilose, ca. 1 mm long	glabrous, ca. 1 mm long
Ratio of anther thecae and tubule	1:2	1:2	1:1

2.6–4 mm, Fig. 3C) distance from petiole, inflorescence with very short peduncle or up to 5 mm long (vs. very short, Fig. 2E), calyx lobes with ciliate margin and a gland at apex (vs. with ciliate and glandular margin and without a gland at apex, Fig. 3D) and, from the latter, by inflorescence with very short peduncle or up to 5 mm long (vs. very short, Fig. 2G), longer (5–7 mm vs. ca. 3 mm) and glabrous (vs. densely pubescent, Fig. 2G) pedicel, glabrous or sparsely villous (vs. densely hispid) calyx tube, glabrous or sparsely pubescent (vs. densely hispid, Fig. 3F) calyx lobes with a gland at apex (vs. without a gland at apex, Fig. 2I) and densely villous (vs. glabrous) filaments. A detailed morphological comparison amongst the three species is summarised in Table 1.


**Type.** CHINA. Guangxi Zhuang Autonomous Region: Baise City, Jingxi City, Renzhuang Town, Bangliang protection station, campsite of Huitun, 850 m a.s.l., 2 June 2021, *S. Y. Nong & P. Yang ZYA00199* (holotype: IBSC!; isotypes: IBK!, IBSC!).

**Description.** Small evergreen shrubs, 10–20 cm tall, usually epilithic, sometimes epiphytic on tree trunks. Stem ascending, with long creeping rhizomes. Roots with bead-like swellings 1–3 cm in diam. Young branches brownish-red, grey when older, densely white villous, glabrescent. Leaves dense; petiole 1–2 mm long, red, adaxially sparsely pubescent, glabrescent, abaxially glabrous; blades obovate or broadly obovate, 1–1.7 × 0.6–1.2 cm, leathery, adaxially near margin sparsely villous when young, glabrescent, abaxially sparsely brown glandular hispidulous, base cuneate, basal gland 1 per side, at 0.3–0.8 mm distance from petiole, margin entire, revolute, apex obtuse, slightly retuse; mid-vein impressed adaxially, raised abaxially; lateral veins 2–4 pairs, flat and inconspicuous or impressed adaxially, raised abaxially. Inflorescence shortly racemose, 2-flowered or solitary, axillary or borne on leafless old stems; peduncle very short or 3–5 mm long, base with several bracts, glabrous or sparsely white villous on distal part; bracts ovate, 0.5–1 mm long, ca. 0.5 mm wide, margin ciliate, apex glandular; pedicel 5–7 mm long, glabrous, thickening towards the apex, articulate with the calyx tube; bracteoles 2, adnate to 1–3 mm



**Figure 1.** *Vaccinium bangliangense* **A** habitat **B** habit **C** young branch, showing the leaves with a sparsely villous margin **D** fruiting branch **E** flowering branch with tinged reddish flowers **F** inflorescence borne on leafless old stem with yellowish-green flowers **G** flower (lateral view) **H** flower (vertical view) **I** flower bud **J, K** flowers with opened corolla **L** flower with corolla and stamens removed **M** bracteole **N** transection of ovary **O** calyx tube and disc **P** androecium and adaxial (left), lateral (middle) and abaxial (right) view of a stamen **Q** infructescence **R** seeds. Scale bars: 5 mm (**G–I, P, Q**); 1 mm (**M–O, R**).





**Figure 2.** A–C *Vaccinium bangliangense* **A** flowering branch with tinged reddish flowers **B** flowering branch with yellowish-green flowers **C** fruiting branch **D–F** *V. pseudotonkinense* **D** flowering branch with opened flower **E** flowering branch with unopened flower **F** fruiting branch **G–I** *V. sciaphilum* **G** flowering branch **H** leafy branch **I** fruiting branch. All by Yi-Hua Tong, except **G** by Xin-Xin Zhu.

above the pedicle base, triangular-ovate, ca.  $1 \times 0.5$  mm, margin ciliolate, apex glandular; calyx tube green or purple green, obconical, ca.  $1.5 \times 2$  mm, glabrous or sparsely villous; calyx limb divided nearly to the base; lobes 5, triangular-ovate, ca.  $2 \times 1.5$  mm, both surfaces glabrous or sparsely pubescent abaxially, margin ciliolate, apex glandular; corolla


yellowish-green or tinged reddish, broadly campanulate, ca.  $9 \times 5$  mm, both surfaces glabrous, 5-lobed; lobes triangular-ovate, apical part reflexed, apex acute, pubescent abaxially, glabrous adaxially; stamens 10, 5.5–6 mm long; filaments white, tinged reddish in reddish flowers, flat, incurved, ca. 2 mm long, densely villous; anthers yellow, 4–5 mm long, thecae ca. 1.5 mm long, tubules 3–3.5 mm long, with 2 spurs at the base abaxially, spurs ca. 1.3 mm long, interlocked (that is: the spurs on antesealous stamens extending laterally outside of antepetalous anthers and strongly overlapping with spurs of next antesealous stamens, those on antepetalous anthers strongly hooked outward below spurs of antesealous stamens); disc yellowish, annular, glabrous; style greenish, tinged reddish in reddish flowers, cylindrical, ca. 6.5 mm long, glabrous, stigma truncate; ovary pseudo-10-locular, each locule with several ovules. Berry globose, ca. 8 mm in diam., glabrous, greenish when young, white when mature, fruiting calyx persistent, narrowly triangular-ovate and slightly inflexed; seeds reniform, ventrally compressed, 1.5–2 mm long, testa brownish, cells elongated, with thickened anticlinal walls.

**Phenology.** *Vaccinium bangliangense* was observed flowering from May to June and fruiting from August to October (and up to January of the following year in indoor cultivated plants).


**Etymology.** The specific epithet is derived from the type locality, Bangliang Gibbon National Nature Reserve of Guangxi, China. The Chinese name is given as “邦亮越橘 (pinyin: bāng liàng yuè jú)”.

**Distribution and habitat.** Thus far, *Vaccinium bangliangense* was found only in Bangliang Gibbon National Nature Reserve of Guangxi, China. It usually grows on rocks of limestone hillside or peak at an elevation from 850–900 m, sometimes on the trunks of *Pistacia weinmanniifolia* J. Poisson ex Franchet (Anacardiaceae). The slope direction is to the south and the slope gradient is ca.  $30^\circ$ . The tree layer is up to 8 m tall with a canopy cover of 70% and the shrub and herb layer covers are 85% and 20%, respectively. The associated species include *Quercus phillyreoides* A. Gray (Fagaceae), *Sinosideroxylon pedunculatum* (Hemsl.) H. Chuang var. *pubifolium* H. Chuang (Sapotaceae), *Ardisia pseudocrispa* Pit. (Primulaceae), *Schefflera pesavis* R. Vig. (Araliaceae), *Tetradium calcicola* (Chun ex C.C. Huang) T.G. Hartley (Rutaceae), *Sageretia camelliifolia* Y.L. Chen & P. K. Chou (Rhamnaceae), *Paraboea swinhoei* (Hance) B.L. Burtt (Gesneriaceae), *Bonia amplexicaulis* (L.C. Chia et al.) N.H. Xia (Poaceae), *Bulbophyllum andersonii* (Hook. f.) J.J. Smith (Orchidaceae) etc.

**Conservation status.** *Vaccinium bangliangense* has only been found in Bangliang Gibbon National Nature Reserve of Guangxi, China. As a new species, more subpopulations of *V. bangliangense* could probably be found in similar habitats of surrounding limestone areas in the future. However, wild surveys have been conducted for more than ten years in the area where the new species was found. Only two subpopulations were found in the protected region with a total of fifteen individuals and seven of these are mature. Based on the current data, its population size is very small, and the area of occupancy (AOO) is restricted. According to Guidelines for Using the IUCN Red List Categories and Criteria (IUCN Standards and Petitions Committee 2022), the conservation status of *V. bangliangense* should be assessed as Critically Endangered (CR), based on criteria D of (IUCN 2012).



**Figure 3.** **A, B** *Vaccinium bangliangense* **A** basal gland **B** calyx, arrows showing apical glands **C, D** *V. pseudotonkinense* **C** basal gland **D** calyx, showing lobes with a ciliate and glandular margin **E, F** *V. sciaphilum* **E** basal gland **F** calyx. Scale bars: 2 mm.



**Figure 4.** *Vaccinium bangliangense* **A** SEM micrograph of seed **B** elongated cells of testa. Scale bars: 100 µm.

**Additional specimens examined (paratypes).** CHINA. Guangxi Zhuang Autonomous Region: Baise City, Jingxi City, Renzhuang Town, Bang Liang protection station, 900 m a.s.l., 2 June 2021, *S.Y. Nong & P. Yang* NSY2021060201 (IBK); Guilin City, cultivated in Botany Garden of Guilin, collected from the same locality as above, 16 January 2022, *Y.S. Huang* 2022011601 (IBK).



## Discussion

In Guangxi, another species of *V.* sect. *Conchophyllum*, namely *V. triflorum* Rehder, is also somewhat similar to this new species in the small and dense leaves, short racemes and campanulate corollas, but can be readily distinguished by its thickly leathery and elliptic or obovate-elliptic leaf blades with a strongly rugose adaxial surface. The two species also have allopatric distribution in Guangxi: *V. triflorum* is distributed in Huanjiang County, north Guangxi, while *V. bangliangense* occurs in Jingxi County, southwest Guangxi.

Two kinds of flower colour of *Vaccinium bangliangense* were observed in the wild: the plants growing in shaded habitat always bear inflorescences with yellowish-green flowers and longer peduncles (Figs 1F, 2B), while inflorescences with tinged reddish flowers and shorter peduncles (Figs 1E, 2A) are normally found in sunlit habitat. In addition, the leaf blades of plants growing in shady habitat are thinner and with sparsely white villous margin (Fig. 2B), while those of plants growing in sunlit habitat are thicker and with less hairy or glabrous margins (Fig. 2A). It is speculated that these variations on flower colour, peduncle length and texture and indumentum of leaf blades may be caused by different light intensity.

*Vaccinium bangliangense* is a small shrub with a beautiful tree form and thus an excellent species for landscaping. It is adaptive to limestone areas and has important application value in limestone mountain greening. The first author once inserted one branch of this species into clear water for more than 3 months and surprisingly found that it grew new roots, which indicates that this plant is relatively easy to be cultivated.

## Acknowledgements

The authors are grateful to Mr. Ping Yang for assistance in fieldwork, to Ms. Yu Jing Wei for preparing the SEM micrographs of seed and to Dr. Xin-Xin Zhu for providing the photo of *Vaccinium sciaphilum*. This study was supported by the project of accurate collection of botanical specimen resources in the China-Vietnam border area of Guangxi, the Science & Technology Basic Resources Investigation Program of China (Grant No. 2017FY100100) and National Natural Science Foundation of China (grant no. 31870180).

## References

- Dop P (1930) Vacciniacées. In: Lecomte H (Ed.) Floré générale de l'Indo-Chine, T. 3. Masson et Cie, 698–714.
- Fang RC (1986) Studies on Chinese *Vaccinium*. Acta Botanica Yunnanica 8: 239–258.
- Fang RC (1991) *Vaccinium*. In: Fang RC (Ed.) Flora Reipublicae Popularis Sinicae, Vol. 57(3). Science Press, Beijing, 75–164.
- Fang RC, Stevens PF (2005) *Vaccinium*. In: Wu ZY, Raven PH (Eds) Flora of China, Vol. 14. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 476–504.



- Fang RC, Wu CY (1987) New taxa of the genus *Vaccinium*. Acta Botanica Yunnanica 9(4): 379–395.
- Huang XY, Liu J, Lu ZC, Xu WB (2015) Supplements to Checklist of Vascular Plants of Guangxi, China (III). Journal of Guangxi Normal University: Natural Science Edition 33(2): 115–119.
- IUCN (2012) IUCN Red List Categories and Criteria: Version 3.1. Second edition. IUCN, Gland, Switzerland and Cambridge, [iv +] 32 pp.
- IUCN Standards and Petitions Committee (2022) Guidelines for using the IUCN Red List Categories and Criteria. Version 15. Prepared by the Standards and Petitions Committee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Newman M, Ketphanh S, Svengsuksa B, Thomas P, Sengdala K, Lamxay V, Armstrong K (2007) A checklist of the vascular plants of Lao PDR. Royal Botanic Garden Edinburgh, Edinburgh, 401 pp.
- Nguyen TH (2005) 83. Ericaceae Juss. 1789. In: Ban NT (Ed.) Checklist of plant species in Vietnam, Vol. 3. Agriculture Publishing House, Hanoi, 437–450.
- Pham HH (1999) An Illustrated Flora of Vietnam, Vol. 1. Youth Publishing House, Hanoi.
- Qin HN, Liu Y (2010) A checklist of vascular plants of Guangxi. Science Press, Beijing.
- Sleumer HO (1941) Vaccinioideen-Studien. Botanische Jahrbücher für Systematik 71(4): 375–510.
- Tang SC (2011) Vacciniaceae. In: Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences (Ed.) Flora of Guangxi, Vol. 3. Guangxi Science & Technology Publishing House, Nanning, 644–657.
- Tong YH, Xia NH (2015) *Vaccinium damingshanense* sp. nov. (Ericaceae) from Guangxi, China. Nordic Journal of Botany 33(1): 74–78. <https://doi.org/10.1111/njb.00581>
- Tong YH, Ye XE, Wu L, Nguyen TTH, Xia NH (2018) A new record of Ericaceae from China: *Vaccinium eberhardtii* Dop. Guihaia 38: 1595–1598. <https://doi.org/10.11931/guihaia.gxzw201709012>
- Tong YH, Huang YS, Ye XH, Cai ZY, Xia NH (2020) *Vaccinium napoense*, a new species of *V.* sect. *Conchophyllum* (Ericaceae) from Guangxi, China. Nordic Journal of Botany 38(12): e02773. <https://doi.org/10.1111/njb.02773>
- Tong YH, Zhao WL, Wang BM, Liu ED, Cai J, Guo YJ (2021a) *Vaccinium motuoense* (Ericaceae), a new species from Xizang, China. PhytoKeys 181: 105–111. <https://doi.org/10.3897/phytokeys.181.71522>
- Tong YH, Zhu YY, Ye XH, Ye XE, Yang CZ, Xia NH (2021b) *Vaccinium zhangzhouense*, a new species endemic to Fujian, China. Nordic Journal of Botany 39(7): e03091. <https://doi.org/10.1111/njb.03091>
- Tong YH, Fritsch PW, Nguyen TTH, Averyanov LV, Kuznetsov AN, Kuznetsova SP, Nuraliev MS (2022) Novelties in Vietnamese *Craibiodendron*, *Lyonia* and *Vaccinium* (Ericaceae). Phytotaxa 538(1): 21–34. <https://doi.org/10.11646/phytotaxa.538.1.2>
- Vander Kloet SP, Dickinson TA (2009) A subgeneric classification of the genus *Vaccinium* and the metamorphosis of *V.* section *Bracteata* Nakai: More terrestrial and less epiphytic inhabit, more continental and less insular in distribution. Journal of Plant Research 122(3): 253–268. <https://doi.org/10.1007/s10265-008-0211-7>
- Wattana S (2015) Ericaceae. In: Newman M, Barfor A (Eds) Flora of Thailand, Vol. 13, Part 1. The Forest Herbarium, Bangkok, 101–141.