

PROFILING THE CAROTENOIDS OF MICROALGA (*Scenedesmus obliquus*) EXTRACT BY HPLC AND ITS ANTIOXIDANT CAPACITY

 Yaşar Aluç¹,  Osman Kök²,  İlhami Tüzün³

¹ Kırıkkale University, Scientific and Technological Research Application and Research Center, Kırıkkale, Turkey

² Kırıkkale University Graduate School of Natural and Applied Sciences, Department of Biology, Kırıkkale, Turkey

³ Kırıkkale University, Faculty of Arts and Sciences, Department of Biology, Kırıkkale, Turkey

*Corresponding Author:

E-mail: alucyasar@kku.edu.tr

(Received 15th December 2021; accepted 17th January 2022)

ABSTRACT. The extracts of the algae species *Scenedesmus obliquus* isolated from Kapulukaya Reservoir (Kırıkkale, Turkey) were analyzed to determine the composition, content and antioxidant properties of carotenoids. The HPLC analysis was applied to identify carotenoids species and to determine their contents. Methanol–acetonitrile–water (84:14:2, v/v/v) and methylene chloride (100%) used in the gradient solvent system resulted in an acceptable separation. Of the 22 resolution peaks assigned to carotenoids either tentatively or positively, the first eight were identified to be epoxy containing while the remaining ones as main carotenoids namely, all-*trans* forms of lutein, β -carotene, α carotene, zeaxanthin, 13- or 13'-*cis*-lutein, 9- or 9'-*cis*-lutein, *cis*-lutein, 13- or 13'-*cis*- β -Carotene, 9- or 9'-*cis*- β -Carotene, 9- or 9'-*cis*- α -carotene and *cis*-neoxanthin. Quantitatively, *S. obliquus* was predominated by all-*trans* isomers of lutein and β -carotene, being 83.74 % (2.52 mg g⁻¹) in total carotenoids (TC). Antioxidant capacity assays of DPPH and FRAP showed considerably low effects of *S. obliquus* extracts compared to standards. This was attributed to the lesser existence of *cis* isomers within derivatives. Multiple regression was utilised to partition the antioxidant effect of both assays and, revealed an estimation of FRAP dominating over DPPH. A need was arisen on the exploration of isomerisation mechanisms of the carotenoids compounds in order to better evaluate the species for a probable potential of further practical applications.

Keywords: *Microalgae, Scenedesmus obliquus, carotenoids, HPLC analysis, DPPH and FRAP assays.*

INTRODUCTION

Owing to their structural and functional roles, carotenoids are indispensable in photosynthetic life. Their task in plants is ample. During photosynthesis, they harvest the light, transfer the energy and protect the photosynthetic apparatus against photooxidative damage [1]. The antioxidant property is in general operated via scavenging singlet molecular oxygen and peroxy radicals and increasing the stability of the photosynthetic apparatus [2]. Protective functions of carotenoids via their anti-oxidant activities [3, 4], have also been well accounted for the human [5]. The natural carotenoids are well known to have a unique complexity owing to a mixture of various isomers in their structure and chemical formations with other bioactive compounds. Therefore, they are preferred over

synthetic counterparts dominated by all-trans compounds [6] in direct human consumption.

Boosted by increasing results about the limitations of the use of synthetic carotenoids due to potential safety concerns [7, 8] microalgae have recently received particular attention as a natural source of carotenoids [9], amongst other bioactive compounds such as carbohydrates, lipids, proteins, alkaloids, fatty acids etc. that can be utilized for commercial use in a wide variety of industries (i.e. food and feeds, health, agriculture, cosmetic and energy (biofuel) [10, 11, 12].

Numerous studies reported sufficient isolation of various types of antioxidant carotenoids compounds including β -carotene from *Dunaliella salina* [13, 14], zeaxanthin and lutein from *Scenedesmus almeriensis*, [15, 16] astaxanthin and lutein in *Chlorella zofingensis* and *Chlorella vulgaris* [17, 18, 19, 20], astaxanthin from *Haematococcus pluvialis* [21, 22] and *Chlorococum* sp. [23], α -tocopherol from *Nanochloropsis oculata* [24]. Various technological and manipulative applications, such as environmental cues that induce carotenoid accumulation through the regulation of carotenogenesis, are still pursued on the aforementioned and other potential species to obtain compounds at desired content and efficacy [25]. In this study, *Scenedesmus obliquus* which has recently gained attention for its potential to be a model species [26, 27] are therefore explored by means of profiling its carotenoids and antioxidant activity performance.

MATERIALS AND METHODS

Collection and culturing of microalgae

Scenedesmus obliquus were isolated from fresh water samples of the River Kızılırmak in Kırıkkale Province (Turkey). The species identification was based upon molecular characterization, using 18S rRNA, ITS, *rbcl* and 16S molecular markers [19].

Basal Bold Medium (BBM) [28] was selected for the growth of *S. obliquus* using an illumination of 16:8 h light–dark cycle of 4000 lux light intensity under temperature conditions kept at 25 ± 1 °C. Cell counting, chlorophyll *a* and dry biomass were measured to monitor the growth of cultures until the stationary phase, at the middle of which (Day 18) the harvesting was executed. Centrifugated biomass at 3000 rpm for 10 min. was lyophilized for 48 hours at -83 °C and 1.33 Pa [29] and stored at -80 °C until used.

Extraction and profiling Carotenoids

The details of both the extraction procedure [30] and the analysis of carotenoids were given in a previous study by Aluç et al., 2018 [19]. Briefly, saponification was first applied to a 0.1 g microalgae sample using 3 mL hexane–ethanol–acetone–toluene (10:6:7:7, v/v) for 1h and then the separation of carotenoids was achieved by consecutive usage of hexane and sodium sulphate under dimmed light and nitrogen gas. Collected extracts dried by evaporation were subjected to HPLC analysis in mobile phase solvent. Two mobile phases (A and B) of methanol–acetonitrile–water (84:14:2, v/v/v) and methylene chloride (100%) with an arranged flow rate of 0.6 mL min⁻¹ were used in a gradient manner from 100% A and 0% B. Reproducibility, the limits of detection (LOD) and quantification were calculated as described in Aluç et al. [19].

Antioxidant capacity assays

For the extraction of antioxidants, the microalgae biomass (0.1 g) was mixed with 1 ml methanol-toluene (3:1) in a volumetric flask. Following homogenization, insoluble biomass was separated from supernatant by centrifugation at 13500 rpm for 10 min. [30].

DPPH (2,2-Diphenyl-1-picrylhydrazyl) activity of the *S. obliquus* extract solutions was determined based on the method described by Blois, 1958 [31] Volumes of 62.5 μL from each extract solutions prepared with methanol at concentrations of 2, 10, 25, 50, 100 $\mu\text{g mL}^{-1}$, respectively, were added to a mixture of methanol (125 μL) and DPPH (62.5 μL) and then left for incubation in the dark at room temperature. The reaction values measured at 515 nm against methanol extract as blank were used to calculate the percentage radical scavenging activity as below.

$$\% \text{ RSC} = \frac{(\text{A control} - \text{A sample})}{\text{A control}} \times 100$$

So that the control included DPPH and methanol while sample consisted of microalgae extract, DPPH and methanol. The EC_{50} values which were calculated from the plot of scavenging activity against the concentration of sample represented the half maximal potency of microalgal extract to scavenge DPPH radicals. For the Ferric Reducing Antioxidant Power Assay (FRAP), a 1 ml of *S. obliquus* extract solutions prepared at graded concentrations (2, 5, 10, 25, 50, 100 $\mu\text{g mL}^{-1}$) was mixed with phosphate buffer (2.5 mL, 0.2 M, pH 6.6) and potassium ferricyanide [$\text{K}_3\text{Fe}(\text{CN})_6$] (200 μL , 1%) and left for incubation at 50 °C for 20 min. Trichloroacetic acid (TCA, 10%, 200 μL) was added to the mixture and centrifuged at 3750 rpm for 10 min. [32]. A 125 μL of the upper layer was separated and mixed with 25 μL distilled water and 20 μL FeCl_3 (0.1 %). The absorbance of the mixture was measured at 665 nm by a UV-spectrophotometer.

Statistics

Means of data were obtained from triplicate analysis during HPLC analysis. Results from antioxidant assays were obtained via triplicate experiments. Relationships between antioxidant activity and concentrations of algal extracts and standards were depicted by regression analysis. Stepwise multiple regression analysis was used to explore substantially the partial contribution of antioxidant activity at each assay (DPPH and FRAP), using STATISTICA (version 7) statistical software.

RESULTS AND DISCUSSION

Identification of carotenoid peaks

For the identification of carotenoids in *S. obliquus*, the photoisomerized lutein and β -carotene standards were subjected to HPLC analysis in order to obtain spectral characteristics and Q ratio values, according to which the peaks obtained from *S. obliquus* samples were designated. Considering the results obtained from these standards in the previous study by Aluç et al., 2018 [19], solvent selection was considered reliable, indicated by the range of retention time and Q ratio values of these standards [33, 34]. The used solvent system was efficient implied by the the retention factor (k values between 0.76 and 7.76) [34, 35] and the mobile phase used for the separation of carotenoid compositions was adequate decided by the separation factor (α) values always

being higher than 1. All peaks and standards also exhibited a high percentage of purities being higher than 90% [19].

The chromatogram of the extract solution from the *S. obliquus* cells revealed 22 resolution peaks assigned to carotenoids (Figure 1 and Table 1). All-*trans* forms of lutein, zeaxanthin, α -carotene and β -carotene were assigned to peaks 12, 13, 19 and 20, respectively (Figure 1 and Table 1). They were all positively identified based on retention time and Q ratio values of standards except for all-*trans* α -carotene, designation of which was tentative and made by the comparison of retention time and absorption spectra values given in the literature (Table 1 and 2) [34, 35, 36]. The peaks 1 through 8 were again tentatively identified to be the epoxy-containing compounds relying upon spectral characteristics and retention behavior reported in the literature (Figure 1 and Table 2) [36, 37, 38].

As given in our previous study [19], during HPLC analysis of standards, an acceptable reproducibility was reached since RSD values were found to be lower than 2.37 and 4.76 for retention times and integrated areas, respectively. Linearities with their markedly high correlation coefficients (> 99%) also indicated that a good purity in the analysis was attained. The calculated LOD values were 0.013, 0.019, 0.024, 0.022 $\mu\text{g/ml}$ for lutein, zeaxanthin, β -cryptoxanthin and β -carotene, respectively while the corresponding values of LOQ were found to be 0.041, 0.058, 0.071 and 0.068 $\mu\text{g mL}^{-1}$.

Table 1. Retention time, retention factor (*k*), separation factor (α), peak purity and resolution of carotenoids in *S. Obliquus*.

Peak no.	Compound	Retention time (min)	k	α	Peak purity (%)	Resolution
1	Auroxanthin	8.61	0.76	0.00	96.61	9.58
2	Auroxanthin	9.99	1.04	1.37	99.74	2.34
3	Neochrome	11.26	1.31	1.26	98.01	2.48
4	Neoxanthin	12.24	1.50	1.14	98.75	1.63
5	<i>Cis</i> -neoxanthin	13.82	1.82	1.22	99.93	2.97
6	Neoxanthin	14.39	1.94	1.06	99.81	0.95
7	Neoxanthin	15.47	2.16	1.12	99.49	1.68
8	Neochrome	16.52	2.37	1.10	96.54	1.91
9	<i>Cis</i> -neoxanthin	17.67	2.61	1.10	97.67	2.09
10	13-or 13'- <i>cis</i> -lutein	20.60	3.20	1.23	99.83	4.47
11	13-or 13'- <i>cis</i> -lutein	22.01	3.49	1.09	99.99	2.39
12	All- <i>trans</i> -lutein	23.47	3.79	1.09	96.08	3.40
13	All- <i>trans</i> -zeaxanthin	26.19	4.34	1.15	99.99	6.88
14	9-or 9'- <i>cis</i> -lutein	27.12	4.53	1.04	99.97	1.90
15	9-or 9'- <i>cis</i> -lutein	28.35	4.78	1.06	93.64	2.09
16	<i>Cis</i> -lutein	29.52	5.02	1.05	99.91	2.05
17	<i>Cis</i> -lutein	31.07	5.34	1.06	97.92	3.33
18	13-or 13'- <i>cis</i> - β -carotene	36.93	6.53	1.18	99.70	10.22
19	All- <i>trans</i> - α -carotene	37.77	6.70	1.03	98.40	1.80
20	All- <i>trans</i> - β -carotene	41.32	7.43	1.11	99.81	7.86
21	9-or 9'- <i>cis</i> - α -carotene	41.93	7.55	1.02	98.75	1.48
22	9-or 9'- <i>cis</i> - β -carotene	42.93	7.76	1.03	98.85	2.40

Table 2. Assignment data for all-*trans* and *cis* forms of carotenoids their amounts (mg g⁻¹) in *S. Obliquus*.

Peak no.	Compound	Retention time (min.)	λ (nm, inline)	λ (nm, reported)	Q-ratio ^a	Amount (mg g ⁻¹)
1	Auroxanthin	8.61	400	421	-	0.000
2	Auroxanthin	9.99	400	422	-	0.000
3	Neochrome	11.26	399	421	326	0.000
4	<i>cis</i> -Neoxanthin	12.24	401	425	326	0.000
5	<i>cis</i> -Neoxanthin	13.82	405	429	456	0.000
6	Neoxanthin	14.39	418	439	-	0.000
7	Neoxanthin	15.47	418	440	469	0.000
8	Neochrome	16.52	400	422	448	0.000
9	<i>cis</i> -Neoxanthin	17.67	415	436	465	0.000
10	13-or 13'- <i>cis</i> -Lutein	20.60	331	442	468	0.000
11	13-or 13'- <i>cis</i> -Lutein	22.01	331	440	467	0.079
12	All- <i>trans</i> -lutein	23.47	333	446	473	1.368
13	All- <i>Trans</i> -zeaxanthin	26.19	-	452	477	0.008
14	9-or 9'- <i>cis</i> -Lutein	27.12	-	441	469	0.000
15	9-or 9'- <i>cis</i> -Lutein	28.35	327	444	470	0.016
16	<i>cis</i> -Lutein	29.52	328	442	470	0.032
17	<i>cis</i> -Lutein	31.07	-	447	473	0.004
18	13-or 13'- <i>cis</i> - β -carotene	36.93	343	451	-	0.227
19	All- <i>trans</i> - α -Carotene	37.77	340	448	475	0.000
20	All- <i>trans</i> - β -Carotene	41.32	344	454	479	1.151
21	9-or 9'- <i>cis</i> - α -Carotene	41.93	-	444	470	0.000
22	9-or 9'- <i>cis</i> - β -Carotene	42.93	346	449	474	0.121
Total						3.008

a: A gradient mobile phase of methanol-acetonitrile-water (84:14:2, v/v/v) and methylene chloride (from 100:0, v/v to 45:55, v/v) was used.
 b: A gradient mobile phase of methanol-acetonitrile-water (84:14:2, v/v/v) and methylene chloride (from 100:0, v/v to 45:55, v/v) was used by Inbaraj et al.(2006)
 c: A gradient mobile phase of methanol-2-propanol (99:1, v/v) and methylene chloride (from 100:0, v/v to 70:30, v/v) was used by Chen et al., (2004)
 d: A gradient mobile phase of methanol-2-propanol (99:1, v/v) and methylene chloride (from 100:0, v/v to 70:30, v/v) was used by Liu et al., (2004)




Fig. 1. The HPLC chromatogram of the carotenoids obtained from *Scenedesmus obliquus* (left) and the profiles of all-trans carotenoids (right)

Quantification of carotenoids

The *S. obliquus* eliquates contained 3.09 mg g^{-1} of total carotenoids (TC), dominated by the two main derivatives, trans and cis forms of lutein and β -carotene, with an equal contribution of each being 49.83% (1.49 mg g^{-1}). All-trans-zeaxanthin was represented by a small amount in TC, being 0.26 % (0.008 mg g^{-1}), (Table 2). In a previous study by Aluç et al. [19], another species of the same genus, *Scenedesmus regularis* was represented with a similar carotenoids composition whereas the total amount of carotenoids was markedly higher (7.14 mg g^{-1}) than the species *S. obliquus*, used in the present study.

Various factors are involved in the biosynthesis of carotenoids or the distribution of derivatives in algae. It is not only species-specific [5] but also may differ within the different strings of the same species [39]. Additionally, variations may emerge due to different culture conditions as stated either in studies of direct determination of carotenoids [40] or in those applying stress factors to increase carotenoids content within a potential [39, 41]. In spite of such facts, our findings can still be made comparable to those given in previous studies. For example, in a study carried out on 36 different microalgae biomass [40], a total carotenoid of $0.44 \pm 0.06 \text{ mg g}^{-1}$ D.W. in ethanol/water extracts of *Scenedesmus obliquus* was reported, being comparatively lesser than that in our study. But, relatively high phenolic content was also found as to be $1.94 \pm 0.16 \text{ mg GAE. g}^{-1}$ D.W. for the same species. Patnaik and Mallick, 2015 [26] conducted a research on the potential of *S obliquus* in β -carotene production under growth conditions optimized by the manipulations of citrate. Total carotenoids content was found to be around 2.80 mg g^{-1} under control and 2.5 mg g^{-1} under optimized conditions. The corresponding values of β -carotene were around 0.82 mg g^{-1} . and 0.57 mg g^{-1} . These findings are all similar or slightly less than that obtained in our study. In another study by Ho et al., 2014 [39], the cell growth and lutein increment in six *S. obliquus* isolates were enhanced under different light exposures. Lutein content varied between strings with the lowest value of $1.86 \pm 0.11 \text{ mg g}^{-1}$ (*S. obliquus* ESP-5) and the highest of $3.63 \pm 0.12 \text{ mg g}^{-1}$ (*S. obliquus* FSP-3). Three strains of *Scenedesmus* sp. among four classes of chlorophyta were also presented to have concentrations ranging between $0.53\text{--}5.57 \text{ mg TC g}^{-1}$ DCW [42]. Distinctively

high yield of TC (10.45 mg g^{-1} DCW) was also noted for *Scenedesmus* sp. [43] Guedes et al., 2011 [44] used a strain (M2-1) of *S. obliquus* to determine the optimal antioxidant capacity in relations to the carotenoid content varying during growth phases under cross-combination of pH and temperature conditions. They found markedly higher lutein ($203.57 \pm 1.41 \text{ mg mL}^{-1}$) and β -carotene ($18.2 \pm 0.33 \text{ mg mL}^{-1}$) contents from the biomass at the early exponential growth phase.

Antioxidant capacity

As a general rule, the background in antioxidant assays are to imitate the biochemical reactions within a cell. Yet, a wide variety of assays which has been evolved within this approach sustain some degree of complexity and complications during applications due to, amongst others, multifunctional nature of the natural antioxidants or phytochemicals [45]. The composition of the biological component, oxidizable substrate, the media used during oxidation experiment and methods to measure antioxidant activity can all vary, and multitude of tests are indeed needed to take account of the chemistry that would better respond to the targeted chemical [46]. Moreover, variations also occur in mechanisms viz., radical scavenging, metal ion transition, hydrogen peroxide decomposition etc. through which antioxidants exert their effect. Therefore, in this study, the assays were selected to measure antioxidant activity regarding two different mechanisms to reveal the whole picture as much as possible as suggested by Apak et al. [46] DPPH, represented the hydrogen atom transfer reactions or (HAT) and FRAP the single electron transfer reactions (ET). The antioxidant activities of the algal extract determined by DPPH and FRAP assays were evaluated against those of ASC and BHT. Concentrations of *S. obliquus* extract and both standard solutions produced positive and significant correlations in DPPH and FRAP assays, the latter being represented with stronger regression coefficients. (Figure 2a, b). The IC_{50} values calculated for DPPH activities followed an increasing order of ASC ($IC_{50} 0.91 \text{ mg mL}^{-1}$) < BHT ($IC_{50} 3.59 \text{ mg mL}^{-1}$) < *S. obliquus* extract ($IC_{50} 115.62 \text{ mg mL}^{-1}$). ASC and BHT equivalent values of microalgae extract are also calculated as given in Table 3.

Table 3. IC_{50} values of *S. obliquus* and standards (BHT and ASC) in DPPH assay, and equivalent BHT and ASC of *S. obliquus* in both assays.

	IC_{50} (mg mL ⁻¹)	Equivalent $\mu\text{mol g}^{-1}$ DW (DPPH)	Equivalent $\mu\text{mol g}^{-1}$ DW (FRAP)
<i>S. obliquus</i> extract	115.617		
BHT	1.393	3.59	1.02
ASC	0.211	0.91	0.48




Fig. 2. Collective display of DPPH radical sweeping (a) and FRAP inhibition (b) effect of *S. obliquus* extract. Column legends and x axis title are shown in graph b.

In order to explore the respective contribution of each mechanism (DPPH and FRAP) on the antioxidant reactions, stepwise multiple regression analysis was applied. Despite the existence of collinearity, a well known problem encountered in multiple regressions, the analysis was proceeded to obtain substantive information. Each explanatory variable alone showed linearity on the carotenoid concentration but they differed from each other (Friedman ANOVA, $p=0.0253$). Thus, the aim was set to find out the best subset of explanatory variable(s) with the strongest contribution rather than attaining a forecasting statistic. This was achieved best with the (forward) stepwise elimination procedure since it allows the inclusion of all variables initially and then removes the variable that contributes the least (usually that with the smallest partial correlations) to explaining the response variable (Carotenoid concentration), until explanatory variable(s) remaining in the model having a significant partial correlation coefficient (Table 4) [47]. As for our data, it was found that DPPH mechanism was excluded but the FRAP mechanisms remained in the model indicating its significant contribution on the carotenoid concentration ($R^2= 0.998$, $p = 0.035$). Semipartial correlations of FRAP and DPPH were 0.965 and 0.865, respectively, indicating also the approval of selection. Cook’s distance values were used to detect the influential data observed in linear regression, and analysis revealed that residuals of all observed cases were validated by their accepted values within the model except for case 1 (concentration of 2 mg mL⁻¹), which was excluded. Overall, it could therefore be concluded that the effect of antioxidant activity of

carotenoids in the *S. obliquus* occurred predominantly by the mechanisms exerted via reduction of metal ions (FRAP).

Table 4. Multiple regression results between dependent (*S. obliquus* extract concentration) and independents (DPPH and FRAP values). Significant values were shown in bold

	Beta	Std.Err.	B	Std.Err.	t(2)	p-level
Intercept			2.44	1.30	1.88	0.20
DPPH	0.32	0.13	0.75	0.31	2.45	0.13
FRAP	0.68	0.13	1.58	0.30	5.18	0.04
Regression Summary for Dependent Variable (<i>S. obliquus</i> extract concentration)						
R= 0.999, R ² = 0.998, Adjusted R ² = 0.997						
F(2,2)=851.65, p<0.001						

It is also clearly understood that considering the DPPH assay, the methanolic extracts of *S. obliquus* exhibited markedly low radical scavenging activity compared to the standards, BHT and ASC. Considerably high amounts of carotenoid extracts (i.e. 100mg mL⁻¹) reached only to a maximum of 42.69% scavenging ability (IC₅₀:115.62 mg mL⁻¹) whereas standards of ASC (IC₅₀: 0.21 mg mL⁻¹) and BHT (IC₅₀:1.39) had strong scavenging effects around 80%) despite their small amounts i.e. 0.1 and 3.0 mg mL⁻¹, respectively.

The FRAP assay also generated similar results; the effect of carotenoid extract being the lowest (i.e. 0.68 at 100 mg mL⁻¹) followed by BHT and ASC (0.65 and 1.39 at 0.1 mg mL⁻¹), respectively. Higher antioxidant effects of standards compared to the sample may be an expected result as it is often found in previous studies. For example, Hu et al., 2008 [48] utilized DPPH assay and found that antioxidant capacities of the algal (*Dunaliella salina*) carotenoid extract was relatively lower than α -Tocopherol used as a standard. But, still the extent of the difference between algal carotenoids extracts and standards is so high and this may make our species, *Scenedesmus obliquus*, doubtful about its efficiency or its preference over chemical counterparts in practical applications. Despite considerable amount of total carotenoids, the composition consisting of only trans forms (All-*E* isomers) and/or lacking cis forms (*Z* isomers) might be the underlying reason for the relatively low antioxidant activity of *Scenedesmus obliquus*. As demonstrated by numerous studies [49, 50, 51], antioxidant activity showed variations between both isomers of the same compound as well as between different compounds of the carotenoid. In a well documented study by Honda et al., 2019 [52], the supporting results from researches are overwhelming to conclude that the *Z* isomers display antioxidant activity as well as bioavailability and functionality greater than or equal to those of all-*E* counterparts in carotenoids. Especially, *Z* isomers of lutein are explicitly evidenced in favour of higher antioxidant activities. Consequently, the results about antioxidant potential in our study are susceptible to the composition of carotenoids that has occurred in specific cultur conditions and timing of harvesting as much as extraction solution and the type of antioxidant assay deployed. The study of Guedes et al., 2011 [44] supports well the variations of compounds and changes in antioxidant activity throughout the

culture. They proved that antioxidant activity of *S. obliquus* showed variations during 17 days of incubation period, depending on the variations occurring in the rate of carotenoid compounds. Despite no isomer data was given, in another study conducted by Guedes et al., 2013 [53], particularly the high lutein content (*i.e.*, $2.69 \pm 0.09 \text{ mg g}^{-1}$) recorded in the *Scenedesmus obliquus* M2-1 extract was responsible for the observed high antioxidant capacity but together with the putative synergistic effect with β -carotene and neoxanthin recorded to a lesser amount. They also demonstrated that the intracellular extracts of *Scenedesmus obliquus* outperformed those of 18 species of cyanobacteria (prokaryotic microalgae) and 23 species of (eukaryotic) microalgae almost four fold the highest value in total antioxidant activity to scavenge ABTS. Hu et al., 2008 [48] found that the antioxidant activity of carotenoid extract from *D. Salina* was higher than the each all-trans forms carotenes such as all *trans*-zeaxanthin, all *trans*-lutein, all *trans*- β -carotene and all *trans*- α -carotene used as standards. They commented that this was caused by the *cis* forms of carotenoids existed additionally in the total carotenoids content of *D. salina*. Similar findings were presented also by other studies conducted on algae [54] and on vegetables and fruits [55]. Regarding the few but successful findings present in the literature, the evaluation of our results on the antioxidant ability of this species could be made more sufficient by further attempts that would consider certain other factors such as culture conditions, extract solution, isomerization and different antioxidant assays.

CONCLUSION

The amount of total carotenoids that the string of a local species, *Scenedesmus obliquus*, contained could be considered as sufficient, relying upon comparisons to other species of *Scenedesmus* sp., or some other microalgal species. Very high domination of lutein and β -caroten in the composition of total carotenoids ascertained for the species might be due partly to conditions selected under study such as growth, extraction and harvesting time which were of significance in variations.

Relatively low antioxidant capacity of the extract compared to standards was attributed to the absence of *cis*-isomers of the compounds, inducing further studies that would encompass an understanding on the mechanism of isomerisation and its control for practical usage in order to maximize both the amount and effect of the targeted compound.

Conflict of Interest. The author declared that there is no conflict of interest.

Authorship Contributions. Concept: Y.A., O.K., I.T., Design: Y.A., O.K., I.T., Data Collection or Processing: Y.A., O.K., Analysis or Interpretation: Y.A., O.K., Literature Search: Y.A., O.K., I.T., Writing: Y.A., O.K., I.T.

Financial Disclosure. The authors gratefully thank Kırıkkale University Research Fund for their financial support, Project number 2019/054.

REFERENCES

- [1] Henríquez, V., Escobar, C., Galarza, J., Gimpel, J. (2016): Carotenoids in Microalgae (Chapter 8) in Carotenoids in Nature, Subcellular Biochemistry 79, C. Stange (ed.),

- Springer International Publishing Switzerland pp: 219-36. doi 10.1007/978-3-319-39126-7_8.
- [2] Grossman, A. R., Lohr, M., Im, C. S. (2004): *Chlamydomonas reinhardtii* in the Landscape of Pigments. *Annual Review of Genetics* 38: 119-173. doi: 10.1146/annurev.genet.38.072902.092328.
 - [3] Fiedor, J., Burda, K. (2014): Potential Role of Carotenoids as Antioxidants in Human Health and Disease. *Nutrients* 6: 466-488. doi: 10.3390/nu6020466.
 - [4] Manayi, A., Abdollahi, M., Raman, T., Nabavi, S. F., Habtemariam, S., Daglia, M., Nabavi, S. M. (2015): Lutein and Cataract: From Bench to Bedside. *Critical Reviews Biotechnology* 8551: 1-11. doi: 10.3109/07388551.2015.1049510.
 - [5] Gong, M., Bassi, A. (2016): Carotenoids from Microalgae: A Review of Recent Developments. *Biotechnology Advances* 34: 1396–1412. <http://dx.doi.org/10.1016/j.biotechadv.2016.10.005>.
 - [6] Patrick, L. (2000): Beta-Carotene: The Controversy Continues. *Journal Alternative Medicine Review* 5(6): 530-545. PMID: 11134976.
 - [7] Demming-Adams, B., Adams, W. W. (2002): Antioxidants in Photosynthesis and Human Nutrition. *Science* 298: 2149-2153. doi: 10.1126/science.1078002.
 - [8] Praveenkumar, R., Lee, K., Lee, J., Oh, Y-K. (2015): Breaking Dormancy: An Energy-Efficient Means of Recovering Astaxanthin from Microalgae. *Green Chemistry* 17: 1226-1234. <https://doi.org/10.1039/C4GC01413H>.
 - [9] Sun, Y., Wang, H., Guo, G., Pu, Y., Yan, B. (2014): The Isolation and Antioxidant Activity of Polysaccharides from the Marine Microalgae *Isochrysis galbana*. *Carbohydrate Polymers* 113: 22-31. <https://doi.org/10.1016/j.carbpol.2014.06.058>.
 - [10] Zhang, Y., Fang, H., Xie, Q., Sun, J., Liu, R., Hong, Z., Yi, R., Wu, H. (2014): Comparative Evaluation of the Radical-Scavenging Activities of Fucoxanthin and its Stereoisomers. *Molecules* 19(2): 2100-2113. <https://doi.org/10.3390/molecules19022100>.
 - [11] Wells, M. L., Potin, P., Craigie, J. S., Raven, J. A., Merchant, S. S., Helliwell, K. E., Smith, A. G., Camire, M. E., Brawley, S. H. (2017): Algae as Nutritional and Functional Food Sources: Revisiting our Understanding. *Journal of Applied Phycology* 29: 949-982. <https://doi.org/10.1007/s10811-016-0974-5>.
 - [12] Zhou, Y., Chen, Y., Li, M., Hu, C. (2020): Production of High-Quality Biofuel via Ethanol Liquefaction of Pretreated Natural Microalgae. *Renewable Energy* 147: 293-301. doi: 10.1007/978-3-319-32232-2.
 - [13] Herrero, M., Jaime, L., Martin-Alvarez, P. J., Cifuentes, A., Ibanez, E. (2006): Optimization of the Extraction of Antioxidants from *Dunaliella salina* Microalga by Pressurized Liquids. *Journal of Agricultural Food Chemistry* 54: 5597-5603. doi: 10.1021/jf060546q.
 - [14] Kyriakopoulou, K., Papadaki, S., Krokida, M. (2015): Life Cycle Analysis of β -Carotene Extraction Techniques. *Journal of Food Engineering* 167: 51-58. <https://doi.org/10.1016/j.jfoodeng.2015.03.008>.
 - [15] Sanchez, J. F., Fernandez, J. M., Acien, F. G., Ceron, M. C., Perez, J., & Molina, E. (2008): Biomass and Lutein Productivity of *Scenedesmus almeriensis*: Influence of Irradiance, Dilution Rate and Temperature. *Applied Microbiology and Biotechnology* 79: 719-729. doi: 10.1007/s00253-008-1494-2.
 - [16] Granado-Lorencio, F., Herrero-Barbudo, C., Acien-Fernandez, G., Molina-Grima, E., Fernandez-Sevilla, J. M., Perez-Sacristan, B., Blanco-Navarro, I. (2009): In Vitro Bioaccessibility of Lutein and Zeaxanthin from the Microalgae *Scenedesmus almeriensis*. *Food Chemistry* 114: 747-752. doi: 10.1016/j.foodchem.2008.10.058.
 - [17] Del Campo, J. A., Rodriguez, H., Moreno, J., Vargas, M. A., Rivas, J., Guerrero M. G. (2004): Accumulation of Astaxanthin and Lutein in *Chlorella zofingiensis* (Chlorophyta). *Applied Microbiology and Biotechnology* 64: 848-854. doi: 10.1007/s00253-003-1510-5.

- [18] Prommuak, C., Pavasant, P., Quitain, A. T., Goto, M., Shotipruk, A. (2013): Simultaneous Production of Biodiesel and Free Lutein from *Chlorella vulgaris*. *Chemical Engineering & Technology* 5: 733-739. <https://doi.org/10.1002/ceat.201200668>.
- [19] Aluç, Y., Başaran-Kankılıç, G., Tüzün, İ. (2018): Determination of Carotenoids in Two Algae Species from the Saline Water of Kapulukaya reservoir by HPLC. *Journal of Liquid Chromatography & Related Technologies* 41(2): 93-100. doi:10.1080/10826076.2017.1418376.
- [20] Kartal, M., Orhan, İ., Abu-Asaker, M., Şenol, F.S., Atıcı, T., Şener, B. (2009): Antioxidant and Anticholinesterase Assets and Liquid Chromatography-Mass Spectrometry Preface of Various Fresh-Water and Marine Macroalgae. *Pharmacognosy Magazine* 5: 291-297. DOI: 10.4103/0973-1296.58147.
- [21] Mendes-Pinto, M. M., Raposo, M. F. J., Bowen, J., Young, A. J., Morais, R. (2001): Evaluation of Different Cell Disruption Processes on Encysted Cells of *Haematococcus pluvialis*: Effects on Astaxanthin Recovery and Implications for Bio-availability. *Journal of Applied Phycology* 13: 19-24. <https://doi.org/10.1023/A:1008183429747>.
- [22] Kang, C. D., An, J. Y., Park, T. H., Sim, S. J. (2006): Astaxanthin Biosynthesis from Simultaneous N and P Uptake by the Green Alga *Haematococcus pluvialis* in Primary-Treated Wastewater. *Biochemical Engineering Journal* 31: 234-238. doi: 10.1016/j.bej.2006.08.002.
- [23] Li, H-B., Chen, F. (2001): Preparative Isolation and Purification of Astaxanthin from the Microalgae *Chlorococcum* sp. by High-Speed Counter-Current Chromatography. *Journal of Chromatography A* 952: 133-137. doi: 10.1016/s0021-9673(01)01022-6.
- [24] Durmaz, Y. (2007): Vitamin E (α -tocopherol) Production by the Marine Microalgae *Nanochloropsis oculata* (Eustigmatophyceae) in nitrogen limitation. *Aquaculture* 272: 717-722. doi:10.1016/j.aquaculture.2007.07.213.
- [25] Varela, J. C., Pereira, H., Vila, M., Leo'n, R. (2015): Production of Carotenoids by Microalgae: Achievements and Challenges. *Photosynthesis Research* 125: 423-436. doi: 10.1007/s11120-015-0149-2.
- [26] Patnaik, R., Mallick, N. (2015): Utilization of *Scenedesmus obliquus* Biomass as Feedstock for Biodiesel and other Industrially Important Co-products: An Integrated Paradigm for Microalgal Biorefinery. *Algal Research* 12: 328-336. doi: 10.1016/j.algal.2015.09.009.
- [27] Breuer, G., Martens, D. E., Draaisma, R. B., Wijffels, R. H., Lamers, P. P. (2015): Photosynthetic Efficiency and Carbon Partitioning in Nitrogen-Starved *Scenedesmus obliquus*. *Algal Research* 9: 254-262. <https://doi.org/10.1016/j.algal.2015.03.012>.
- [28] Bischoff, H. W., Bold, H. C. (1963): *Phycological Studies IV. Some Soil Algae from Enchanted Rock and Related Algal Species*. University of Texas Publication, Austin.
- [29] Richmond, A., Hu, Q. (2013): *Handbook of microalgal culture: Biotechnology and Applied Phycology* 2nd edition. Wiley Blackwell ISBN: 978-0-470-67389-8.
- [30] Salvador, N., Garreta, A. G., Lavelli, L., Ribera, M. (2007): Antimicrobial Activity of Iberian Macroalgae. *Scientia Marina* 71(1): 101-113. doi: <https://doi.org/10.3989/scimar.2007.71n1101>.
- [31] Blois, M. S. (1958): Antioxidant Determinations by the Use of a Stable Free Radical. *Nature* 26: 1199-1200. <https://doi.org/10.1038/1811199a0>.
- [32] Oyaizu, M. (1986): Studies on products of browning reaction: Antioxidative Activities of Products of Browning Reaction Prepared from Glucosamine. *Japan Journal of Nutrition* 44: 307-315. <https://doi.org/10.5264/eiyogakuzashi.44.307>.
- [33] Yuan, J. P., Chen, F., Liu, X., Li, X. Z. (2002): Carotenoid Composition in the Green Microalga *Chlorococcum*. *Food Chemistry* 76: 319-325. doi: 10.1016/s0308-8146(01)00279-5.
- [34] Inbaraj, B. S., Chien, J. T., Chen, B. H. (2006): Improved High Performance Liquid Chromatographic Method for Determination of Carotenoids in the Microalga *Chlorella*

- pyrenoidosa*. Journal of Chromatography A 1102: 193-199. doi: 10.1016/j.chroma.2005.10.055.
- [35] Liu, H. L., Kao, T. H., & Chen, B. H. (2004): Determination of Carotenoids in the Chinese Medical Herb Jiao-Gu-Lan (*Gynostemma pentaphyllum* MAKINO) by Liquid Chromatography. Chromatographia 60: 411-417. doi: 10.1016/s0021-9673(04)01406-2.
- [36] Chen, J. P., Tai, C. Y., Chen, B. H. (2004): Improved liquid chromatographic method for determination of carotenoids in Taiwanese Mango (*Mangifera indica* L.). Journal of Chromatography A 1054(1-2): 261-268.
- [37] Glass, A., Keogh, J., Meyskens, F., Valanis, B., Williams, J. et al. (1996). Effects of a Combination of Beta Carotene and Vitamin A on Lung Cancer and Cardiovascular Disease. New England Journal of Medicine 334(18): 1150-1155. doi: 10.1056/nejm199605023341802.
- [38] Jayappriyan, K. R., Rajkumar, R., Venkatakrishnan, V., Nagaraj, S., Rengasamy, R. (2013): In Vitro Anticancer Activity of Natural β -Carotene from *Dunaliella salina* EU5891199 in PC-3 Cells. Biomedicine & Preventive Nutrition 3(2): 99-105. doi: 10.1016/j.bionut.2012.08.003.
- [39] Ho, S-H., Chan, M-C., Liu, C-C., Chen, C-Y., Lee, W-L., Lee, D-J., Chang, J-S. (2014): Enhancing Lutein Productivity of an Indigenous Microalga *Scenedesmus Obliquus* FSP-3 Using Light-Related Strategies. Bioresource Technology 152: 275-282. doi: 10.1016/j.biortech.2013.11.031.
- [40] Goiris, K., Muylaert, K., Fraeye, I., Foubert, I., De Brabanter, J., De Cooman, L. (2012): Antioxidant Potential of Microalgae in Relation to Their Phenolic and Carotenoid Content. Journal of Applied Phycology 24: 1477-1486. doi: 10.1007/s10811-012-9804-6.
- [41] Cerón-García, M. C., González-López, C. V., Camacho-Rodríguez, J., López-Rosales L, García-Camacho F., Molina-Grima, E. (2018): Maximizing Carotenoid Extraction from Microalgae Used as Food Additives and Determined by Liquid Chromatography (HPLC). Food Chemistry 257: 316-324. doi: 10.1016/j.foodchem.2018.02.154.
- [42] Paliwal, C., Ghosh, T., George, B., Pancha, I., Maurya, R., Chokshi, K., Ghosh, A., Mishra, S. (2016): Microalgal Carotenoids: Potential Nutraceutical Compounds with Chemotaxonomic Importance. Algal Research 15: 24-31.
- [43] Aburai, N., Ohkubo, S., Miyashita, H., Abe, K. (2013): Composition of Carotenoids and Identification of Aerial Microalgae Isolated from the Surface of Rocks in Mountainous Districts of Japan. Algal Research 2: 237-243. doi:10.1016/j.algal.2016.01.017.
- [44] Guedes, A. C., Amaro, H. M., Malcata, F. X. (2011): Microalgae as Sources of Carotenoids. Marine Drugs 9: 625-644. <https://doi.org/10.3390/md9040625>.
- [45] Huang, D., Ou, B., Prior, R. L. (2005): The Chemistry Behind Antioxidant Capacity Assays. Journal of Agricultural and Food Chemistry 53: 1841-1856. <https://doi.org/10.1021/jf030723c>.
- [46] Apak, R., Özyürek, M., Güçlü, K., Çapanoğlu, E. (2016): Antioxidant Activity/Capacity Measurement. 1. Classification, Physicochemical Principles, Mechanisms and Electron Transfer (ET)-Based Assays. Journal of Agricultural and Food Chemistry 64: 997-1027.
- [47] Legendre, P, Legendre, L. (2012): Numerical Ecology. Elsevier (3rd edition), Pp: 1006. ISBN: 9780444538680.
- [48] Hu, C-C., Lin, J-T., Lu, F-J., Chou, F-P., Yang, D-J. (2008): Determination of Carotenoids in *Dunaliella Salina* Cultivated in Taiwan and Antioxidant Capacity of the Algal Carotenoid Extract. Food Chemistry 109: 439-446. doi: 10.1016/j.foodchem.2007.12.043.
- [49] Zhang, M., Li, R., Cao, L., Shi, J., Liu, H., Huang, Y., Shen, Q., (2014): Algal Sludge from Taihu Lake can be Utilized to Create Novel PGPR-Containing Bio-Organic Fertilizers. Journal of Environmental Management 132: 230-236. doi: 10.1016/j.jenvman.2013.10.031.
- [50] Yang, C., Zhang, H., Liu, R., Zhu, H., Zhang, L., Tsao, R. (2017): Bioaccessibility, Cellular Uptake, and Transport of Astaxanthin Isomers and Their Antioxidative Effects in Human

- Intestinal Epithelial Caco-2 Cells. *Journal of Agricultural and Food Chemistry* 65: 10223-10232. <https://doi.org/10.1021/acs.jafc.7b04254>.
- [51] Kopec, R. E., Failla, M. L. (2018): Recent Advances in the Bioaccessibility and Bioavailability of Carotenoids and Effects of Other Dietary Lipophiles. *Journal of Food Composition and Analysis* 68: 16-30. <https://doi.org/10.1016/j.jfca.2017.06.008>.
- [52] Honda, M., Kageyama, H., Hibino, T., Zhang, Y., Diono, W. (2019): Improved Carotenoid Processing With Sustainable Solvents Utilizing Z-Isomerization-Induced Alteration in Physicochemical Properties: A Review and Future Directions. *Molecules* 24: 2149, 1-19. Doi:10.3390/molecules24112149.
- [53] Guedes, A. C., Gião, M. S., Seabra, R., Silva Ferreira, A. C., Tamagnini, P., Moradas-Ferreira, P., Xavier Malcata, F. (2013): Evaluation of the Antioxidant Activity of Cell Extracts from Microalgae. *Marine Drugs* 11: 1256-1270. doi:10.3390/md11041256.
- [54] Jimenez, C., Pick, U. (1993): Differential Reactivity of β -Carotene Isomers from *Dunaliella Bardawil* toward Oxygen Radicals. *Plant Physiology* 101: 385-390. doi: 10.1104/pp.101.2.385.
- [55] Müller, L., Fröhlich, K., & Böhm, V. (2011): Comparative Antioxidant Activities of Carotenoids Measured by Ferric Reducing Antioxidant Power (FRAP), ABTS Bleaching Assay (α TEAC), DPPH Assay and Peroxyl Radical Scavenging Assay. *Food Chemistry* 129: 139-148. doi: 10.1016/j.foodchem.2011.04.045.