

Note on the Odd Perfect Numbers

Frank Vega

CopSonic, 1471 Route de Saint-Nauphary 82000 Montauban, France

Abstract

The Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. We state the conjecture that $\frac{\pi^2}{6.4} \times e^{0.0712132519795} \times \log x \geq e^\gamma \times \log(x - K \times \sqrt{x})$ is satisfied for infinitely many natural numbers $x > 10^8$ where $K > 0$ is a constant. Under the assumption of this conjecture and the Riemann Hypothesis, we prove that there is not any odd perfect number at all.

Keywords: Riemann Hypothesis, Prime numbers, Odd perfect numbers, Superabundant numbers, Sum-of-divisors function

2000 MSC: 11M26, 11A41, 11A25

1. Introduction

The Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. As usual $\sigma(n)$ is the sum-of-divisors function of n :

$$\sum_{d|n} d$$

where $d | n$ means the integer d divides n , $d \nmid n$ means the integer d does not divide n and $d^k || n$ means $d^k | n$ and $d^{k+1} \nmid n$. Define $f(n)$ and $G(n)$ to be $\frac{\sigma(n)}{n}$ and $\frac{f(n)}{\log \log n}$ respectively, such that \log is the natural logarithm. We know these properties from these functions:

Proposition 1.1. [1]. Let $\prod_{i=1}^r q_i^{a_i}$ be the representation of n as a product of primes $q_1 < \dots < q_r$ with natural numbers as exponents a_1, \dots, a_r . Then,

$$f(n) = \left(\prod_{i=1}^r \frac{q_i}{q_i - 1} \right) \times \prod_{i=1}^r \left(1 - \frac{1}{q_i^{a_i+1}} \right).$$

Proposition 1.2. For every prime power q^a , we have that $f(q^a) = \frac{q^{a+1}-1}{q^a \times (q-1)}$ [2]. If $m, n \geq 2$ are natural numbers, then $f(m \times n) \leq f(m) \times f(n)$ [2]. Moreover, if p is a prime number, and a, b two positive integers, then [2]:

$$f(p^{a+b}) - f(p^a) \times f(p^b) = -\frac{(p^a - 1) \times (p^b - 1)}{p^{a+b-1} \times (p - 1)^2}.$$

Email address: vega.frank@gmail.com (Frank Vega)

Preprint submitted to Elsevier

June 1, 2022

Say Robins(n) holds provided

$$G(n) < e^\gamma$$

where the constant $\gamma \approx 0.57721$ is the Euler-Mascheroni constant. The importance of this property is:

Proposition 1.3. Robins(n) holds for all natural numbers $n > 5040$ if and only if the Riemann Hypothesis is true [3].

The Chebyshev function $\theta(x)$ is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

with the sum extending over all prime numbers p that are less than or equal to x [4]. We state the following properties about this function:

Proposition 1.4. [4]. For $x \geq 89909$:

$$\theta(x) > \left(1 - \frac{0.068}{\log(x)}\right) \times x.$$

Proposition 1.5. [5]. There is a constant $K > 0$ such that there are infinitely many natural numbers x :

$$\theta(x) < x - K \times \sqrt{x}.$$

In mathematics, $\Psi = n \times \prod_{q|n} \left(1 + \frac{1}{q}\right)$ is called the Dedekind Ψ function. Say Dedekinds(q_n) holds provided

$$\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right) > \frac{e^\gamma}{\zeta(2)} \times \log \theta(q_n)$$

where q_n is the n th prime number, $\zeta(x)$ is the Riemann zeta function and $\zeta(2) = \prod_{i=1}^{\infty} \frac{q_i^2}{q_i^2 - 1} = \frac{\pi^2}{6}$. The importance of this inequality is:

Proposition 1.6. Dedekinds(q_n) holds for all prime numbers $q_n > 3$ if and only if the Riemann Hypothesis is true [6].

Let $q_1 = 2, q_2 = 3, \dots, q_k$ denote the first k consecutive primes, then an integer of the form $\prod_{i=1}^k q_i^{a_i}$ with $a_1 \geq a_2 \geq \dots \geq a_k \geq 0$ is called an Hardy-Ramanujan integer [7]. A natural number n is called superabundant precisely when, for all natural numbers $m < n$

$$f(m) < f(n).$$

Proposition 1.7. If n is superabundant, then n is an Hardy-Ramanujan integer [8]. Let n be a superabundant number, then $p \parallel n$ where p is the largest prime factor of n [8]. For large enough superabundant number n , we have that $q^{a_q} < 2^{a_2}$ for $q > 11$ where $q^{a_q} \parallel n$ and $2^{a_2} \parallel n$ [8]. For large enough superabundant number n , we obtain that $\log n < \left(1 + \frac{0.5}{\log p}\right) \times p$ where p is the largest prime factor of n [4]. Moreover, for large enough superabundant n , we know that $2^{a_2} < 2 \times p \times \log p$ such that p is the largest prime factor of n where $p \parallel n$ and $2^{a_2} \parallel n$ [8]. Let n be a superabundant number, then $f(n) > (1 - \varepsilon(p)) \times \prod_{q|n} \frac{q}{q-1}$ where $\varepsilon(p) = 1 - \frac{1}{\log p} \times \left(1 + \frac{1.5}{\log p}\right)$ and p is the largest prime factor of n [4].

On the sum of the reciprocals of power prime numbers not exceeding x , we have these results:

Proposition 1.8. [9]. For $x \geq 2278383$:

$$\sum_{p \leq x} \frac{1}{p} \geq \log \log x + B - \frac{1}{5 \times \log^3 x}$$

where $B \approx 0.261497212847642$ is the Meissel-Mertens constant [10].

Proposition 1.9. [11]. For $y \geq 10^8$:

$$\sum_{p \geq x} \frac{1}{p^2} \leq \frac{1}{y \times \log y} - \frac{1}{y \times \log^2 y} + \frac{2}{y \times \log^3 y} - \frac{2.07}{y \times \log^4 y}.$$

In addition, we will use these properties:

Proposition 1.10. [6]. For $n \geq 2$:

$$\prod_{q > q_n} \frac{q^2}{q^2 - 1} \leq e^{\frac{2}{q_n}}.$$

Proposition 1.11. [12]. For $x \geq 1$:

$$\frac{1}{x + 0.5} < \log\left(1 + \frac{1}{x}\right).$$

In number theory, a perfect number is a positive integer n such that $f(n) = 2$. Euclid proved that every even perfect number is of the form $2^{s-1} \times (2^s - 1)$ whenever $2^s - 1$ is prime. It is unknown whether any odd perfect numbers exist, though various results have been obtained:

Proposition 1.12. Any odd perfect number N must satisfy the following conditions: $N > 10^{1500}$ and the largest prime factor of N is greater than 10^8 [13], [14].

Say $\text{Vegas}(x)$ holds provided

$$\frac{\pi^2}{6.4} \times e^{0.0712132519795} \times \log x \geq e^y \times \log(x - K \times \sqrt{x})$$

where $K > 0$ is a constant.

Conjecture 1.13. $\text{Vegas}(x)$ holds for infinitely many natural numbers $x > 10^8$.

Under the assumption of this conjecture and the Riemann Hypothesis, we prove that there is not any odd perfect number at all.

2. Numerical Calculations

Theorem 2.1.

$$\sum_q \left(\frac{1}{q \times (q + 0.5)} \right) < 0.380503927189989469441$$

Proof. Using the Proposition 1.9, we check by computer that,

$$\begin{aligned} \sum_q \left(\frac{1}{q \times (q + 0.5)} \right) &< \sum_{q < 10^8} \left(\frac{1}{q \times (q + 0.5)} \right) + \sum_{q \geq 10^8} \left(\frac{1}{p^2} \right) \\ &< 0.380503926673572 + \frac{1}{10^8 \times \log 10^8} - \frac{1}{10^8 \times \log^2 10^8} + \frac{2}{10^8 \times \log^3 10^8} - \frac{2.07}{10^8 \times \log^4 10^8} \\ &< 0.380503927189989469441. \end{aligned}$$

□

3. Central Lemma

Lemma 3.1. *For all prime numbers $q_n > 10^8$, we have that*

$$\prod_{q \leq q_n} \left(1 + \frac{1}{q} \right) > e^{0.0712132519795} \times \log q_n$$

is satisfied.

Proof. We apply the logarithm to the both sides of the inequality,

$$\sum_{q \leq q_n} \log \left(1 + \frac{1}{q} \right) > 0.0712132519795 + \log \log q_n.$$

We use the Proposition 1.11,

$$\sum_{q \leq q_n} \frac{1}{q + 0.5} > 0.0712132519795 + \log \log q_n.$$

This is the same as

$$\sum_{q \leq q_n} \left(\frac{1}{q} \right) - \sum_{q \leq q_n} \left(\frac{1}{q} - \frac{1}{q + 0.5} \right) > 0.0712132519795 + \log \log q_n.$$

We know that

$$\frac{1}{q} - \frac{1}{q + 0.5} = \frac{1}{2 \times q \times (q + 0.5)}.$$

Hence,

$$\sum_{q \leq q_n} \left(\frac{1}{q} \right) - \log \log q_n > 0.0712132519795 + \sum_{q \leq q_n} \left(\frac{1}{2 \times q \times (q + 0.5)} \right).$$

We use that Proposition 1.8,

$$B - \frac{1}{5 \times \log^3(q_n)} > 0.0712132519795 + \sum_{q \leq q_n} \left(\frac{1}{2 \times q \times (q + 0.5)} \right)$$

that is equivalent to

$$B > 0.0712132519795 + \sum_{q \leq q_n} \left(\frac{1}{2 \times q \times (q + 0.5)} \right) + \frac{1}{5 \times \log^3(q_n)}.$$

Using the numerical computation in the Lemma 2.1, we only need to prove that

$$B > 0.0712132519795 + \frac{0.380503927189989469441}{2} + \frac{1}{5 \times \log^3(10^8)}$$

since $\frac{1}{5 \times \log^3(q_n)}$ decreases as q_n increases. In this way, we obtain that

$$B > 0.261497212847634$$

and thus, the proof is done. \square

4. Main Insight

Lemma 4.1. *Under the assumption of the Conjecture 1.13, we prove that*

$$\frac{\pi^2}{6.4} \times \prod_{q \leq q_n} \left(1 + \frac{1}{q}\right) > e^\gamma \times \log \theta(q_n)$$

is satisfied for infinitely many prime numbers $q_n > 10^8$.

Proof. We know there is a constant $K > 0$ such that there are infinitely many prime numbers $q_n > 10^8$:

$$\theta(q_n) < q_n - K \times \sqrt{q_n}$$

according to the Proposition 1.5. Hence, it is enough to show there are infinitely many prime numbers $q_n > 10^8$ such that

$$\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right) > \frac{e^\gamma}{\frac{\pi^2}{6.4}} \times \log (q_n - K \times \sqrt{q_n}).$$

The previous inequality will be satisfied when

$$e^{0.0712132519795} \times \log q_n \geq \frac{e^\gamma}{\frac{\pi^2}{6.4}} \times \log (q_n - K \times \sqrt{q_n})$$

due to the Lemma 3.1. That is equivalent to

$$\frac{\pi^2}{6.4} \times e^{0.0712132519795} \times \log q_n \geq e^\gamma \times \log (q_n - K \times \sqrt{q_n})$$

which is true for infinitely many prime numbers $q_n > 10^8$ under the assumption of the Conjecture 1.13. \square

5. Main Theorem

Theorem 5.1. *Under the assumption of the Conjecture 1.13 and the Riemann Hypothesis, we prove that there is not any odd perfect number at all.*

Proof. Suppose that N is the smallest odd perfect number, then we will show its existence implies that the Conjecture 1.13 or the Riemann Hypothesis is false. There is always a large enough superabundant number n such that n is a multiple of N . We would have

$$f(n) \leq f(N) \times f\left(\frac{n}{N}\right)$$

according to the Proposition 1.2. That is the same as

$$f(n) \leq 2 \times f\left(\frac{n}{N}\right)$$

since $f(N) = 2$, because N is a perfect number. Hence,

$$\begin{aligned} \frac{f(n)}{2} &= \frac{\left(2 - \frac{1}{2^{a_2}}\right) \times f\left(\frac{n}{2^{a_2}}\right)}{2} \\ &= f\left(\frac{n}{2^{a_2}}\right) \times \frac{\left(2 - \frac{1}{2^{a_2}}\right)}{2} \\ &= f\left(\frac{n}{2^{a_2}}\right) \times \frac{2^{a_2+1} - 1}{2^{a_2+1}} \end{aligned}$$

when $2^{a_2} \parallel n$ due to the Proposition 1.2. In this way, we have

$$\frac{f\left(\frac{n}{2^{a_2}}\right)}{f\left(\frac{n}{N}\right)} \leq \frac{2^{a_2+1}}{2^{a_2+1} - 1}.$$

However, we know that $p < 2^{a_2}$ because of $p > 10^8 > 11$ and the Propositions 1.7 and 1.12, where p is the largest prime factor of n . Consequently,

$$\frac{2^{a_2+1}}{2^{a_2+1} - 1} \leq \frac{2 \times p}{2 \times p - 1}$$

since $\frac{x}{x-1}$ decreases when $x \geq 2$ increases. In addition, we know that

$$\frac{2 \times p}{2 \times p - 1} \leq f(p)$$

where we know that $f(p) = \frac{p+1}{p}$ from the Proposition 1.2. Certainly,

$$\begin{aligned} 2 \times p^2 &\leq (p+1) \times (2 \times p - 1) \\ &= 2 \times p^2 + 2 \times p - p - 1 \\ &= 2 \times p^2 + p - 1 \end{aligned}$$

where this inequality is satisfied for every prime number p . So,

$$\frac{f\left(\frac{n}{2^{a_2}}\right)}{f\left(\frac{n}{N}\right)} \leq f(p)$$

where we know that $p \parallel n$ from the Proposition 1.7. Under the assumption of the Riemann Hypothesis, we have that

$$\begin{aligned} e^\gamma &> G(n) \\ &= \frac{f\left(\frac{n}{p}\right) \times f(p)}{\log \log n} \\ &\geq \frac{f\left(\frac{n}{p}\right) \times f\left(\frac{n}{2a_2}\right)}{f\left(\frac{n}{N}\right) \times \log \log n} \end{aligned}$$

since $f(\dots)$ is multiplicative and as a consequence of the Propositions 1.3. This is equivalent to

$$\frac{f\left(\frac{n}{p}\right)}{f\left(\frac{n}{N}\right)} < \frac{e^\gamma}{f\left(\frac{n}{2a_2}\right)} \times \log \log n.$$

Under the assumption of the Conjecture 1.13 and using the Lemma 4.1 and the Proposition 1.12:

$$\frac{\pi^2}{8} \times \prod_{q \leq p} \left(1 + \frac{1}{q}\right) > e^\gamma \times \log((\theta(p))^{0.8}).$$

From the Propositions 1.1 and 1.7, we know that

$$f\left(\frac{n}{2a_2}\right) = \left(\prod_{i=2}^k \frac{q_i}{q_i - 1}\right) \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right)$$

where $q_k = p$ and $q_1 = 2$. We know that

$$\frac{q_i}{q_i - 1} = \frac{q_i + 1}{q_i} \times \frac{q_i^2}{q_i^2 - 1}.$$

Using the previous inequality and the Conjecture 1.13, we obtain that

$$\begin{aligned} e^\gamma \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) \times \log((\theta(p))^{0.8}) &< \frac{\pi^2}{8} \times \prod_{q \leq p} \left(1 + \frac{1}{q}\right) \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) \\ &= f\left(\frac{n}{2a_2}\right) \times \frac{3}{2} \times \prod_{q > p} \frac{q^2}{q^2 - 1} \\ &\leq f\left(\frac{n}{2a_2}\right) \times \frac{3}{2} \times e^{\frac{2}{p}} \end{aligned}$$

according to the Proposition 1.10. Taking into account that $p > 10^8 > 3$ and n is superabundant:

$$\frac{\frac{3}{2} \times e^{\frac{2}{p}}}{\log((\theta(p))^{0.8})} > \frac{e^\gamma}{f\left(\frac{n}{2a_2}\right)} \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right).$$

We use the previous inequality to show that

$$\frac{f\left(\frac{n}{p}\right)}{f\left(\frac{n}{N}\right)} \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) < \frac{\frac{3}{2} \times e^{\frac{2}{p}}}{\log((\theta(p))^{0.8})} \times \log \log n.$$

For large enough superabundant number n and $p > 10^8$, then

$$\frac{\frac{3}{2} \times e^{\frac{2}{p}}}{\log((\theta(p))^{0.8})} \times \log \log n \leq \frac{\frac{3}{2} \times e^{\frac{2}{10^8}}}{\log\left(\left(1 - \frac{0.068}{\log 10^8}\right) \times 10^8\right)^{0.8}} \times \log\left(\left(1 + \frac{0.5}{\log 10^8}\right) \times 10^8\right)$$

because of the Propositions 1.4 and 1.7. We obtain that

$$\frac{\frac{3}{2} \times e^{\frac{2}{10^8}}}{\log\left(\left(1 - \frac{0.068}{\log 10^8}\right) \times 10^8\right)^{0.8}} \times \log\left(\left(1 + \frac{0.5}{\log 10^8}\right) \times 10^8\right) < 1.87811.$$

Thus,

$$\frac{f\left(\frac{n}{p}\right)}{f\left(\frac{n}{N}\right)} \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) < 1.87811.$$

For every prime p_i that divides N such that $p_i^{a_i} \parallel N$ and $p_i^{a_i+b_i} \parallel n$ for a_i, b_i two natural numbers, we have that

$$f(p_i^{a_i+b_i}) - f(p_i^{a_i}) \times f(p_i^{b_i}) = -\frac{(p_i^{a_i} - 1) \times (p_i^{b_i} - 1)}{p_i^{a_i+b_i-1} \times (p_i - 1)^2}$$

in the Proposition 1.2. This is equal to

$$\frac{f(p_i^{a_i+b_i})}{f(p_i^{b_i})} = f(p_i^{a_i}) - \frac{(p_i^{a_i} - 1) \times (p_i^{b_i} - 1)}{f(p_i^{b_i}) \times p_i^{a_i+b_i-1} \times (p_i - 1)^2}.$$

Hence,

$$\begin{aligned} \frac{f\left(\frac{n}{p}\right)}{f\left(\frac{n}{N}\right)} \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) &= \prod_i \left(\frac{f(p_i^{a_i+b_i})}{f(p_i^{b_i})}\right) \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) \\ &= \prod_i \left(f(p_i^{a_i}) - \frac{(p_i^{a_i} - 1) \times (p_i^{b_i} - 1)}{f(p_i^{b_i}) \times p_i^{a_i+b_i-1} \times (p_i - 1)^2}\right) \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) \\ &\approx \prod_i (f(p_i^{a_i})) \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) \\ &= f(N) \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) \\ &= 2 \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) \\ &> 2 \times \left(1 - \frac{1}{\log p} \times \left(1 + \frac{1.5}{\log p}\right) - \log\left(1 - \frac{1}{2^{a_2+1}}\right)\right) \\ &> 2 \times \left(1 - \frac{1}{\log p} \times \left(1 + \frac{1.5}{\log p}\right) - \log\left(1 - \frac{1}{4 \times p \times \log p}\right)\right) \\ &> 2 \times \left(1 - \frac{1}{\log 10^8} \times \left(1 + \frac{1.5}{\log 10^8}\right) - \log\left(1 - \frac{1}{4 \times 10^8 \times \log 10^8}\right)\right) \\ &> 1.88 \\ &> 1.87811 \end{aligned}$$

using the Propositions 1.7 and 1.1 since we know that the expression

$$\frac{(p_i^{a_i} - 1) \times (p_i^{b_i} - 1)}{f(p_i^{b_i}) \times p_i^{a_i+b_i-1} \times (p_i - 1)^2}$$

tends to 0 as b_i tends to infinity for every odd prime p . Certainly, the fraction $\frac{f(\frac{n}{p})}{f(\frac{n}{N})}$ gets closer to 2 as long as we take n bigger and bigger. However,

$$1.87811 < \frac{f(\frac{n}{p})}{f(\frac{n}{N})} \times \prod_{i=2}^k \left(1 - \frac{1}{q_i^{a_i+1}}\right) < 1.87811$$

is a contradiction. By contraposition, the number N does not exist under the assumption of the Conjecture 1.13 and the Riemann Hypothesis. The smallest counterexample N must comply that $N > 10^{1500}$ and therefore, we will always be capable of obtaining a large enough superabundant number n that is multiple of N . Note that, this proof fails for even perfect numbers. \square

Acknowledgments

The author would like to thank his mother, maternal brother and his friend Sonia for their support.

References

- [1] A. Hertlein, Robin's Inequality for New Families of Integers, *Integers* 18, (2018).
- [2] R. Vojak, On numbers satisfying Robin's inequality, properties of the next counterexample and improved specific bounds, arXiv preprint arXiv:2005.09307(2020).
- [3] G. Robin, Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann, *J. Math. pures appl* 63 (2) (1984) 187–213.
- [4] S. Nazardonyavi, S. Yakubovich, Superabundant numbers, their subsequences and the Riemann hypothesis, arXiv preprint arXiv:1211.2147(2012).
- [5] E. Schmidt, Über die Anzahl der Primzahlen unter gegebener Grenze, *Mathematische Annalen* 57 (2) (1903) 195–204.
- [6] P. Solé, M. Planat, Extreme values of the Dedekind ψ function, *Journal of Combinatorics and Number Theory* 3 (1) (2011) 33–38.
- [7] Y. Choie, N. Lichiardopol, P. Moree, P. Solé, On Robin's criterion for the Riemann hypothesis, *Journal de Théorie des Nombres de Bordeaux* 19 (2) (2007) 357–372. doi:doi:10.5802/jtnb.591.
- [8] L. Alaoglu, P. Erdős, On highly composite and similar numbers, *Transactions of the American Mathematical Society* 56 (3) (1944) 448–469. doi:doi:10.2307/1990319.
- [9] C. Axler, New Estimates for Some Functions Defined Over Primes, *Integers* 18, (2018).
- [10] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie., *J. reine angew. Math.* 1874 (78) (1874) 46–62. doi:10.1515/crll.1874.78.46.
- [11] J.-L. Nicolas, The sum of divisors function and the Riemann hypothesis, *The Ramanujan Journal* (2021) 1–45.
- [12] A. Ghosh, An asymptotic formula for the Chebyshev theta function, *Notes on Number Theory and Discrete Mathematics* 25 (4) (2019) 1–7. doi:10.7546/nntdm.2019.25.4.1-7.
- [13] P. Ochem, M. Rao, Odd perfect numbers are greater than 10^{1500} , *Mathematics of Computation* 81 (279) (2012) 1869–1877. doi:doi:10.1090/S0025-5718-2012-02563-4.
- [14] T. Goto, Y. Ohno, Odd perfect numbers have a prime factor exceeding 10^8 , *Mathematics of Computation* 77 (263) (2008) 1859–1868.