

Modulation of immune responses by nutritional ligands of Aryl hydrocarbon Receptor

Alba De Juan¹, Elodie Segura^{1*}

¹ Institut Curie, PSL Research University, INSERM, U932, 26 rue d'Ulm, 75005 Paris, France

* Correspondence:

Elodie Segura
elodie.segura@curie.fr

Keywords: aryl hydrocarbon receptor, AHR, tryptophan, immunity, microbiota, intestine

Abstract

Accumulating evidence indicates that nutrition can modulate the immune system through metabolites, either produced by host digestion or by microbiota metabolism. In this review, we focus on dietary metabolites that are agonists of the Aryl hydrocarbon Receptor (AhR). AhR is a ligand-activated transcription factor, initially characterized for its interaction with xenobiotic pollutants. Numerous studies have shown that AhR also recognizes indoles and tryptophan catabolites originating from dietary compounds and commensal bacteria. Here, we review recent work employing diet manipulation to address the impact of nutritional AhR agonists on immune responses, both locally in the intestine and at distant sites. In particular, we examine the physiological role of these metabolites in immune cell development and functions (including T lymphocytes, innate-like lymphoid cells, and mononuclear phagocytes) and their effect in inflammatory disorders.

1 Introduction

Food represents not only a source of nutrients for the maintenance of essential biological functions, but also contains dietary components that regulate immune cell populations. These include microbiota-derived short-chain fatty acids, polyamines, and indoles derivatives, which are ligands of the Aryl Hydrocarbon Receptor (AhR) (1).

AhR is a ligand-activated transcription factor residing in the cytosol. Upon binding to an agonist, AhR translocates to the nucleus where it forms an active heterodimer with ARNT and promotes the transcription of genes that are under its control. AhR is expressed in multiple immune cells such as myeloid cells, innate lymphoid cells, B lymphocytes and certain subtypes of T cells. AhR activation has an overall anti-inflammatory and immunoregulatory role in innate and adaptive immunity, both in steady-state or in inflammatory scenarios such as autoimmunity or infection (2). However, a number of these observations were made *in vitro* or *in vivo* by injecting AhR agonists in non-physiological routes or concentrations. Some studies have also used AhR ligands of xenobiotic origin, which are known to induce aberrant AhR signaling (3). This has led to some contradictory findings, in particular in T cell biology (4).

In this review, we examine the physiological role of nutritional AhR ligands in immune cells and immune responses by focusing on experimental results obtained by direct intestinal exposure or diet manipulation.

2 What are the nutritional AhR ligands ?

AhR was initially described as a receptor for xenobiotic pollutants, mostly aromatic hydrocarbons. However, over the years, physiological ligands have been identified. There are other exhaustive reviews of AhR agonists (5–7), here we focus on nutritional AhR ligands (Figure 1 and Table 1).

AhR ligands are present in the diet, as natural compounds in food. Flavonoids derived from fruits and vegetables such as quercetin and resveratrol have been identified as AhR ligands based on *in vitro* assays, but their physiological relevance is unclear (8–10). The main class of dietary AhR ligands is indoles, including Indole-3-acetonitrile, Indole-3-carbinole (I3C), 3,3'-diindolylmethane (DIM) and Indolo(3,4)carbazole, which are found mainly in cruciferous vegetables like broccoli or Brussel sprouts (11). After consumption, I3C is converted in the stomach by acid-mediated condensation into various byproducts, including high affinity AhR ligands DIM and indole[3,2-b]carbazole (ICZ) (11).

Another source of nutritional AhR ligands is microbiota metabolism, particularly tryptophan catabolism. Some species of bacteria such as *Lactobacillus* can use tryptophan instead of glucose as a source of energy, and produce AhR ligands such as indole-3-acetic acid (IAA), tryptamine (TA) and 3-methyl indole (12,13). A well-described example is *Lactobacillus reuteri*, producing the AhR ligand indole-3-aldehyde (IAld) (13). Moreover, Tryptophanase-expressing bacteria, which are mostly ampicillin-sensitive and vancomycin-resistant, degrade tryptophan into indole that is further metabolized by host liver cells into AhR ligands, such as indoxyl-3-sulfate (I3S) and indole-3-propionic acid (IPA) (14). In addition, it has been reported that butyrate and other short-chain fatty acids (SCFA), which originate from the fermentation of dietary fibers by the microbiota, can activate AhR signaling in reporter cell lines (15). Whether this observation holds true in other cell types remains to be confirmed. In particular, another study failed to detect AhR activation in B cells upon *in vitro* exposure to butyrate, while 5-hydroxyindole-3-acetic acid (5-HIAA), a serotonin metabolite, activated AhR signaling in B cells *in vitro* and *in vivo* (16).

3 What is the biodistribution of nutritional AhR ligands ?

Dietary AhR ligands are released locally in the digestive track but also distributed to other sites via the blood. After oral administration, the biodistribution of I3C and its acid condensation products has been analyzed by high-performance liquid chromatography. I3C is absorbed from the gut and distributed systemically into a number of well-perfused tissues (17). I3C level peaks at 15min and decreases considerably 1h after administration in the plasma, liver, kidney, lung, heart and brain. However, I3C products DIM, LTR₁ and HI-IM are detectable in these organs from 15 min and persist after 6h, and up to 24h in the liver. ICZ, another product from I3C, was also identified in the liver, but not in plasma or other organs, at 6h and 24h after I3C administration (17). In line with those observations, I3C was detected in serum 15 min after gavage, but not when administered as supplement in the chow diet. By contrast, DIM was detectable in the serum of mice fed with I3C-supplemented chow diet (18). In a study carried out in women given a single oral I3C dose, DIM, but not I3C itself, was detected in plasma peaking at 2h and returning to basal levels 24h after administration (19). These observations suggest that I3C is rapidly cleared from the circulation, while its condensation products can reach distant organs and exert longer lasting effects.

It has also been proposed that nutritional AhR ligands can cross the blood-brain barrier. After intra-peritoneal injection, I3S was detected in the brain (20), however oral administration was not examined.

Finally, microbiota-derived AhR ligands have been detected in breastmilk. After gavage of pregnant mice with radiolabelled *E.Coli*, a species which expresses Tryptophanase, labelled AhR ligands were detected in maternal milk, including indole-3-lactic acid, showing the transfer from maternal intestinal microbiota to milk (21).

4 Do nutritional AhR ligands influence microbiota composition?

Tryptophan content in the diet shapes significantly the microbiota composition. Tryptophan supply relies exclusively on the diet since the host cannot synthesize it. After 2 or 3 weeks of dietary intervention, mice fed with a tryptophan-deprived diet display an increase in fecal *Actinobacteria* and *Proteobacteria*, and lower relative abundance of *Bacteroidetes* and bacteria belonging to *Firmicutes* phylum such as *Lactobacillus* and *Staphylococcus* (22,23). In addition, after 4 weeks of tryptophan-low diet, the abundance of *Lactobacillus reuteri* is also decreased in the stomach (13). Of note, tryptophan is metabolised in the gut not only into AhR ligands by microbiota, but also through the serotonin and kynurenine pathways by host cells (5,24). Whether the observed alterations in microbiota composition are entirely dependent on changes in AhR ligands availability remains to be confirmed. In addition, caution should be exercised when interpreting *in vivo* experiments employing tryptophan-low diets, as modification in microbiota diversity by itself may impact the outcome.

By contrast, I3C content in the diet only causes a relatively minor change in intestinal microbiota composition. Normal chow contains phytochemicals that can act as precursors of AhR ligands. Switching from normal chow to a synthetic diet alters fecal microbiota diversity (25,26). Mice fed with a AhR ligand-free synthetic diet show a decreased abundance in fecal *Bacteroidetes* and increased abundance in *Actinobacteria* and *Firmicutes* compared to mice fed with the same diet supplemented with I3C (25,26). Relative change observed in *Erysipelotrichaceae* is inconsistent between studies (25,26). Importantly, most of these alterations in fecal microbiome composition were also observed in AhR-deficient mice, showing that this occurs independently of AhR signaling (26). In a study employing a synthetic diet supplemented with DIM, no difference were observed in fecal microbiota between groups (27). However, mice fed with the synthetic diet had increased *Bacteroidetes* and decreased *Firmicutes* abundance in the small intestine compared to mice fed with the DIM-supplemented synthetic diet. These differences in microbiota diversity induced by DIM supplementation were abrogated in AhR-deficient mice (27).

These results highlight a complex interplay between microbiota composition and the supply of dietary AhR ligands, with tryptophan having a more pronounced effect than single AhR ligands.

5 What is the impact of nutritional AhR ligands on intestinal immunity ?

Nutritional AhR ligands are essential for the maintenance of intestinal intraepithelial lymphocytes (IELs) and type 3 innate lymphoid cells (ILC3).

Small intestine IELs are a specialized population of T cells composed of several subsets (TCR $\gamma\delta$, CD4⁻CD8 $\alpha\alpha$ ⁺ TCR $\alpha\beta$ and CD4⁺CD8 $\alpha\alpha$ ⁺ TCR $\alpha\beta$). Mice fed with a AhR ligand-free synthetic diet have lower numbers of TCR $\gamma\delta$ and CD4⁻CD8 $\alpha\alpha$ ⁺ TCR $\alpha\beta$ IELs compared to mice on I3C-supplemented synthetic diet (28,29). However, these IELs can develop normally in AhR-deficient, indicating that dietary AhR ligands are required for their maintenance rather than differentiation (29). In addition, CD4⁺CD8 $\alpha\alpha$ ⁺ TCR $\alpha\beta$ IELs are decreased in the intestine of mice fed with a tryptophan-low synthetic diet compared to standard diet and tryptophan-high diet (30). *Lactobacillus reuteri* was identified as essential for CD4⁺CD8 $\alpha\alpha$ ⁺ TCR $\alpha\beta$ IELs development. However, a tryptophan-high diet in conjunction

with *L.reuteri* colonization was not sufficient to induce these IELs in germ-free mice, suggesting the participation of additional factors (30).

ILC3 produce lymphotoxin, controlling the development of intestinal lymphoid follicles. Mice fed with a AhR ligand-free synthetic diet display decreased ILC3 and intestinal lymphoid follicles numbers compared to mice fed on I3C-supplemented synthetic diet (28,31). Of note, I3C supplementation had no impact when given to AhR-deficient mice (31). In addition, development of ILC3 and intestinal lymphoid follicles are normal in germ-free mice (32), suggesting that dietary AhR ligands, produced independently of microbiota metabolism, are sufficient for ILC3 differentiation or maintenance. ILC3 are the main producers of intestinal IL22, which is critical for the secretion of antimicrobial peptides by intestinal epithelial cells and the defense against intestinal infections (33). Consistent with decreased ILC3 numbers, mice fed on a AhR ligand-free synthetic diet express lower levels in the intestine of antimicrobial peptides such as C-type lectin regenerating islet-derived protein 3 (RegIII) (29,34), and are more susceptible to infections with *Citrobacter Rodentium* (35) or *Clostridium difficile* (28).

AhR has been proposed to control Treg differentiation. Tregs in the gut express higher levels of AhR than other Tregs in the body (36). AhR expression is necessary for Treg gut homing and function, and is independent of microbiota, since it is not affected in antibiotics-treated mice or germ-free mice. Whether nutritional AhR ligands play a role is unclear, as feeding mice with an AhR ligand-free synthetic diet was reported to be inconclusive regarding Treg development (36).

Lack of dietary AhR ligands worsens the symptoms of intestinal inflammation. In the model of DSS-induced colitis, mice on a synthetic AhR ligand-free diet show more severe symptoms, such as weight loss and tissue damage, compared to mice fed with a synthetic diet supplemented with I3C (26,29) or with tryptophan (34). Tryptophan supplementation did not modify the severity of symptoms in AhR-deficient mice, confirming the dependency on AhR activation (34). Consistent with these observations, symptoms of DSS-induced colitis were mildly ameliorated when mice fed on normal chow were given tryptophan supplementation in the drinking water (37) or I3C by oral gavage (38). In mice fed on AhR ligand-free synthetic diet, increased epithelial damage may be due to the lack of IL22 production in response to DSS-induced inflammation (26,34). Importantly, AhR activation in intestinal epithelial cells is also involved in barrier repair during colitis (39).

Finally, in a model of oral tolerance to ovalbumin, mice fed with I3C-enriched chow diet have lower levels of serum anti-ovalbumin IgG1 antibodies, indicating better induction of oral tolerance (18). This was correlated with increased expression in the small intestine of retinaldehyde dehydrogenase, a molecule known to promote Treg differentiation, but the target cells of dietary AhR ligands in this model remain unclear.

Collectively, these observations show an essential role for nutritional AhR ligands in maintaining intestinal lymphoid populations and homeostasis (Figure 2).

6 What is the impact of nutritional AHR ligands on immune responses at distant sites ?

Nutritional AhR ligands can also modulate the differentiation of immune cells outside of the intestinal mucosa, as shown for monocytes. Monocytes circulate in the blood and are recruited to tissues where they differentiate into dendritic cells or macrophages. In mice fed with a synthetic AhR ligand-free diet, monocyte differentiation into dendritic cells is reduced in the skin, compared to mice on an I3C-supplemented diet (40). This is consistent with *in vitro* observations that AhR activation skews monocyte differentiation from macrophages to dendritic cells by controlling the expression of

the transcription factors Irf4 and Blimp-1. In addition, monocyte differentiation into dendritic cells in the peritoneum is impaired in antibiotics-treated mice and could be restored by I3C diet supplementation (40), suggesting a role for microbiota-derived AhR ligands in physiological conditions (Figure 3).

Deficit in dietary AhR ligands increases the severity of inflammation in the central nervous system, as evidenced in the experimental autoimmune encephalomyelitis (EAE) model. After induction of EAE, mice placed on a synthetic tryptophan-free diet display worse disease scores and delayed recovery compared to mice fed with tryptophan-supplemented diet (20,41). This difference was abolished in mice deficient for AhR specifically in astrocytes or in microglia, showing that dietary AhR ligands can exert their effect on brain-resident populations. Lack of dietary tryptophan results in the increased expression of pro-inflammatory molecules in the brain such as *Ccl2*, *Nos2* and *Tnfa* (20,41). Treatment with ampicillin also delays disease recovery and increases *Ccl2* and *Nos2* expression, which could be reverted by diet supplementation with IPA, IAld, indole or Tryptophanase (20), suggesting a major role in this phenomenon for microbiota-derived AhR ligands.

AhR ligands from microbiota metabolism also influence the differentiation of IL-10 producing regulatory B cells, which are found in lymphoid organs. Gavage with 5-HIAA increases the expression of *Il10* in spleen B cells and reduces the severity of joint swelling in a model of antigen-induced arthritis (16). This effect is abolished in mice deficient for AhR in B cells, confirming the role of AhR signaling.

These observations indicate that AhR ligands participate in the communication between gut, microbiota and distant tissues such as brain and skin.

7 Conclusion and Perspectives

There is accumulating evidence that nutritional AhR ligands play an essential role in the maintenance of intestinal immune homeostasis and the control of intestinal inflammation. Several studies also suggest a similar role in distant tissues such as skin and brain. Whether dietary AhR ligands impact other organs or mucosal sites remains to be investigated.

Circadian rhythms regulate some essential aspects of immune activity, such as leukocyte trafficking or inflammatory cytokine secretion (42). AhR has been reported to interact with circadian clock proteins and suppress their transcriptional activity (43). In addition, the supply of nutritional AhR ligands fluctuates across time with feeding behavior, as does microbiota mass and the release of nutritional metabolites (44). It will be important to decipher the possible connection between AhR, circadian rhythms and immune cells.

Impaired production of AhR ligands has been observed in the intestinal microbiota of patients suffering from inflammatory bowel disease (45) and celiac disease (46). Given the critical impact of dietary AhR ligands on ILC3 numbers and the maintenance of barrier integrity, diet supplementation with AhR ligands, or AhR ligand-producing bacteria, is an attractive strategy to improve the treatment of inflammatory gastrointestinal diseases (24). However, a better understanding of the role of nutritional AhR ligands on the immune homeostasis of distant tissues and on myeloid cells will be essential to optimize these therapeutic approaches.

Conflict of Interest

The authors declare the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

ADJ and ES wrote the manuscript.

Funding

This work was funded by INSERM, Institut Curie, Agence Nationale de la Recherche (ANR-10-LABX-0043 and ANR-17-CE15-0011-01) and European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 842535.

References

1. Postler TS, Ghosh S. Understanding the holobiont: how microbial metabolites affect human health and shape the immune system. *Cell Metab* (2017) **26**:110–130. doi:10.1016/j.cmet.2017.05.008
2. Rothhammer V, Quintana FJ. The aryl hydrocarbon receptor: an environmental sensor integrating immune responses in health and disease. *Nat Rev Immunol* (2019) **19**:184–197. doi:10.1038/s41577-019-0125-8
3. Stockinger B, Di Meglio P, Gialitakis M, Duarte JH. The aryl hydrocarbon receptor: multitasking in the immune system. *Annu Rev Immunol* (2014) **32**:403–432. doi:10.1146/annurev-immunol-032713-120245
4. Zhou L. AHR function in lymphocytes: emerging concepts. *Trends Immunol* (2016) **37**:17–31. doi:10.1016/j.it.2015.11.007
5. Hubbard TD, Murray IA, Perdew GH. Indole and tryptophan metabolism: endogenous and dietary routes to ah receptor activation. *Drug Metab Dispos* (2015) **43**:1522–1535. doi:10.1124/dmd.115.064246
6. Nguyen LP, Bradfield CA. The search for endogenous activators of the aryl hydrocarbon receptor. *Chem Res Toxicol* (2008) **21**:102–116. doi:10.1021/tx7001965
7. Dolciemi D, Ballarotto M, Gargaro M, López-Cara LC, Fallarino F, Macchiarulo A. Targeting Aryl hydrocarbon receptor for next-generation immunotherapies: Selective modulators (SAhRMs) versus rapidly metabolized ligands (RMAhRLs). *Eur J Med Chem* (2020) **185**:111842. doi:10.1016/j.ejmech.2019.111842
8. Ciolino HP, Daschner PJ, Yeh GC. Resveratrol inhibits transcription of CYP1A1 in vitro by preventing activation of the aryl hydrocarbon receptor. *Cancer Res* (1998) **58**:5707–5712.
9. Ciolino HP, Daschner PJ, Yeh GC. Dietary flavonols quercetin and kaempferol are ligands of the aryl hydrocarbon receptor that affect CYP1A1 transcription differentially. *Biochem J* (1999) **340** (Pt 3):715–722.
10. Jin U-H, Park H, Li X, Davidson LA, Allred C, Patil B, Jayaprakasha G, Orr AA, Mao L, Chapkin RS, et al. Structure-Dependent Modulation of Aryl Hydrocarbon Receptor-Mediated

Activities by Flavonoids. *Toxicol Sci* (2018) **164**:205–217. doi:10.1093/toxsci/kfy075

11. Bjeldanes LF, Kim JY, Grose KR, Bartholomew JC, Bradfield CA. Aromatic hydrocarbon responsiveness-receptor agonists generated from indole-3-carbinol in vitro and in vivo: comparisons with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Proc Natl Acad Sci USA* (1991) **88**:9543–9547. doi:10.1073/pnas.88.21.9543
12. Jin U-H, Lee S-O, Sridharan G, Lee K, Davidson LA, Jayaraman A, Chapkin RS, Alaniz R, Safe S. Microbiome-derived tryptophan metabolites and their aryl hydrocarbon receptor-dependent agonist and antagonist activities. *Mol Pharmacol* (2014) **85**:777–788. doi:10.1124/mol.113.091165
13. Zelante T, Iannitti RG, Cunha C, De Luca A, Giovannini G, Pieraccini G, Zecchi R, D'Angelo C, Massi-Benedetti C, Fallarino F, et al. Tryptophan catabolites from microbiota engage aryl hydrocarbon receptor and balance mucosal reactivity via interleukin-22. *Immunity* (2013) **39**:372–385. doi:10.1016/j.immuni.2013.08.003
14. Wikoff WR, Anfora AT, Liu J, Schultz PG, Lesley SA, Peters EC, Siuzdak G. Metabolomics analysis reveals large effects of gut microflora on mammalian blood metabolites. *Proc Natl Acad Sci USA* (2009) **106**:3698–3703. doi:10.1073/pnas.0812874106
15. Marinelli L, Martin-Gallausiaux C, Bourhis J-M, Béguet-Crespel F, Blottière HM, Lapaque N. Identification of the novel role of butyrate as AhR ligand in human intestinal epithelial cells. *Sci Rep* (2019) **9**:643. doi:10.1038/s41598-018-37019-2
16. Rosser EC, Piper CJM, Matei DE, Blair PA, Rendeiro AF, Orford M, Alber DG, Krausgruber T, Catalan D, Klein N, et al. Microbiota-Derived Metabolites Suppress Arthritis by Amplifying Aryl-Hydrocarbon Receptor Activation in Regulatory B Cells. *Cell Metab* (2020) **31**:837–851.e10. doi:10.1016/j.cmet.2020.03.003
17. Anderton MJ, Manson MM, Verschoyle RD, Gescher A, Lamb JH, Farmer PB, Steward WP, Williams ML. Pharmacokinetics and tissue disposition of indole-3-carbinol and its acid condensation products after oral administration to mice. *Clin Cancer Res* (2004) **10**:5233–5241. doi:10.1158/1078-0432.CCR-04-0163
18. Hammerschmidt-Kamper C, Biljes D, Merches K, Steiner I, Daldrup T, Bol-Schoenmakers M, Pieters RHH, Esser C. Indole-3-carbinol, a plant nutrient and AhR-Ligand precursor, supports oral tolerance against OVA and improves peanut allergy symptoms in mice. *PLoS One* (2017) **12**:e0180321. doi:10.1371/journal.pone.0180321
19. Reed GA, Arneson DW, Putnam WC, Smith HJ, Gray JC, Sullivan DK, Mayo MS, Crowell JA, Hurwitz A. Single-dose and multiple-dose administration of indole-3-carbinol to women: pharmacokinetics based on 3,3'-diindolylmethane. *Cancer Epidemiol Biomarkers Prev* (2006) **15**:2477–2481. doi:10.1158/1055-9965.EPI-06-0396
20. Rothhammer V, Mascanfroni ID, Bunse L, Takenaka MC, Kenison JE, Mayo L, Chao C-C, Patel B, Yan R, Blain M, et al. Type I interferons and microbial metabolites of tryptophan modulate astrocyte activity and central nervous system inflammation via the aryl hydrocarbon receptor. *Nat Med* (2016) **22**:586–597. doi:10.1038/nm.4106

21. Gomez de Agüero M, Ganal-Vonarburg SC, Fuhrer T, Rupp S, Uchimura Y, Li H, Steinert A, Heikenwalder M, Hapfelmeier S, Sauer U, et al. The maternal microbiota drives early postnatal innate immune development. *Science* (2016) **351**:1296–1302. doi:10.1126/science.aad2571
22. Lamas B, Hernandez-Galan L, Galipeau HJ, Constante M, Clarizio A, Jury J, Breyner NM, Caminero A, Rueda G, Hayes CL, et al. Aryl hydrocarbon receptor ligand production by the gut microbiota is decreased in celiac disease leading to intestinal inflammation. *Sci Transl Med* (2020)
23. Sonner JK, Keil M, Falk-Paulsen M, Mishra N, Rehman A, Kramer M, Deumelandt K, Röwe J, Sanghvi K, Wolf L, et al. Dietary tryptophan links encephalogenicity of autoreactive T cells with gut microbial ecology. *Nat Commun* (2019) **10**:4877. doi:10.1038/s41467-019-12776-4
24. Modoux M, Rolhion N, Mani S, Sokol H. Tryptophan metabolism as a pharmacological target. *Trends Pharmacol Sci* (2020) doi:10.1016/j.tips.2020.11.006
25. Brawner KM, Yeramilli VA, Duck LW, Van Der Pol W, Smythies LE, Morrow CD, Elson CO, Martin CA. Depletion of dietary aryl hydrocarbon receptor ligands alters microbiota composition and function. *Sci Rep* (2019) **9**:14724. doi:10.1038/s41598-019-51194-w
26. Schanz O, Chijiwa R, Cengiz SC, Majlesain Y, Weighardt H, Takeyama H, Förster I. Dietary ahr ligands regulate ahrr expression in intestinal immune cells and intestinal microbiota composition. *Int J Mol Sci* (2020) **21**: doi:10.3390/ijms21093189
27. Korecka A, Dona A, Lahiri S, Tett AJ, Al-Asmakh M, Braniste V, D'Arienzo R, Abbaspour A, Reichardt N, Fujii-Kuriyama Y, et al. Bidirectional communication between the Aryl hydrocarbon Receptor (AhR) and the microbiome tunes host metabolism. *npj Biofilms and Microbiomes* (2016) **2**:16014. doi:10.1038/npjbiofilms.2016.14
28. Julliard W, De Wolfe TJ, Fechner JH, Safdar N, Agni R, Mezrich JD. Amelioration of *Clostridium difficile* Infection in Mice by Dietary Supplementation With Indole-3-carbinol. *Ann Surg* (2017) **265**:1183–1191. doi:10.1097/SLA.0000000000001830
29. Li Y, Innocentin S, Withers DR, Roberts NA, Gallagher AR, Grigorieva EF, Wilhelm C, Veldhoen M. Exogenous stimuli maintain intraepithelial lymphocytes via aryl hydrocarbon receptor activation. *Cell* (2011) **147**:629–640. doi:10.1016/j.cell.2011.09.025
30. Cervantes-Barragan L, Chai JN, Tianero MD, Di Luccia B, Ahern PP, Merriman J, Cortez VS, Caparon MG, Donia MS, Gilfillan S, et al. *Lactobacillus reuteri* induces gut intraepithelial CD4⁺CD8 α ⁺ T cells. *Science* (2017) **357**:806–810. doi:10.1126/science.aah5825
31. Kiss EA, Vonarbourg C, Kopfmann S, Hobeika E, Finke D, Esser C, Diefenbach A. Natural aryl hydrocarbon receptor ligands control organogenesis of intestinal lymphoid follicles. *Science* (2011) **334**:1561–1565. doi:10.1126/science.1214914
32. Lee JS, Cella M, McDonald KG, Garlanda C, Kennedy GD, Nukaya M, Mantovani A, Kopan R, Bradfield CA, Newberry RD, et al. AHR drives the development of gut ILC22 cells and postnatal lymphoid tissues via pathways dependent on and independent of Notch. *Nat Immunol* (2011) **13**:144–151. doi:10.1038/ni.2187

33. Zheng Y, Valdez PA, Danilenko DM, Hu Y, Sa SM, Gong Q, Abbas AR, Modrusan Z, Ghilardi N, de Sauvage FJ, et al. Interleukin-22 mediates early host defense against attaching and effacing bacterial pathogens. *Nat Med* (2008) **14**:282–289. doi:10.1038/nm1720
34. Islam J, Sato S, Watanabe K, Watanabe T, Ardiansyah, Hirahara K, Aoyama Y, Tomita S, Aso H, Komai M, et al. Dietary tryptophan alleviates dextran sodium sulfate-induced colitis through aryl hydrocarbon receptor in mice. *J Nutr Biochem* (2017) **42**:43–50. doi:10.1016/j.jnutbio.2016.12.019
35. Wu Y, He Q, Yu L, Pham Q, Cheung L, Kim YS, Wang TTY, Smith AD. Indole-3-Carbinol Inhibits *Citrobacter rodentium* Infection through Multiple Pathways Including Reduction of Bacterial Adhesion and Enhancement of Cytotoxic T Cell Activity. *Nutrients* (2020) **12**: doi:10.3390/nu12040917
36. Ye J, Qiu J, Bostick JW, Ueda A, Schjerven H, Li S, Jobin C, Chen Z-ME, Zhou L. The aryl hydrocarbon receptor preferentially marks and promotes gut regulatory T cells. *Cell Rep* (2017) **21**:2277–2290. doi:10.1016/j.celrep.2017.10.114
37. Wang B, Sun S, Liu M, Chen H, Liu N, Wu Z, Wu G, Dai Z. Dietary L-Tryptophan Regulates Colonic Serotonin Homeostasis in Mice with Dextran Sodium Sulfate-Induced Colitis. *J Nutr* (2020) **150**:1966–1976. doi:10.1093/jn/nxaa129
38. Alkarkoushi RR, Hui Y, Tavakoli AS, Singh U, Nagarkatti P, Nagarkatti M, Chatzistamou I, Bam M, Testerman TL. Immune and microRNA responses to *Helicobacter muridarum* infection and indole-3-carbinol during colitis. *World J Gastroenterol* (2020) **26**:4763–4785. doi:10.3748/wjg.v26.i32.4763
39. Metidji A, Omenetti S, Crotta S, Li Y, Nye E, Ross E, Li V, Maradana MR, Schiering C, Stockinger B. The Environmental Sensor AHR Protects from Inflammatory Damage by Maintaining Intestinal Stem Cell Homeostasis and Barrier Integrity. *Immunity* (2018) **49**:353–362.e5. doi:10.1016/j.immuni.2018.07.010
40. Goudot C, Coillard A, Villani A-C, Gueguen P, Cros A, Sarkizova S, Tang-Huau T-L, Bohec M, Baulande S, Hacohen N, et al. Aryl Hydrocarbon Receptor Controls Monocyte Differentiation into Dendritic Cells versus Macrophages. *Immunity* (2017) **47**:582–596.e6. doi:10.1016/j.immuni.2017.08.016
41. Rothhammer V, Borucki DM, Tjon EC, Takenaka MC, Chao C-C, Ardura-Fabregat A, de Lima KA, Gutiérrez-Vázquez C, Hewson P, Staszewski O, et al. Microglial control of astrocytes in response to microbial metabolites. *Nature* (2018) **557**:724–728. doi:10.1038/s41586-018-0119-x
42. Scheiermann C, Gibbs J, Ince L, Loudon A. Clocking in to immunity. *Nat Rev Immunol* (2018) **18**:423–437. doi:10.1038/s41577-018-0008-4
43. Jaeger C, Tischkau SA. Role of aryl hydrocarbon receptor in circadian clock disruption and metabolic dysfunction. *Environ Health Insights* (2016) **10**:133–141. doi:10.4137/EHI.S38343
44. Zarrinpar A, Chaix A, Yooseph S, Panda S. Diet and feeding pattern affect the diurnal dynamics of the gut microbiome. *Cell Metab* (2014) **20**:1006–1017.

doi:10.1016/j.cmet.2014.11.008

45. Lamas B, Richard ML, Leducq V, Pham H-P, Michel M-L, Da Costa G, Bridonneau C, Jegou S, Hoffmann TW, Natividad JM, et al. CARD9 impacts colitis by altering gut microbiota metabolism of tryptophan into aryl hydrocarbon receptor ligands. *Nat Med* (2016) **22**:598–605. doi:10.1038/nm.4102
46. Lamas B, Hernandez-Galan L, Galipeau HJ, Constante M, Clarizio A, Jury J, Breyner NM, Caminero A, Rueda G, Hayes CL, et al. Aryl hydrocarbon receptor ligand production by the gut microbiota is decreased in celiac disease leading to intestinal inflammation. *Sci Transl Med* (2020) **12**: doi:10.1126/scitranslmed.aba0624

Figure legends




Figure 1. Sources of nutritional AhR ligands. Nutritional AhR ligands are derived either from the breakdown of food components or from tryptophan catabolism by intestinal microbiota. Several types of indoles are present in cruciferous vegetables. In particular, indole-3-carbinol (I3C) is converted in the stomach into high affinity AhR ligands Diindolylmethane (DIM) and indole[3,2-b]carbazole (ICZ). Tryptophan is metabolised by *Lactobacillus* bacteria into indole-3-acetic acid (IAA), tryptamine (TA) and 3-methyl indole. In addition, tryptophanase-expressing bacteria degrade tryptophan into indole, which is metabolised by host liver cells into indoxyl-3-sulfate (I3S) and indole-3-propionic acid (IPA).




Figure 2. Nutritional AhR ligands in intestinal immunity. Nutritional AhR ligands are involved in the maintenance of intestinal intra-epithelial lymphocytes (IEL) and type innate-like lymphoid cells (ILC3). ILC3 are the main producers of IL22, which acts on intestinal epithelial cells to induce the secretion of antimicrobial peptides. In the absence of dietary AhR ligands, IEL and ILC3 are reduced, and microbiota diversity is altered.




Figure 3. Nutritional AhR ligands in monocyte differentiation. Nutritional AhR ligands modulate monocyte differentiation in the skin. AhR activation from dietary agonists favors monocyte differentiation towards In the absence of AhR signaling, monocytes differentiate preferentially into macrophages.dendritic cells via the induction of the transcription factors Irf4 and Blimp-1.

Table 1. Nutritional AhR ligands classification. Reported nutritional AhR agonists classified based on their origin.

Compound	Abbreviations	Source	Reference
Quercetin		Dietary ligands	8-10
Resveratrol			
Indole-3-acetonitrile	IAN	Dietary ligands	11
Indolo(3,4)bicarbazole			
Indole-3-carbinole	I3C	Dietary ligands	11
3,3'-diindolylmethane	DIM		
Indole[3,2-b] carbazole		Host metabolism from food components	15
Linear trimer, [2-(indol-3-ylmethyl)-indol-3-yl]indol-3-ylmethane	LTR ₁		
1-(3-hydroxymethyl)-indolyl-3-indolylmethane	HI-IM	Host metabolism from food components	16
5-hydroxyindole-3-acetic acid	5-HIAA		
Indole-3-acetic acid	IAA	Microbiota metabolism from food components	12-13
Tryptamine	TA		
3-methyl indole		Microbiota metabolism from food components	13
Indole-3-aldehyde	IAlD		
Butyrate		Microbiota metabolism from food components	15
Indoxyl-3-sulfate	I3S		
Indole-3-propionic acid	IPA	Host metabolism from tryptophan catabolites from microbiota	14