

Combined spectroscopy and intensity interferometry to determine the distances of the blue supergiants P Cygni and Rigel

E. S. G. de Almeida ^{1,1★†}, M. Hugbart ^{2★}, A. Domiciano de Souza,¹ J.-P. Rivet,¹ F. Vakili,¹ A. Siciak,² G. Labeyrie,² O. Garde,³ N. Matthews,² O. Lai ¹, D. Vernet,⁴ R. Kaiser² and W. Guerin ²

¹Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Laboratoire Lagrange, F-06304 Nice, France

²Université Côte d'Azur, CNRS, Institut de Physique de Nice, F-06103 Nice Cedex 2, France

³2SPOT, F-38690 Chabons, France

⁴Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, UMS Galilée, F-06304 Nice, France

Accepted 2022 May 31. Received 2022 May 31; in original form 2022 March 22

ABSTRACT

In this paper, we report on spatial intensity interferometry measurements within the $H\alpha$ line on two stars: the Luminous Blue Variable supergiant P Cygni and the late-type B supergiant Rigel. The experimental setup was upgraded to allow simultaneous measurement of two polarization channels, instead of one in our previous setup, and the zero baseline correlation function on-sky to validate independent estimates obtained from the stellar spectrum and the instrumental spectral throughput. Combined with simultaneous spectra measurements and based on radiative transfer models calculated with the code CMFGEN, we were able to fit our measured visibility curves to extract the stellar distances. Our distance determinations for both P Cygni (1.61 ± 0.18 kpc) and Rigel (0.26 ± 0.02 kpc) agree very well with the values provided by astrometry with the Gaia and *Hipparcos* missions, respectively. This result for Rigel was obtained by adopting a stellar luminosity of $L_\star = 123\,000 L_\odot$, which is reported in the literature as being consistent with the *Hipparcos* distance to Rigel. However, due to the lack of consensus on Rigel's luminosity, we also explore how the adoption of the stellar luminosity in our models affects our distance determination for Rigel. In conclusion, we support, in an independent way, the distance to Rigel as the one provided by the *Hipparcos* mission, when taking the luminosity of $123\,000 L_\odot$ at face value. This study is the first successful step towards extending the application of the Wind Momentum Luminosity Relation method for distance calibration from an LBV supergiant to a more normal late-type B supergiant.

Key words: techniques: interferometric – stars: distances – stars: massive – stars: individual: P Cygni – stars: individual: Rigel – stars: winds, outflows.

1 INTRODUCTION

Fifty years after Hanbury Brown and his team's pioneering contribution to stellar astrophysics (Hanbury Brown, Davis & Allen 1974) using the Narrabri high angular resolution facility (Hanbury Brown 1974), intensity interferometry has entered a new age of development for several reasons. First, progress in photonics components, efficient detectors that record single photon events, fast electronics, and digital correlators, all offer enhanced sensitivity for the same amount of light collection area (Guérin et al. 2017, 2018). Secondly, large imaging air Cherenkov telescope arrays, primarily built for high-energy astrophysics, have been recently successful in performing stellar intensity interferometry (Abeysekara et al. 2020; Acciari et al. 2020). In comparison with the Narrabri interferometer, these arrays allow faster and more accurate measurements of angular

diameters of hot stars. Hence, future large-scale facilities, such as the Cherenkov Telescope Array, open new perspectives for very high angular resolution synthesis imaging by intensity interferometry, especially at short visible wavelengths (Nuñez & Domiciano de Souza 2015; Dravins 2016).

Our team is following a complementary path by using traditional astronomical telescopes with photon-counting avalanche photodiodes (APDs) that feed a fast time tagger, which computes the temporal correlations in real time (Lai et al. 2018; Rivet et al. 2018). One advantage of our approach is that the optical quality of the telescope allows the collimation of the beam and subsequently a narrow spectral filtering with a bandpass of $\Delta\lambda \sim 1$ nm. This gives the possibility to scrutinize the star under observation within spectral lines, in absorption or emission, and therefore access to the physical conditions in their extended atmospheres or to other effects that finely depend on the wavelength across the visible spectrum. Then, using state-of-the-art radiative transfer models to reproduce high-resolution spectroscopy and photometry (spectral energy distribution, SED), we can constrain the fundamental parameters of the star and thus synthesize intensity maps projected across the sky, from which the computed visibilities can be compared to the measured

* E-mail: elisson.saldanha@oca.eu, elisson.dealmeida@uv.cl (ESGdeA); mathilde.hugbart@inphyni.cnrs.fr (MH)

† Current affiliation: Instituto de Física y Astronomía, Universidad de Valparaíso. Av. Gran Bretaña 1111, Casilla 5030, Valparaíso, Chile.

ones. This approach has been effectively demonstrated with intensity interferometry of the archetype Luminous Blue Variable (LBV) supergiant star P Cygni (P Cyg) to provide its distance (Rivet et al. 2020), independently from OB association distance estimates (Turner et al. 2001) or global astrometry with *Gaia* (Brown et al. 2021).

In this paper, we aim at going beyond this first successful determination of the distance of P Cygni by a second observation at a different epoch of the same star, and by extending the method to the blue supergiant Rigel (β Ori), which presents a much weaker emission in the $H\alpha$ line. Thus, we can examine the application of the so-called Wind Momentum Luminosity Relation (WLR hereafter, Kudritzki, Lennon & Puls 1995; Puls et al. 1996; Kudritzki & Puls 2000) in the context of temporal-spectral variability of LBV stars, here P Cygni and different B supergiants (Rigel), for the use of the WLR as an independent distance indicator for extragalactic sources such as the Virgo cluster (Kudritzki et al. 1999) in the future. For this purpose, the experimental setup has been improved and now exploits the two orthogonal polarizations, instead of one as done in the previous setup. It also allows measuring simultaneously the spatial intensity correlation function with two telescopes and the temporal intensity correlation on one telescope used to calibrate the spatial intensity correlations at zero baseline. Based on the measured spectra and the radiative transfer code CMFGEN, we determine the distance of P Cygni and Rigel from modelling their measured visibilities.

This paper is organized as follows. We first describe our upgraded experimental setup in the next section (Section 2), which allows in particular measuring the polarization-resolved intensity correlation functions, and then we present our observations and the spatial intensity correlation functions measured on P Cygni and Rigel (Section 3). Section 4 describes the radiative transfer code CMFGEN and our modelling approach to determine the distances of P Cygni and Rigel. Finally, our results are compared to the ones found in literature and then summarized in Section 5.

2 EXPERIMENTAL SETUP

2.1 Setup

The details on the experimental setup can be found in Guerin et al. (2017, 2018) and Rivet et al. (2020). Briefly, the light is first collected by two telescopes T_1 and T_2 , as shown in Fig. 1. The observation runs were performed in 2020 at the C2PU facility on the Plateau de Calern site of Observatoire de la Côte d'Azur (OCA). The distance between the telescopes is equal to 15 m, with an almost east–west orientation, which gives access to different projected baselines during the night. Each telescope has a diameter of 1.04 m and a central obstruction of 0.3 m in diameter. The two telescopes with yoke equatorial mounts ensure that there is no field rotation.

The light collected by the telescopes then goes through a coupling assembly (CA) attached to the telescopes and depicted in Fig. 2. This CA has been modified compared to the one previously used in Rivet et al. (2020). It now allows extracting the two orthogonal polarizations, labelled H and V for, respectively, horizontal and vertical in the rest of the paper, thanks to a polarizing beamsplitter (PBS). This PBS is needed to select one polarization mode and was also present in the previous CA. However, while before the photons on the V channel were lost, this new CA allows exploiting all the photons collected by the telescopes. The extinction ratio (ratio of the unpolarized optical power to the optical power with polarization parallel to the polarizer) of the PBS is better than 10^{-3} in transmission, ensuring a high degree of linear polarization for the transmitted beam. However, this extinction ratio can be as high as a




Figure 1. Experimental setup to measure the spatial intensity correlation function on two orthogonal polarizations (labelled H for horizontal and V for vertical). T_1 and T_2 : telescopes, CA: coupling assembly, see Fig. 2 for more details, MMF: multimode fiber, FBS: fibered beamsplitter, APD: avalanche photodiode, CC: 50 Ω coaxial cables, TDC: time-to-digital converter.




Figure 2. Coupling assembly (CA) placed at the telescope Cassegrain ports, to perform spectral filtering, polarization separation, and fiber injection. DBS: Dichroic Beam Splitter, used to send the shortest wavelengths of the input beam to a guiding CMOS camera. L_1 : Diverging lens ($f_1 = -50$ mm) to collimate the input beam on the narrow-band interference filter (bandwidth $\Delta\lambda = 1$ nm, central wavelength $\lambda_0 = 656.3$ nm). PBS: Polarizing Beam Splitter, splitting the beam into a V-polarized beam (reflected beam) and an H-polarized beam (transmitted beam). P: Linear polarizer plate to improve the polarization purity on the reflected beam (V polarization). L_2 : Pair of converging lens ($f_2 = +20$ mm) to focus the two output beams on the tip of 100 μm core multimode fibers (MMF).

few percent in reflection. To overcome this, a second polarizer (P) parallel to the polarization of the reflected beam is added after the PBS. Each polarized beam is then injected in a 100 μm core diameter multimode fiber. A spectral filtering is performed before the PBS on the collimated beam. The bandwidth of the filter is $\Delta\lambda = 1$ nm with a central frequency $\lambda_0 = 656.3$ nm corresponding to the $H\alpha$ line. The two CAs, placed at the Cassegrain focus of each telescope, have been checked in the laboratory on an unresolved artificial source and the correlation functions are the same for each CA as well as for each polarization channel.

The outputs of the CAs are connected to single-photon avalanche photodiodes (APDs). The counts detected by the different APDs are time-tagged by a time-to-digital converter (TDC). The time response τ_{el} of this electronic setup is of the order of a few hundreds of picoseconds, mainly limited by the time resolution of the photodiodes. From

the time-tagged photon stream the correlation function between pairs of detectors with the same polarization state, thus corresponding to six different correlation functions, are computed in real time and saved on a computer. Instrumental path-length differences in the electronic and fiber cabling between correlated detectors are accounted for in software using values measured in the laboratory. Any path-length fluctuations in the cabling are negligible compared to our relative tolerance of ~ 15 cm set by the corresponding light traveltime during a time equivalent to our temporal resolution. Before averaging, each stored correlation function is shifted in time by the fixed instrumental delay, and by the computed geometrical optical path delay that is variable throughout the night, such that the expected signal appears at $\tau = 0$ corresponding to zero optical path delay.

The setup at the output of the two telescopes has been modified compared to Rivet et al. (2020). The new setup now allows measuring the correlation function with the two telescopes at the same time as the correlation function at zero baseline using one telescope. The zero baseline calibration done on-sky reduces systematic uncertainties in comparison to previous methods which required laboratory measurements on an artificial unresolved light source. To do so, the setup on the first telescope (T_1) is slightly different from the one placed on the second telescope (T_2). On T_1 , each CA output is first connected to a fibered beamsplitter (FBS) whose outputs illuminate two APDs. This allows measuring the temporal intensity correlation function for zero baseline $g_V^{(2)}(\tau)$ and $g_H^{(2)}(\tau)$ (Guerin et al. 2017), for two orthogonal polarization states. This provides a calibration of the zero-baseline visibility in real time. The APDs are placed in shielded boxes and are put far apart from each other (typically 2 m far apart). This configuration allows us to avoid spurious correlations that corresponds to unwanted extra peaks above noise and that were previously observed (Rivet et al. 2020) and that needed to be removed with a ‘white’ signal.

Measuring the coincidences between two APDs set on different telescopes and for the same polarization gives the spatial intensity correlation functions $g_V^{(2)}(\tau, r_B)$ or $g_H^{(2)}(\tau, r_B)$, where r_B is the projected baseline. One can note that for a given polarization, one has two possible pairs of photodiodes (one APD on T_2 and two APDs on T_1). The two corresponding intensity correlation functions are expected to be identical. Therefore, we sum the two correlation functions computed by the TDC, before normalization, to obtain one spatial intensity correlation function for each polarization. The final signal-to-noise ratio (SNR) is then the same as if the photon flux were not split into two at one telescope.

2.2 Measured quantities

2.2.1 Temporal intensity correlation function

At zero separation ($r_B = 0$), one measures the temporal intensity correlation function also called the temporal second-order correlation function:

$$g^{(2)}(\tau) = \frac{\langle I(t, 0)I(t + \tau, 0) \rangle}{\langle I(t, 0) \rangle^2}, \quad (1)$$

with $\langle . \rangle$ corresponding to the averaging over the whole observing time t , and $I(t, 0)$ the intensity collected at zero baseline. For chaotic light, $g^{(2)}(\tau)$ is linked to the temporal electric field correlation function $g^{(1)}(\tau)$ through the Siegert relation (Siegert 1943; Loudon 1973; Ferreira et al. 2020):

$$g^{(2)}(\tau) = 1 + |g^{(1)}(\tau)|^2. \quad (2)$$




Figure 3. Temporal intensity correlation functions measured on P Cygni. (a) Horizontal polarization channel, SNR = 6.5 given by the Gaussian fit (dashed line). (b) Vertical polarization channel, SNR = 7.7.

Finally, the Wiener–Khinchine theorem (Wiener 1930; Khinchine 1934) relates $g^{(1)}(\tau)$ and the optical spectrum $S(\omega)$:

$$S(\omega) = \int g^{(1)}(\tau) e^{i\omega\tau} d\tau. \quad (3)$$

For chaotic light such as the one coming from stars and for an infinite electronic bandwidth, the expected contrast $C_{\text{exp}} = g^{(2)}(0) - g^{(2)}(\infty)$ is equal to 1, leading to the so-called bunching effect which corresponds to a peak above 1 on the temporal intensity correlation function at zero delay, as can be seen for example in Fig. 3. The coherence time τ_c , which corresponds to the $g^{(2)}(\tau)$ decay time and thus to the typical width of the theoretical bunching peak, is inversely proportional to the spectral bandwidth, of the order of 1 ps for $\Delta\lambda = 1$ nm at visible wavelengths. This coherence time is thus much smaller than the time response of our experimental setup τ_{el} . The measured bunching peak corresponds to the bunching peak of width τ_c convolved with the mutual time response of our detectors $\tau_{\text{el}} > \tau_c$. This leads to a reduction of the measured contrast $C \simeq \tau_c/\tau_{\text{el}}$ and a $g^{(2)}(\tau)$ decay time mainly limited by τ_{el} . On the other hand, the area of the bunching peak A_{BP} is proportional to the height times the decay time of the bunching peak $C\tau_{\text{el}} \simeq \tau_c$, and is thus independent of the electronic time response.

The APD time response can slightly vary from one detector to another, which leads to a variation of τ_{el} and thus a variation of the contrast depending on the detector pair used to measure the correlation function. Furthermore, we have observed a slight dependency of the electronic time response on the APDs count rate, which means that the contrast can slightly vary during an observational run. On the contrary, as said before, the area of the bunching peak does not depend on the electronic time response and thus is also independent from the count rate, at least at first order. The area, directly related to the coherence time, is therefore a more robust quantity compared to the contrast. This is what will be used throughout this paper.

2.2.2 Spatial intensity correlation function

The spatial intensity correlation function is defined as

$$g^{(2)}(\tau, r_B) = \frac{\langle I(t, 0)I(t + \tau, r_B) \rangle}{\langle I(t, 0) \rangle \langle I(t, r_B) \rangle}, \quad (4)$$

with $I(t, r_B)$ the intensity collected with a second telescope, r_B being also called the projected baseline. The angular size can be inferred from the typical spatial decay of $g^{(2)}(\tau, r_B)$, which depends on the visibility $V(r_B)$, measured in amplitude interferometry, as follows (Loudon 1973; Labeyrie, Lipson & Nisenson 2006):

$$g^{(2)}(\tau, r_B) = 1 + |V(r_B)|^2 |g^{(1)}(\tau)|^2, \quad (5)$$

Table 1. Observing conditions for the runs performed on P Cygni. Begin and end dates are in UTC (*ISO 8601* compact format). a is the airmass range. ϵ is the seeing estimate, provided by the GDIMM instrument (Ziad et al. 2012; Aristidi et al. 2014) of the CATS station (Calern Atmospheric Turbulence Station, Chabé et al. 2016). The numbers are given as median values over the whole nights. A ‘–’ symbol means that no GDIMM measurements were available that night.

| Begin | End | a | ϵ |
|----------------|----------------|--------------------|-------------|
| 20200804T0004Z | 20200804T0202Z | 1.03 → 1.21 | – |
| 20200804T2034Z | 20200805T0257Z | 1.12 → 1.00 → 1.42 | – |
| 20200805T1931Z | 20200806T0340Z | 1.26 → 1.00 → 1.67 | 0.97 arcsec |
| 20200806T2002Z | 20200807T0403Z | 1.17 → 1.00 → 1.87 | 0.74 arcsec |
| 20200807T1928Z | 20200808T0354Z | 1.25 → 1.00 → 1.83 | 0.76 arcsec |
| 20200808T1924Z | 20200809T0343Z | 1.25 → 1.00 → 1.77 | 0.88 arcsec |

$$= 1 + |V(r_B)|^2 (g^{(2)}(\tau) - 1). \quad (6)$$

Based on the same arguments as in the previous section, we measure the area $A(r_B)$ of the bunching peak for different baselines to infer the visibility:

$$A(r_B) = |V(r_B)|^2 A(r_B = 0). \quad (7)$$

The quantity $A(r_B = 0)$ is measured on T_1 with the temporal correlation function, as explained in Section 2.1.

Finally, with an SNR of 17 at best, we did not detect any polarization difference on our measurements on stars. It is beyond the scope of this paper to provide a detailed description of the circumstellar environments of P Cygni and Rigel. Nevertheless, such a lack of polarization difference indicates that we are not able to detect any asymmetry in the circumstellar environments of both P Cygni and Rigel within our error bars. We thus decide to merge the temporal and spatial correlation functions obtained for each polarization. The SNR is increased by typically a factor $\sqrt{2}$ as expected.

3 OBSERVATIONS

3.1 Intensity correlations on P Cygni

P Cygni was observed at C2PU, within the $H\alpha$ line, between 2020 August 3 and 2020 August 9 as reported in Table 1. The total integration time was 40.3 h. We detected in average 320×10^3 cps (counts per second) per detector on telescope 1 (T_1 , where the signal from each polarization channel is split into two) and 715×10^3 cps per detector on telescope 2 (T_2). Those new data will be compared to the ones obtained during our first observations in 2018, also within the $H\alpha$ line, but with only one polarization channel. The results have been published in Rivet et al. (2020), where we estimated the distance of P Cygni by comparing the measured visibilities to simulations computed with the code CMFGEN, with the physical parameters of P Cygni constrained by contemporaneous observed spectra.

3.1.1 Temporal intensity correlations

As mentioned in Section 2, the measurements done with only one telescope allow measuring the temporal intensity correlation function. The results for the two polarization channels are presented in Fig. 3 with a Gaussian fit on top of it. One can see that the width and the contrast are slightly different resulting mainly from a difference in the temporal response of each detector. As stated in Section 2.2, taking the area of the bunching peak allows getting rid of the different




Figure 4. Spectrum of P Cygni (plain curve), with its maximum value normalized to one, reported on the AAVSO database and measured on 2020 August 8. The dotted line corresponds to the $H\alpha$ filter transmission.

electronic time responses. The areas are extracted from the Gaussian fit. One can see in Fig. 3 that the fit is correctly superimposed to the data, with a reduced χ^2 equal to 0.94. We get: $A_H = 1.95 \pm 0.3$, $A_V = 2.3 \pm 0.3$, and $A = 2.1 \pm 0.2$ ps if we merge the two temporal correlation functions (before fitting), with 1σ statistical uncertainties. They are compatible with each other within the error bars.

To calculate the expected area of the bunching peak, we need to numerically compute the $g^{(2)}(\tau)$ function from the spectrum using equations (2) and (3), as explained in Rivet et al. (2020). Fig. 4 presents one spectrum reported on the AAVSO database (AAVSO 2020) and measured on 2020 August 8 using an eShel spectrometer (from Shelyak) with a resolving power $R = 11\,650$. We can observe a strong emission line, slightly weaker than the one reported in 2018 (Rivet et al. 2020). For a point-like source, we get $A_{\text{exp}} = 2.35$ ps (2.55 ps in 2018). This value is compatible with A_V within 1σ and with A_H within 2σ . In the rest of the paper, we will thus consider that we can use the value measured with one telescope as the zero baseline value $A = A(r_B = 0)$.

3.1.2 Spatial intensity correlations

For the spatial intensity correlation functions $g^{(2)}(\tau, r_B)$, we first merge the two correlation functions computed by the TDC, before normalization, and measured with the same polarization, between one detector on T_2 and the two other ones on T_1 , and then the correlations obtained for both polarizations. The procedure to take into account the geometrical optical delay between the telescopes and the variation of the baseline during the night are explained in Guerin et al. (2018) and Rivet et al. (2020). The merged normalized intensity correlations are presented in Fig. 5 for zero baseline and for projected baselines $9.5 < r_B < 13.4$ and $13.4 < r_B < 15$, corresponding to mean baselines of 11.8 and 14.4 m, respectively. These intervals have been chosen to get the same number of individual correlation functions, measured with an exposure time of 10 s, within each baseline interval.

The squared visibility is calculated by dividing the bunching area by the value measured at zero baseline: $V^2 = A(r_B)/A(r_B = 0)$. The results are plotted in Fig. 6 and reported in Table 2, taking into account the uncertainty on $A(r_B)$ and $A(r_B = 0)$. Finally, we can compare our new data to the ones already published in Rivet et al. (2020), represented by the black circles in Fig. 6. At that time, the squared visibilities were computed by dividing the measured contrast by the contrast expected from the spectrum. Our new data are also compatible with the previous ones within the error bars. Fig. 6 also shows the fitted visibility curve from our reference CMFGEN model




Figure 5. Experimental intensity correlations merging the V and H polarization contributions, obtained on P Cygni (grey line), and their Gaussian fits (black dashed line) for different baselines: Intensity correlation measured (a) with one telescope, corresponding to the zero baseline, merging the V and H polarization contributions presented in Fig. 3, (b) with two telescopes for projected baselines $9.5 < r_B < 13.4$ m, and (c) $13.4 < r_B < 15$ m.




Figure 6. Squared visibility measured on P Cygni as a function of projected baseline r_B . Blue points: Squared visibility $V^2 = A(r_B)/A(r_B = 0)$ reported in this paper at $\lambda = 656.3$ nm when the intensity correlation for both polarizations are merged. The vertical errors bars are given at 1σ and are extracted from the Gaussian fit used to calculate the area of $g^{(2)}(\tau, r_B)$: $A(r_B)$ and $A(r_B = 0)$. The horizontal error bars correspond to the minimum and maximum projected baselines. The data are fitted (blue line) from our reference CMFGEN model for P Cygni (Table 5; Section 4.2) using equation (11) (Section 4.4) with the distance d to P Cygni as the only free parameter. The shaded area corresponds to the 1σ uncertainty given by the fit. Black circles: $V^2 = C/C_{\text{exp}}$, with C the contrast of the bunching peak, reported in (Rivet et al. 2020) also at $\lambda = 656.3$ nm in the H channel, and C_{exp} the contrast expected from the measured spectrum. The fit to the data corresponds to the black dashed line.

Table 2. Summary of the observations on P Cygni with r_B the average baseline, A the area of the bunching peak extracted from a Gaussian fit on the correlation function, and V^2 the squared visibility calculated by dividing the bunching area by the value measured at zero baseline: $V^2 = A(r_B)/A(r_B = 0)$.

| r_B (m) | A (ps) | V^2 |
|----------------|-----------------|-----------------|
| 0 | 2.13 ± 0.22 | 1 |
| 11.8 ± 1.2 | 1.0 ± 0.2 | 0.47 ± 0.12 |
| 14.4 ± 0.5 | 0.78 ± 0.15 | 0.37 ± 0.08 |

Table 3. Observing conditions for the runs performed on Rigel. Same symbols as in Table 1.

| Begin | End | a | ϵ |
|----------------|----------------|--|------------|
| 20200129T1912Z | 20200129T2101Z | $1.69 \rightarrow 1.62 \rightarrow 1.66$ | 2.27 |
| 20200131T1955Z | 20200131T2320Z | $1.62 \rightarrow 1.62 \rightarrow 2.65$ | 1.52 |
| 20200201T1749Z | 20200201T2309Z | $1.99 \rightarrow 1.62 \rightarrow 2.54$ | 2.89 |
| 20200203T1823Z | 20200203T2219Z | $1.78 \rightarrow 1.62 \rightarrow 2.08$ | 3.10 |
| 20200204T1736Z | 20200204T2110Z | $2.00 \rightarrow 1.62 \rightarrow 1.74$ | – |
| 20200205T1714Z | 20200205T2300Z | $2.14 \rightarrow 1.62 \rightarrow 2.65$ | 2.36 |
| 20200206T1728Z | 20200206T2300Z | $2.00 \rightarrow 1.62 \rightarrow 2.71$ | 2.07 |
| 20200207T1732Z | 20200207T2233Z | $1.95 \rightarrow 1.62 \rightarrow 2.38$ | 1.81 |
| 20200208T1821Z | 20200208T2255Z | $1.72 \rightarrow 1.62 \rightarrow 2.76$ | – |
| 20200211T1835Z | 20200211T2017Z | $1.66 \rightarrow 1.62 \rightarrow 1.67$ | – |
| 20200212T1842Z | 20200212T2254Z | $1.64 \rightarrow 1.62 \rightarrow 3.07$ | 1.96 |
| 20200214T1755Z | 20200214T2242Z | $1.73 \rightarrow 1.62 \rightarrow 2.97$ | 1.07 |
| 20200215T1738Z | 20200215T2014Z | $1.77 \rightarrow 1.62 \rightarrow 1.70$ | 1.18 |




Figure 7. Temporal intensity correlation functions measured on Rigel. (a) Horizontal polarization channel, SNR = 17.4 given by the Gaussian fit (dashed line). (b) Vertical polarization channel, SNR = 14.

for P Cygni with the distance to this star as the only free parameter which is discussed in Section 4.

3.2 Intensity correlations on Rigel (β Ori)

The observations on Rigel have been performed during 13 nights between 2020 January 29 and 2020 February 15, still within the $H\alpha$ line, with a total integration time of 50.6 h. The mean number of counts was 1.25×10^6 cps per detector on T_1 and 2.9×10^6 cps per detector on T_2 . The observation dates and atmospheric conditions are summarized in Table 3.

3.2.1 Temporal intensity correlations

The bunching peaks are visible on all the correlations functions, either on the $g^{(2)}(\tau)$ functions obtained with one telescope, as shown in Fig. 7, or on the $g^{(2)}(\tau, r_B)$ functions. The areas extracted from the Gaussian fit of $g^{(2)}(\tau)$ are: $A_H = 1.22 \pm 0.07$, $A_V = 1.12 \pm 0.08$, and $A = 1.14 \pm 0.05$ ps when the correlation functions obtained on the two polarizations are merged (before fitting), thus compatible with each other within the error bars. The measured areas must be compared to what we expect from the filtered star spectrum. Fig. 8 presents one spectrum reported in the ARAS database (ARAS 2020) in 2020 between the 1st and 13th of February. We can observe a small absorption and emission line. For a point-like source, we get $A_{\text{exp}} = 1.22$ ps, equivalent actually to what would be obtained for a flat spectrum. This value is compatible with A_H within 1σ and




Figure 8. One of the spectra of Rigel (solid line) reported in the ARAS database in 2020 between the 1st and 13th of February, with its maximum value normalized to one. The dotted line corresponds to the H α filter transmission.




Figure 9. Experimental intensity correlations, merging the V and H polarization contributions, obtained on Rigel (grey line), and their Gaussian fits (black dashed line) for different baselines: (a) Intensity correlation measured with one telescope, corresponding to the zero baseline, merging the V and H polarization contributions presented in Fig. 7, (b) with two telescopes for projected baselines $8.9 < r_B < 13.7$, (c) $13.7 < r_B < 14.7$, and (d) $14.7 < r_B < 15$ m.

with A_V within 2σ . As before, we thus also consider that the areas measured with one telescope can be used as the zero baseline values.

3.2.2 Spatial intensity correlations

To calculate the spatial intensity correlation functions, we use the same procedure as the one detailed in Section 3.1.2. The SNR is higher than the one obtained on P Cygni due to the fact that Rigel is significantly brighter and due to a slightly longer integration time. We divide the baselines in three ranges, $8.9 < r_B < 13.7$ m, $13.7 < r_B < 14.7$ m, and $14.7 < r_B < 15$ m, corresponding to mean baselines of 12.2, 14.3, and 14.90 m, respectively. The different intensity correlation functions are plotted in Fig. 9. Fig. 10 presents the squared visibility $V^2 = A(r_B)/A(r_B = 0)$ as a function of baseline. The SNR is similar for all measurements at large baselines, of the order of 13.5. The results are also summarized in Table 4. Ahead of the discussion (Section 4), Fig. 10 also shows the fitted visibility curve from our reference CMFGEN model for Rigel with the distance to this star as the only free parameter.




Figure 10. Squared visibility $V^2 = A/A(r_B = 0)$ measured on Rigel as a function of the projected baseline. The vertical errors bars are given at 1σ and are extracted from the Gaussian fit used to calculate the area of $g^{(2)}(\tau, r_B)$: $A(r_B)$ and $A(r_B = 0)$. The horizontal error bars correspond to the minimum and maximum projected baselines. The data are fitted (black line) from our reference CMFGEN model for Rigel (Table 5; Section 4.2) using equation (11) (Section 4.4) with the distance d to Rigel as the only free parameter. The shaded area corresponds to the 1σ uncertainty given by the fit.

Table 4. Summary of the observations on Rigel. Same symbols as in Table 2.

| r_B (m) | A (ps) | V^2 |
|------------------|-----------------|-----------------|
| 0 | 1.14 ± 0.05 | 1 |
| 12.2 ± 1.2 | 0.85 ± 0.05 | 0.75 ± 0.06 |
| 14.3 ± 0.3 | 0.79 ± 0.05 | 0.70 ± 0.05 |
| 14.90 ± 0.09 | 0.82 ± 0.04 | 0.72 ± 0.05 |

4 ESTIMATION OF THE STAR DISTANCES BASED ON THE CODE CMFGEN

4.1 The code CMFGEN

To provide a robust interpretation of our interferometric data of P Cygni and Rigel, we used unified photosphere-wind models calculated with the non-LTE (local thermodynamic equilibrium) radiative transfer code CMFGEN (Hillier & Miller 1998). For a set of stellar and wind parameters, CMFGEN solves in an iterative way the radiative transfer, statistical, and equilibrium equations in the comoving frame. This code has been successfully used in the literature to model observables of different types of hot stars and then to determine their stellar and wind parameters (e.g. see Hillier 2012, 2020, and references therein).

It is well-understood that radiative line-driven winds of hot stars show density fluctuations due to local agglomerations of matter, called wind clumps (e.g. Eversberg, Lépine & Moffat 1998). This feature must be taken into account in the modelling of hot stars in order to well reproduce their observables, and then to obtain more accurate estimates of the wind mass-loss rates (Bouret, Lanz & Hillier 2005; Fullerton, Massa & Prinja 2006; Davies, Vink & Oudmaier 2007). The code CMFGEN allows us to implement the effect of wind clumping, using the so-called microclumping approximation (Hillier et al. 2001). This assumes a void interclump medium and wind clumps' sizes smaller than the photon mean-free path for any value of wavelength. In CMFGEN, the wind clumping is parametrized by the volume filling factor, $f(r)$, as follows

$$f(r) = f_\infty + (1 - f_\infty)e^{-\frac{v(r)}{v_{\text{initial}}}}, \quad (8)$$

where r is the distance from the center of the star, f_∞ is the filling factor value at $r \rightarrow \infty$, $v(r)$ is the wind velocity, and v_{initial} is the onset

Table 5. Summary of the main stellar and wind parameters of our CMFGEN reference models for P Cygni and Rigel based on the match to the observed H α line profiles observed in 2020 for P Cygni (from the AAVSO database) and for Rigel (from the ARAS database).

| Parameters | P Cygni | Rigel |
|---|----------------------|----------------------|
| L_* (L_\odot) | 610 000 | 123 000 |
| T_{eff} (K) | 18 700 | 12 500 |
| $\log g$ | 2.25 | 1.75 |
| R_* (R_\odot) | 75 | 75 |
| M_* (M_\odot) | 37 | 12 |
| \dot{M} ($M_\odot \text{ yr}^{-1}$) | 3.3×10^{-5} | 8.1×10^{-8} |
| f_∞ | 0.5 | 0.1 |
| v_∞ (km s^{-1}) | 185 | 300 |
| β | 2.3 | 1.0 |

velocity of clumping in the wind. Despite current efforts to solve the radiative transfer equations in a self-consistent way with the wind hydrodynamics (e.g. Gormaz-Matamala et al. 2021), the wind density and velocity structures are usually adopted in CMFGEN as performed in Rivet et al. (2020). Then the validity of the adopted wind density and velocity is only justified after the match with observations.

The wind velocity $v(r)$ is parametrized by the so-called β -law approximation, shown in its simplest form below:

$$v(r) = v_\infty \left(1 - \frac{R_*}{r} \right)^\beta, \quad (9)$$

where v_∞ is the wind terminal velocity and R_* is the stellar radius (r higher than R_*). Therefore, assuming a stationary symmetric wind and taking into account the clumping factor $f(r)$, the wind density and velocity are related to each other by the equation of mass continuity:

$$\dot{M} = 4\pi r^2 \rho(r) v(r) f(r), \quad (10)$$

where $\rho(r)$ is the wind density and \dot{M} is the wind mass-loss rate, assumed in this case to be constant at any point of the wind.

4.2 Model parameters of P Cygni and Rigel

Both P Cygni and Rigel were previously studied using the code CMFGEN to model different types of observables (e.g. Najarro, Hillier & Stahl 1997; Najarro 2001; Chesneau et al. 2010; Richardson et al. 2013; Chesneau et al. 2014; Rivet et al. 2020). In Table 5, we summarize the main stellar and wind parameters of our reference models for P Cygni and Rigel: stellar luminosity (L_*), effective temperature (T_{eff}), gravity surface acceleration ($\log g$), radius (R_*), mass (M_*), mass-loss rate (\dot{M}), wind clumping factor (f_∞), terminal velocity (v_∞), and the wind velocity law exponent (β).

Following Rivet et al. (2020), we adopted the stellar and wind parameters for P Cygni based on the study of Najarro (2001) that used CMFGEN to model the ultraviolet, visible, and infrared spectrum of P Cygni. As described in Rivet et al. (2020), the chemical composition of our CMFGEN models for P Cygni also follows Najarro (2001).

For Rigel, the model parameters are based on Chesneau et al. (2010, 2014) that used CMFGEN to model interferometric data of Rigel.¹ Solar chemical composition is assumed in our models for Rigel, as in Chesneau et al. (2010, 2014). In turn, these interferometric studies based their analysis on the stellar and wind parameters derived for Rigel by Przybilla et al. (2006) and Markova et al. (2008).

¹See table 1 of Chesneau et al. (2014).

The adopted values for the photospheric parameters T_{eff} and $\log g$ are in good agreement with other spectroscopic studies of Rigel in the visible region. For instance, using models calculated with the non-LTE radiative transfer code FASTWIND (Santolaya-Rey, Puls & Herrero 1997; Puls et al. 2005; Rivero González et al. 2012), Haucke et al. (2018) derived $T_{\text{eff}} = 12\,700 \pm 500$ K and $\log g = 1.7 \pm 0.1$ for Rigel. With respect to the wind mass-loss rate, Chesneau et al. (2014) tested values ranging between $\sim 1.0 \times 10^{-7}$ and $\sim 1.0 \times 10^{-6} M_\odot \text{ yr}^{-1}$. Based on interferometric quantities measured in the H α line, Chesneau et al. (2010) determined $\dot{M} = 1.5 \times 10^{-7} M_\odot \text{ yr}^{-1}$ for Rigel.²

Our model for Rigel shown in Table 5 is a modified version from ‘Chesneau’s model’ for this star (Chesneau et al. 2010, 2014). Instead of assuming $L_* = 279\,000 L_\odot$, as done by these authors, we initially assumed a lower stellar luminosity of $L_* = 123\,000 L_\odot$. From modeling the SED of Rigel, Haucke et al. (2018) derived $L_* = 123\,000 L_\odot$ for Rigel when taking into account the distance of ~ 265 pc from *Hipparcos* parallaxes (van Leeuwen 2007). So, in comparison with Chesneau et al. (2010, 2014), the stellar radius and mass are also changed considering $T_{\text{eff}} = 12\,500$ K and $\log g = 1.75$. Here, we assumed $\dot{M} = 8.1 \times 10^{-8} M_\odot \text{ yr}^{-1}$ in order to have the same wind density parameter for recombination lines [e.g. see equation (39) of Puls, Vink & Najarro 2008] of Chesneau’s model ($\dot{M} = 1.5 \times 10^{-7} M_\odot \text{ yr}^{-1}$). This change on \dot{M} allows our modified model to produce a very similar H α profile in comparison to the original parameter set from Chesneau et al. (2010, 2014) for Rigel.

We followed the approach described above aiming to verify if our distance determination for Rigel is compatible with the results provided by *Hipparcos* parallaxes from van Leeuwen (2007). Nevertheless, as will be discussed in Section 5, we also determined the distance of Rigel by assuming the same model parameters from Chesneau et al. (2010, 2014), that is, considering a higher stellar luminosity of $L_* = 279\,000 L_\odot$ for this star.

It is beyond the scope of this paper to determine the stellar and wind parameters of both P Cygni and Rigel. Nevertheless, the wind parameters of our reference CMFGEN models for these stars are tuned in order to provide a good match to the observed H α line profiles, as discussed in the following.

4.3 Comparison to the observed spectrum of P Cygni and Rigel

We compared our reference models to public spectroscopic data of P Cygni from the AAVSO database and Rigel from the ARAS database (ARAS 2020) observed in 2020. Due to the high H α variability of P Cygni (Markova et al. 2001) and Rigel (Kaufer et al. 1996), we analysed observed H α line profiles that were recorded close in time to our interferometric measurements: August 8th (P Cygni) and February 5th (Rigel). Our reference CMFGEN models for P Cygni and Rigel are compared to their visible spectra around the H α line in Figs 11 and 12, respectively. For the comparison to the spectrum of P Cygni observed in 2018, we used the same CMFGEN model from Rivet et al. (2020).

²As pointed by Chesneau et al. (2014), the mass-loss rate of Rigel derived from Br γ is up to about one order of magnitude higher than the one derived from H α . Since our study is based on observations centred at the H α line, our reference value for Rigel’s mass-loss rate is based on $\dot{M} = 1.5 \times 10^{-7} M_\odot \text{ yr}^{-1}$.




Figure 11. Comparison between the observed visible spectrum of P Cygni (black line) in $\sim 6555\text{--}6686\text{ \AA}$ and our reference CMFGEN models for this star (red line). The synthetic spectrum is convoluted with $v\sin i = 35\text{ km s}^{-1}$ and spectral resolving powers $R = 9000$ (top panels) and $11\,650$ (bottom panel). These reference models are based on the match to the $H\alpha$ line profile of P Cygni observed at different epochs. Top panel: same reference model for P Cygni from Rivet et al. (2020), based on the match to the $H\alpha$ line profile observed on 2018 August 14. Bottom panel: reference model for P Cygni based on the match to $H\alpha$ line profile observed on 2020 August 8 (Table 5). These two reference models for P Cygni (based on different epochs) allow us to refine our distance determination for this star as shown in Section 4.4.1.

4.3.1 P Cygni

From Fig. 11, one sees that the $H\alpha$ line profile of P Cygni observed in 2020 is slightly less intense in comparison with the observations performed in 2018. These spectra were observed with a spectral resolving power of $R = 9000$ (2018) and $R = 11\,650$ (2020). This difference in R is not able to explain the different emission components of $H\alpha$ as observed for P Cygni in 2018 and 2020. With respect to our CMFGEN model used to mimic the $H\alpha$ line profile observed in 2018 (see table 4 of Rivet et al. 2020), we only varied the wind mass-loss rate in order to match the $H\alpha$ line profile observed in 2020. We followed this simple approach on our analysis since the change on this wind parameter has a strong impact on the synthetic emission component of $H\alpha$. In addition, the wind mass-loss rate of P Cygni is thought to be variable over time, combined with a change in its stellar radius and effective temperature (Markova et al. 2001). Having all the other parameters fixed from the model for P Cygni used in (Rivet et al. 2020), we needed to reduce the mass-loss rate by about 18 per cent (from 4.0×10^{-5} in 2018 to $3.3 \times 10^{-5}\text{ M}_{\odot}\text{ yr}^{-1}$ in 2020). The latter value is closer to the mass-loss rate determined from Najjarro (2001) of $2.4 \times 10^{-5}\text{ M}_{\odot}\text{ yr}^{-1}$. Thus, based on the $H\alpha$ spectroscopic data from 2020, the adopted physical model in this paper for P Cygni only differs to the model of Rivet et al. (2020) with respect to the mass-loss rate. The main stellar parameters are listed in Table 5.




Figure 12. Top panel: comparison between the observed visible spectrum of Rigel (black line) in $\sim 6555\text{--}6686\text{ \AA}$ and our reference CMFGEN model for this star (solid red line). Bottom panel: zoom around the $H\alpha$ line profile. The synthetic spectrum is convoluted with $v\sin i = 36\text{ km s}^{-1}$ and spectral resolving power $R = 9000$. The observed spectrum was recorded on 2020 February 5. The parameters of our reference model for Rigel are shown in Table 5. Dashed blue line: synthetic spectrum calculated using Rigel's parameters from Table 5, with exception of the wind velocity law exponent ($\beta = 1.5$ instead of $\beta = 1.0$). In Section 4.4.2, we discuss the effect of varying β (from 1.0 to 1.5) on our distance determination for Rigel.

As pointed out by Markova et al. (2001), the wind mass-loss rate of P Cygni should change by about 19 per cent in a time-scale of about 7 yr. This time-scale is longer than the 2-yr time-span between our analysed spectra of P Cygni (2018 and 2020). In addition, variations of stellar parameters were not taken into account in our modeling of the more recent $H\alpha$ spectroscopic data of P Cygni (2020). In short, despite our ability to reproduce fairly well the $H\alpha$ line profile of P Cygni observed in 2020 using such a less intense wind model, it is beyond the scope of the current paper to state that the intensity of the wind of P Cygni varied in this way during this 2-yr period.

4.3.2 Rigel

In comparison with P Cygni, the blue supergiant Rigel shows a more complex variation of the morphology of the $H\alpha$ line profile over time. Its $H\alpha$ line can be found as classical and inverse P Cygni profiles, double- and single-peak emission, or pure-absorption (e.g. see Morrison, Rother & Kurschat 2008, and references therein). In particular, the $H\alpha$ line profile of Rigel formed a P Cygni profile during the period of our interferometric observations performed in 2020 February (see Fig. 12). The $H\alpha$ emission component of Rigel is much weaker than the one found in P Cygni due to the large difference in the wind mass-loss rate between these stars (see Table 5). Overall our reference CMFGEN model for Rigel reproduces fairly well its observed $H\alpha$ line profile. C II $\lambda\lambda 6580, 6585$ and He I $\lambda 6678$ of Rigel are pure-photospheric lines (almost insensitive to changes on




Figure 13. Effective H α radial profile I_{eff} of the reference CMFGEN model for P Cygni as a function of radial coordinate ϖ given in units of the stellar photospheric radius (clipped at $20R_{\star}$ for better visualization). The curves are normalized to one at $\varpi = 0$. Solid blue line: profile corresponding to the observations made in 2020 within the H α filter; Dotted blue curve: profile in the continuum ($\lambda = 655$ nm), calculated from the spectrum measured in 2020; Dashed black curve: CMFGEN profile corresponding to the observations made in 2018 within the H α filter (Rivet et al. 2020). The spectra of these models (around the H α line) are shown in Fig. 11.

the wind’s parameters) and are well reproduced by our model. This indicates that both the physical conditions of Rigel’s photosphere and wind are well described by our adopted CMFGEN model for this star.

We are aware that values of β much larger than 1.0 (up to ≈ 3.0) can be required to reproduce the H α line of OB supergiants (e.g. see Puls et al. 2008, and references therein). For instance, based on models calculated with the code FASTWIND, Markova et al. (2008) derived β up to 1.5 for their sample of late-type B supergiant (which included Rigel), but without specifying a value for Rigel, while Haucke et al. (2018) derived $\beta = 2.6$ for Rigel also based on spectroscopic modelling using FASTWIND.

As our physical model for Rigel is based on Chesneau et al. (2010, 2014), and the wind velocity law exponent is not specified in these studies, we initially adopted $\beta = 1.0$. We then tested the effect of higher value of β on the modeling of the observed H α profile of Rigel. From Fig. 12, we see that our model with $\beta = 1.5$ tends to reproduce better the absorption component of H α while the emission component is misfitted, considering all the other parameters fixed. It is beyond the scope of this paper to determine the wind velocity law exponent of Rigel. In addition, it is known that matching simultaneously the observed H α absorption and emission components of Rigel is a hard task (e.g. see fig. 1 of Haucke et al. 2018). Nevertheless, in comparison to previous quantitative spectroscopic studies of Rigel (Markova et al. 2008; Chesneau et al. 2010; Haucke et al. 2018), the reference CMFGEN model for Rigel considered in this paper ($\beta = 1.0$) is able to reproduce fairly well the observed overall H α profile. As will be discussed in Section 4.4, we tested how the adoption of two different values of β for Rigel’s wind affects our distance estimation for this star.

4.4 Discussion on luminosities and distances of P Cygni and Rigel from quantitative spectroscopy and intensity interferometry

4.4.1 Distance to P Cygni

Following the same procedure as the one adopted in Rivet et al. (2020), we compute the effective radial intensity profile $I_{\text{eff}}(\varpi)$ within the H α filter from the CMFGEN models [equation (7) in Rivet

et al. (2020)], where the coordinate ϖ is the impact parameter, following the same notation used in Rivet et al. (2020). In the (p , z) coordinate system, the impact parameter is usually denoted by p and is related to the radial coordinate r used in equations (9) and (10) (e.g. see figs 7–29 of Mihalas 1978).

The normalized intensity profile within the H α filter of our reference CMFGEN models for P Cygni (based on 2018 and 2020 observations) is plotted in Fig. 13. For comparison, Fig. 13 also shows the intensity profile in the continuum region (at $\lambda = 655$ nm) close to H α of our model calculated from the spectrum measured in 2020. One sees that the profiles measured in 2020 and in 2018 (Rivet et al. 2020) are similar since the difference in the mass-loss rate between our reference models for P Cygni is not very large, changing from 4.0×10^{-5} (2018) to $3.3 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ (2020). As expected, the width of the intensity profile is larger within the H α line than in the continuum, that is, H α is formed throughout a more extended region in the wind of P Cygni. This happens due to the high value of mass-loss rate of P Cygni’s wind, resulting in a larger flux contribution from the wind in H α than in the continuum.

Then, the corresponding normalized squared visibility V^2 is computed using the Hankel transform (circular symmetry):

$$V^2 = \left| \frac{\int_0^{\infty} I_{\text{eff}}(x) J_0(2\pi x q) 2\pi x \, dx}{\int_0^{\infty} I_{\text{eff}}(x) 2\pi x \, dx} \right|^2, \quad (11)$$

where J_0 is the zeroth-order Bessel function of the first kind, $x = \varpi/d$ is the radial angular coordinate, with d being the distance to the star and used as a free parameter to fit the data. The radial spatial frequency coordinate associated to x is $q = r_B/\lambda_{\text{eff}}$, corresponding to the average projected baseline r_B divided by the effective wavelength of the observations λ_{eff} .

Using equation (11) and $\lambda_{\text{eff}} = 6562.9 \text{ \AA}$ derived from the spectrum observed in 2020 and the adopted filter, the fit to our data is shown in Fig. 6, with d being the only free parameter. We derived $d_{\text{PCygni, 2020}} = 1.67 \pm 0.26$ kpc in good agreement with the value obtained in 2018 of $d_{\text{PCygni, 2018}} = 1.56 \pm 0.25$ kpc (Rivet et al. 2020). Finally, we refine our distance estimate to P Cygni from averaging $d_{\text{PCygni, 2018}}$ and $d_{\text{PCygni, 2020}}$: $d_{\text{PCygni, averaged}} = 1.61 \pm 0.18$ kpc.

4.4.2 Distance to Rigel

The normalized intensity profile of Rigel within the H α filter, associated with the adopted CMFGEN model, is plotted in Fig. 14 in addition to the one obtained in the continuum ($\lambda = 655$ nm). In comparison with P Cygni (see Fig. 13), one sees that the intensity profile of our model for Rigel within the H α filter quickly drops as a function of impact parameter since Rigel’s wind has a much lower mass-loss rate than P Cygni (up to about two orders of magnitude). Nevertheless, as our model for Rigel shows a weak emission component in the H α line profile, one can still see a higher I_{eff} in H α than in the continuum region at the innermost part of the wind up to $\sim 2\text{--}3 R_{\star}$.

In Fig. 10, we show the squared visibility V^2 for Rigel, also fitted using equation (11) from the effective profile. From that, we derived the distance to Rigel as $d_{\text{Rigel, } \beta=1.0} = 0.26 \pm 0.02$ kpc, considering the parameters of our CMFGEN model listed in Table 5, that is, with $\beta = 1.0$.

As discussed in Section 4.3.2, our model with $\beta = 1.0$ better reproduces the emission component of the H α line, while a larger value of β , namely, 1.5, better reproduces the absorption component. When considering our model with $\beta = 1.5$ (having all the other parameters fixed), we derived $d_{\text{Rigel, } \beta=1.5} = 0.28 \pm 0.02$ kpc, still




Figure 14. Effective $H\alpha$ radial profile I_{eff} of the reference CMFGEN model for Rigel as a function of radial coordinate w given in units of the stellar photospheric radius (clipped at $3R_{\star}$ for better visualization). The curves are normalized to one at $w = 0$. Solid line: profile corresponding to the observations within the $H\alpha$ filter; Dotted curve: CMFGEN profile in the continuum ($\lambda = 655$ nm), calculated from the spectrum measured in 2020. The spectrum of this model (around the $H\alpha$ line) is shown in Fig. 12.

compatible at 1σ with the distance obtained for $\beta = 1.0$. In conclusion, since these distance estimates are in good agreement, we consider, in this paper, that the distance to Rigel is $d_{\text{Rigel}} = 0.26 \pm 0.02$ kpc, based on our reference CMFGEN model for Rigel presented in Table 5.

4.4.3 Discussion on Rigel's luminosity

As discussed in Section 4.2, instead of adopting $L_{\star} = 279\,000 L_{\odot}$ from Chesneau's model for Rigel, we initially adopted the stellar luminosity for Rigel according to the value provided by Haucke et al. (2018) of $L_{\star} = 123\,000 L_{\odot}$, which is based on the fit to Rigel's SED taking into account $d_{\text{Rigel, Hipparcos}}$. We followed this approach since *Hipparcos* parallaxes are usually considered reliable for close stars (up to ~ 500 pc), as Rigel, and should be taken at face value when compared to other distance determination methods.

However, quite discrepant values for the stellar luminosity and distance of Rigel are reported in the literature. For instance, the spectroscopic study of Przybilla et al. (2006) determined $\log L_{\star}/L_{\odot} = 5.34 \pm 0.08$ for Rigel, that is, with a luminosity ranging from 182 000 to 263 000 L_{\odot} . These authors adopted a distance of ~ 360 pc for Rigel based on Hoffleit & Jaschek (1982) considering the membership of Rigel to the τ Ori R1 complex. An even larger distance value up to ~ 500 pc has been considered due to its membership of the Ori OB1 association (Humphreys 1978).

We evaluated the impact of the adopted stellar luminosity on our distance determination of Rigel. For this purpose, we derived its distance considering the same parameters as used by the studies of Chesneau et al. (2010, 2014). In comparison with the parameters for Rigel listed in Table 5, the following parameters are changed: L_{\star} from 123 000 to 279 000 L_{\odot} , R_{\star} from 75 to 113 R_{\odot} , M_{\star} from 12 to 26 M_{\odot} , and \dot{M} from 8.1×10^{-8} to $1.5 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$. The latter parameter is changed in order to have the same wind density parameter than our CMFGEN model shown in Table 5.

Following the method described in Section 4.4, we fitted the theoretical visibility curve to our data of Rigel, but considering Chesneau's model for Rigel. From that, we derived the distance to Rigel as $d_{\text{Rigel, Chesneau}} = 0.42 \pm 0.03$ kpc.³ As expected, when

³Here, we use the subscript 'Chesneau' to denote that the distance value of Rigel was derived considering $L_{\star} = 279\,000 L_{\odot}$.




Figure 15. Comparison between the observed SED of Rigel from ~ 0.35 to $\sim 20 \mu\text{m}$ (black points) and synthetic SEDs of our reference CMFGEN models (red lines). SED is shown in units of Jy and μm . Model SEDs are calculated in the continuum. Top panel: synthetic SED calculated from our model for Rigel as listed in Table 5 and considering the distance value of $d_{\text{Rigel}} = 0.26$ kpc. Bottom panel: synthetic SED calculated from our adopted model for Rigel as used in Chesneau et al. (2010, 2014) and considering the distance value of $d_{\text{Rigel, Chesneau}} = 0.42$ kpc. See text for discussion.

assuming a higher luminosity in our modelling, the derived distance to Rigel is quite larger than the one found from *Hipparcos* parallaxes, being closer to other results in the literature, for instance, as reported in Przybilla et al. (2006) that considered a stellar luminosity for Rigel up to $\sim 263\,000 L_{\star}$.

Fig. 15 compares the observed SED⁴ of Rigel with our model SEDs for Rigel considering different values of luminosity: $L_{\star} = 123\,000 L_{\odot}$ and $L_{\star} = 279\,000 L_{\odot}$. For each case, we take into account the derived distance associated with each model: $d_{\text{Rigel}} = 0.26$ kpc ($L_{\star} = 123\,000 L_{\odot}$) and $d_{\text{Rigel, Chesneau}} = 0.42$ kpc ($L_{\star} = 279\,000 L_{\odot}$). The effect of interstellar medium extinction is included in the model SEDs following the reddening law from Cardelli, Clayton & Mathis (1989), assuming a colour excess $E(B - V) = 0.05$ (Przybilla et al. 2006) and a total to selective extinction ratio $R_V = 3.1$ as a typical value for Galactic stars. One sees that in both cases our distance estimates of Rigel are consistent with the stellar luminosity in order to reproduce well the observed SED. In conclusion, the adoption of the stellar luminosity in our CMFGEN models highly affects the distance determination when fitting our interferometric data. Nevertheless, we verify that our derived distances are self-consistent with the adopted luminosity when looking at other observables than interferometry, as photometry, and therefore providing an independent check to our results.

5 CONCLUSION

In this paper, we have observed P Cygni within the $H\alpha$ line, which allowed us to determine the distance based on the CMFGEN model.

⁴Public data available in the Centre de Données astronomiques de Strasbourg: <https://cds.u-strasbg.fr/>.

Taking into account the observations done in 2018 (Rivet et al. 2020) and in 2020 for this paper, we get $d_{\text{PCyg, averaged}} = 1.61 \pm 0.18$ kpc, improving the uncertainty by a factor of 1.4 compared to our published distance in Rivet et al. (2020). The comparison to other distance determinations has already been done in our previous paper (Rivet et al. 2020), as well as the discussion on the controversy on this distance measurement. Since then, a new distance has been given by the *Gaia* global astrometry mission in its third early data release (EDR3), with $d_{\text{PCyg, EDR3}} = 1.60^{+0.21}_{-0.17}$ kpc (Brown et al. 2021), in excellent agreement with our result.

Rigel’s parallax has not been measured by the *Gaia* mission. With an apparent magnitude of 0.13 (Ducati 2002) in the *V* band (500–600 nm), Rigel exceeds *Gaia*’s detector saturation limit,⁵ which is of about 3 (*G* band, 330–1050 nm). From the fit to our interferometric data using a self-consistent physical model of Rigel, our distance determination to Rigel, $d_{\text{Rigel}} = 0.26 \pm 0.02$ kpc, agrees very well with the one found from *Hipparcos* parallaxes of $d_{\text{Rigel, Hipparcos}} = 0.27 \pm 0.03$ kpc (van Leeuwen 2007).

Therefore, when compared with results provided by direct parallax measurements, our distance estimate method works well for both P Cygni and Rigel in spite of these hot supergiant stars showing quite different H α line profiles: P Cygni shows a strong and fully developed P Cygni profile in H α , while our analysed spectrum of Rigel shows a much weaker emission in H α .

Due to the lack of consensus on the luminosity of Rigel, we also fitted our interferometric data of this star using a higher luminosity than the initially fixed value of $123\,000 L_{\odot}$: $279\,000 L_{\odot}$ from Chesneau et al. (2014). As expected, in this case, we infer a larger distance to Rigel of $d_{\text{Rigel, Chesneau}} = 0.42 \pm 0.03$ kpc. This result is in line with some distance estimations that are reported for Rigel in the literature, indicating a larger distance for this star (up to ~ 0.5 kpc) than the one found from *Hipparcos* parallaxes. Both our lower and higher luminosity models for Rigel are self-consistent with the inferred distances when looking the observed SED of Rigel. However, we point out that parallax measurements from the *Hipparcos* mission are very usually considered reliable for nearby stars as Rigel. In conclusion, when taking the luminosity of $123\,000 L_{\odot}$ at face value for Rigel, our results support, in an independent way, the distance to Rigel as the one provided by the *Hipparcos* mission. Said differently, our study supports that Rigel’s luminosity of $123\,000 L_{\odot}$ is consistent with its distance provided by the *Hipparcos* mission.

Previous spectroscopic studies of OBA supergiants (used due to their high values of luminosity) showed that the WLR is a promising tool to derive extragalactic distances (e.g. see Bresolin & Kudritzki 2004, and references therein). On the other hand, it is well-known that there are disagreements among both theoretical and measured⁶ (modified) wind momentum for different types of hot stars (e.g. see Kudritzki et al. 1999; Vink, de Koter & Lamers 2000; Marcolino et al. 2009; Haucke et al. 2018; de Almeida et al. 2019; Björklund et al. 2021). Based on that, the employment of the WLR to derive stellar distances should be taken with caution. Nevertheless, it is still important to evaluate its consistency as distance indicator since it can bring new insights on the wind properties of hot stars such as their real values of mass-loss rates.

It is beyond the scope of this paper to provide a robust quantitative evaluation of the WLR since we studied only two stars. Nevertheless,

based on our results discussed above, we can thus claim having achieved a first successful step towards extending the application of the WLR method for distance calibration from an LBV supergiant to a more *normal* late-type B supergiant, including the temporal variability of P Cygni. The latter has been observed at multiepochs in 2005 and 2008 by the NPOI amplitude interferometer in the light of H α emission (Balan et al. 2010) across a 10 nm spectral filter. It was found that its diameter did not change between these epochs within 10 per cent, which is also our present measurement uncertainties. It clearly appears that we need to gain an order of magnitude in visibility precision on our intensity interferometry observations in order to robustly establish the WLR method application to cosmological distance measurements beyond the local Virgo and Fornax clusters of galaxies, taking advantage of their extreme luminosities compared to standard candles. For example, P Cygni absolute magnitude is smaller by about 4 compared to the one of δ Cep (Classical Cepheid).

To push further the precision of our visibility measurements, some improvements have been already implemented on our setup and presented in this paper. First, we now measure the correlation functions on the two polarization channels. Since the telescopes are mounted on equatorial mounts and the CA are attached to the telescopes, this can be used to detect any effect that would depend on polarization. In this paper, since we do not measure any polarization difference within our experimental uncertainties, we used these two channels to improve our SNR. As expected for white noise, we obtain an improvement of a factor of $\sqrt{2}$, paving the way to multichannel (two polarizations and multispectral) measurements. The second improvement comes from the fact that we measure, at the same time, the spatial correlation function with two telescopes and the spatial correlation function at zero baseline with one telescope. This calibration at zero baseline, that was done before in the laboratory on an artificial unresolved star, leads to smaller systematic error. This is not the goal of this paper to discuss in details systematic errors, especially since we are mainly limited by statistical uncertainties. However, for high-precision measurements that will be done in the future, with large telescopes in particular, one needs to take this aspect into account. We will need for example to characterize the impact on systematic uncertainties of the filter calibration or of the beam collimation on this filter.

A future goal of this study is to propose an independent method to estimate distances beyond the classical cosmological indicators such as Classical Cepheids that are limited to ~ 30 Mpc. This value should be extended up to ~ 50 Mpc with the *James Webb Space Telescope* (Riess et al. 2009). To reach larger distances, a more recent method has been proposed based on Ultra Long Period Cepheids (e.g. see Fiorentino et al. 2012; Musella et al. 2021), which however needs to be further tested. As for the WLR method, following our recent results, we will engage a systematic intensity interferometric survey of a few tens of the closest and brightest OBA supergiants. Those angular determinations, once combined with linear sizes determination with simultaneous multi-epoch spectrometry and consequent CMFGEN modelling, could establish the luminosity versus wind momentum relation and its application to cosmology.

ACKNOWLEDGEMENTS

We thank the referee Chris Haniff for helping us to improve this paper. We acknowledge the financial support of the UCA-JEDI project ANR-15-IDEX-01, the Doebelin Federation, the OPTIMAL platform, the Région PACA (project I2C) and the French ANR (project I2C, ANR-20-CE31-0003).

⁵A summary of the photometric system and magnitude limits of *Gaia* EDR3 can be found at <https://www.cosmos.esa.int/web/gaia/earlydr3>.

⁶By ‘measured’ we mean modified wind momentum ($\dot{M}v_{\infty}\sqrt{R_{*}}$) that are derived from quantitative spectroscopic analysis.

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- AAVSO, 2020, AAVSO spectral database, Public data available at: <https://app.aavso.org/avspec/obs/7783>
- ARAS, 2020, ARAS spectral database, Public data available at: <http://www.astrosurf.com/aras/>
- Abeysekera A. U. et al., 2020, *Nat. Astron.*, 4, 1164
- Acciari V. A. et al., 2020, *MNRAS*, 491, 1540
- Aristidi E., Fantei-Caujolle Y., Ziad A., Dimur C., Chabé J., Roland B., 2014, in Stepp L. M., Gilmozzi R., Hall H. J., eds, Proc. SPIE Conf. Ser. Vol. 9145, Ground-based and Airborne Telescopes V. SPIE, Bellingham, p. 91453G
- Balan A., Tycner C., Zavala R. T., Benson J. A., Hutter D. J., Templeton M., 2010, *AJ*, 139, 2269
- Björklund R., Sundqvist J. O., Puls J., Najarro F., 2021, *A&A*, 648, A36
- Bouret J. C., Lanz T., Hillier D. J., 2005, *A&A*, 438, 301
- Bresolin F., Kudritzki R. P., 2004, in McWilliam A., Rauch M., eds, Origin and Evolution of the Elements. WORLD SCIENTIFIC, Japan, p. 283
- Brown A. G. A. et al., 2021, *A&A*, 650, C3
- Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
- Chabé J., Ziad A., Fantei-Caujolle Y., Aristidi E., Renaud C., Blary F., Marjani M., 2016, in Hall H. J., Roberto G., Heather M. K., eds, Proc. SPIE Conf. Ser. Vol. 9906, Ground-based and Airborne Telescopes VI. SPIE, Bellingham, p. 99064Z
- Chesneau O. et al., 2010, *A&A*, 521, A5
- Chesneau O., Kaufer A., Stahl O., Colvinter C., Spang A., Dessart L., Prinja R., Chini R., 2014, *A&A*, 566, A125
- Davies B., Vink J. S., Oudmaijer R. D., 2007, *A&A*, 469, 1045
- Dravins D., 2016, in Fabien M., Creech-Eakman M. J., Tuthill P. G., eds, Proc. SPIE Conf. Ser. Vol. 9907, Optical and Infrared Interferometry and Imaging V. SPIE, Bellingham, p. 0M
- Ducati J. R., 2002, VizieR Online Data Catalog: Catalogue of Stellar Photometry in Johnson's 11-color system VizieR Online Data Catalog II/237
- Eversberg T., Lépine S., Moffat A. F. J., 1998, *ApJ*, 494, 799
- Ferreira D., Bachelard R., Guerin W., Kaiser R., Fouché M., 2020, *Am. J. Phys.*, 88, 831
- Fiorentino G. et al., 2012, *Ap&SS*, 341, 143
- Fullerton A. W., Massa D. L., Prinja R. K., 2006, *ApJ*, 637, 1025
- Gormaz-Matamala A. C., Curé M., Hillier D. J., Najarro F., Kubátová B., Kubát J., 2021 *ApJ*, 920, 64
- Guerin W., Dussaux A., Fouché M., Labeyrie G., Rivet J.-P., Vernet D., Vakili F., Kaiser R., 2017, *MNRAS*, 472, 4126
- Guerin W., Rivet J.-P., Fouché M., Labeyrie G., Vernet D., Vakili F., Kaiser R., 2018, *MNRAS*, 480, 245
- Hanbury Brown R., 1974, *The Intensity Interferometer: Its Application to Astronomy*. Taylor & Francis, London
- Hanbury Brown R., Davis J., Allen L. R., 1974, *MNRAS*, 167, 121
- Haucke M., Cidale L. S., Venero R. O. J., Curé M., Kraus M., Kanaan S., Arcos C., 2018, *A&A*, 614, A91
- Hillier D. J., 2012, in Richards M. T., Hubeny I., eds, *From Interacting Binaries to Exoplanets: Essential Modeling Tools*, Vol. 282. p. 229
- Hillier D. J., 2020, *Galaxies*, 8, 60
- Hillier D. J., Miller D. L., 1998, *ApJ*, 496, 407
- Hillier D. J., Davidson K., Ishibashi K., Gull T., 2001, *ApJ*, 553, 837
- Hoffleit D., Jaschek C., 1982, *The Bright Star Catalogue*. Fourth revised edition. (Containing data compiled through 1979)
- Humphreys R. M., 1978, *ApJS*, 38, 309
- Kaufer A. et al., 1996, *A&A*, 314, 599
- Khintchine A., 1934, *Math. Ann.*, 109, 604
- Kudritzki R., Puls J., 2000, *ARA&A*, 38, 613
- Kudritzki R. P., Lennon D. J., Puls J., 1995, in Walsh J. R., Danziger I. J., eds, *Science with the VLT*. Springer, Heidelberg, p. 246
- Kudritzki R., Puls J., Lennon D., Venn K., Reetz J., Najarro F., McCarthy J., Herrero A., 1999, *A&A*, 350, 970
- Labeyrie A., Lipson S. G., Nisenson P., 2006, *An Introduction to Optical Stellar Interferometry*. Cambridge Univ. Press, Cambridge
- Lai O. et al., 2018, in Creech-Eakman M. J., Tuthill P. G., Mérand A., eds, Proc. SPIE Conf. Ser. Vol. 10701, Optical and Infrared Interferometry and Imaging VI. SPIE, Bellingham, p. 1070121
- Loudon R., 1973, *The Quantum Theory of Light*. Oxford Science Publications, New York
- Marcolino W. L. F., Bouret J. C., Martins F., Hillier D. J., Lanz T., Escolano C., 2009, *A&A*, 498, 837
- Markova N., Morrison N., Kolka I., Markov H., 2001, *A&A*, 376, 898
- Markova N., Prinja R. K., Markov H., Kolka I., Morrison N., Percy J., Adelman S., 2008, *A&A*, 487, 211
- Mihalas D., 1978, *Stellar Atmospheres*. W.H. Freeman & Co Ltd, San Francisco
- Morrison N. D., Rother R., Kurschat N., 2008, in Hamann W.-R., Feldmeier A., Oskinova L. M., eds, *Clumping in Hot-Star Winds*. Universitätsverlag Potsdam, Potsdam, p. 155
- Musella L., Marconi M., Molinaro R., Fiorentino G., Ripepi V., De Somma G., Moretti M. I., 2021, *MNRAS*, 501, 866
- Najarro F., 2001, in de Groot M., Sterken C., eds, *ASP Conf. Proc. Vol. 233, P Cygni 2000: 400 years of Progress*. Astron. Soc. Pac., San Francisco, p. 133
- Najarro F., Hillier D. J., Stahl O., 1997, *A&A*, 326, 1117
- Núñez P. D., Domiciano de Souza A., 2015, *MNRAS*, 453, 1999
- Przybilla N., Butler K., Becker S. R., Kudritzki R. P., 2006, *A&A*, 445, 1099
- Puls J. et al., 1996, *A&A*, 305, 171
- Puls J., Urbaneja M. A., Venero R., Repolust T., Springmann U., Jokuthy A., Mokiem M. R., 2005, *A&A*, 435, 669
- Puls J., Vink J. S., Najarro F., 2008, *A&AR*, 16, 209
- Richardson N. D. et al., 2013, *ApJ*, 769, 118
- Riess A. G. et al., 2009, *ApJS*, 183, 109
- Rivero González J. G., Puls J., Najarro F., Brott I., 2012, *A&A*, 537, A79
- Rivet J.-P., Vakili F., Lai O., Vernet D., Fouché M., Guerin W., Labeyrie G., Kaiser R., 2018, *Exp. Astron.*, 46, 531
- Rivet J.-P. et al., 2020, *MNRAS*, 494, 218
- Santolaya-Rey A. E., Puls J., Herrero A., 1997, *A&A*, 323, 488
- Siegert A. J. F., 1943, *Report: Radiation Laboratory, On the Fluctuations in Signals Returned by Many Independently Moving Scatterers*. Massachusetts Institute of Technology, Cambridge
- Turner D., Welch G., Graham M., Fairweather D., 2001, *J. Am. Assoc. Var. Star Obs.*, 29, 73
- Vink J. S., de Koter A., Lamers H. J. G. L. M., 2000, *A&A*, 362, 295
- de Almeida E. S. G., Marcolino W. L. F., Bouret J. C., Pereira C. B., 2019, *A&A*, 628, A36
- van Leeuwen F., 2007, *A&A*, 474, 653
- Wiener N., 1930, *Acta Math.*, 55, 117
- Ziad A., Borgnino J., Dali Ali W., Berdja A., Maire J., Martin F., 2012, *J. Opt.*, 14, 045705

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.