



DOI: <http://dx.doi.org/10.5281/zenodo.6874345>

Araştırma Makalesi / Research Article

Effects of Different Storage Periods on The Vase Life of Goldenrod (*Solidago x Hybrida*) Cut Flower

Veysel AYDIN¹ (Orcid ID: 0000-0001-9496-7711), Emine KIRBAY² (Orcid ID: 0000-0002-0343-0829),
Soner KAZAZ³ (Orcid ID: 0000-0002-6644-9690)

¹Batman University, Sason Vocational School, Department of Plant and Animal Production, Batman

²Afyonkarahisar Health Sciences University, Ataturk Health Services Vocational School, Department of
Plant and Animal Production, Afyonkarahisar

³Ankara University, Faculty of Agriculture, Department of Horticulture, Ankara

* Corresponding author (Sorumlu yazar): zmveysel@gmail.com

Geliş Tarihi (Received): 20.06.2022

Kabul Tarihi (Accepted): 22.07.2022

Abstract

Goldenrod (*Solidago x hybrida*) is one of the most traded cut foliage in the world, used as a popular filler in flower arrangements and bouquets. In cut flower, it is desirable to have a long post-harvest storage period and vase life. However, long storage periods shorten vase life. Managing the storage period is important to prevent quality and quantity loss of the product. In this study, it was aimed to determine the effects of different storage periods on the vase life of cut goldenrod flower. *Solidago x hybrida* cv 'Golden Glory' was used as plant material in the study. The flowers were stored dry at different periods (0, 5, 10 and 15 days) in a cold storage (2-4 °C) and then their vase life was measured. The vase life of the flowers used as control was measured without storage. In the study, chlorophyll density in leaves (SPAD) and relative fresh weight at pre and post storage stages were measured. In the research, the longest vase life was found with no-storage application (control) with 15.27 days. This was followed by 12.13 days of application stored for 5 days. Extending the storage period resulted with significant leaf yellowing and reduction of vase life. The results showed that the longest vase life of cut goldenrod flowers was at no-storage application condition. However, although storing for 5 days decreased the vase life of the flowers by 20% compared to the control, it was concluded that cut goldenrod flowers could be stored for 5 days.

Keywords: Solidago, goldenrod, vase life, storage, leaf yellowing, cut flower

INTRODUCTION

Plant parts with and/or without flowers such as fruity and fruitless branches, shoots and leaves are frequently used to prepare all kinds of flower arrangements such as baskets, bouquets, and wreaths to provide freshness and color diversity to improve appearance (Özzambak, 2009; Kazaz, 2012; Mabini and Acedo, 2013). Solidago is well known in the world and in Turkey for a long time, but its use as a cut flower is recent. The name “*Solidago*” is the combination of the Latin words “solid” meaning firm and “ago” meaning to strengthen. It was translated into Turkish as “Altınbaşak” in the “Büyük Bitkiler Klavuzu” in 1952. About 100 goldenrod species naturally grow in the North America and intersection zone of the Europe-Asia Continent (Anonymous, 2008; Kazaz and Karagüzel 2010). *Solidago x hybrida* is among the most traded cut foliage in the world (Ergür et al., 2016). This flower has conical clusters formed by long-lasting and voluminous flowers and is widely used in bouquets and arrangements with its durable and long stem structure. Leaf yellowing is an important problem in cut flower species such as *solidago*, *lilium*, *chrysanthemum*, *alstromeria* and *Matthiola*. Leaf yellowing reduces the quality of flowers and shortens the vase life (Philosoph-Hadas et al., 1996; Hassan et al., 2003; Çelikel, 2013). The economic value of many ornamental plants is directly related to leaf color, and yellowed leaved flowers does not have commercial value. Unfavorable environmental conditions (water loss, temperature, nitrogen deficiency, insufficient light, diseases and pathogen attacks), ethylene, plant growth regulators, genetic structure, chlorophyll and carotenoid pigments are affective on leaf yellowing (Nowak and Rudnicki

1990; Van Doorn, 1997; He et al., 2002; Looze and Van Staaveren, 2003; Ferrante et al., 2004; Shiva, 2006; Woltering, 2008; Christiansen and Gregersen, 2014; Penfold and Buchanan, 2014). Leaf yellowing can be controlled by the use of plant growth regulators at various concentrations (Thomas and Stoddart, 1980, Thimann, 1985). High temperatures increase respiration, sugar/starch consumption, ethylene synthesis, and leaf yellowing. Low temperatures, on the other hand, limit the slowdown of respiration, the reduction of water losses through transpiration, destruction reactions, ethylene biosynthesis, and pathogen attacks (Çelikel and Reid, 2002; Teixeira, 2003; Armitage and Laushman, 2003). The term vase life refers to the length of time from when the flower stem are placed in the vase solution until the loss of their appeal and aesthetic value (Halevy and Mayak 1981; Fanourakis et al. 2013). In the cut flower industry, vase life is an important quality criterion and affects consumer satisfaction and demand for flowers. Amounts of cut flower products get lost due to different reasons from the producer to the consumer in the world. Although it varies according to the species and varieties, globally 25% of the cut flowers produced are lost during the storage and transportation process. Many studies show the positive effects of storage at temperatures above freezing. Dry storage is preferred for long-term storage, and storage in water containing a preservative solution is preferred for short-term storage. Many cut flower species are stored at 0°C just above the freezing point during their storage. Since tropical flowers show cold damage at temperatures below 10 °C, they are stored above 10 °C during storage and transportation (Jones and Moody 1993; Sacalis, 1993; Reid, 2002; Kazaz et al., 2003; Macnish et al., 2009; Çelikel,

2020). The protection of the commercial value of the cut flowers grown in Turkey is important for the development of ornamental plants sector and the country's economy. Cut goldenrod cultivation and its export is becoming more widespread in Turkey. *Solidago x hybrida* 'Golden Glory' variety is one of the popular varieties produced both in Turkey and in the world. Knowing the post-harvest storage strength and vase life of the variety is important for the sector. For this reason, the effects of

different storage periods on the vase life of the cut goldenrod flower were investigated in this study.

MATERIALS and METHODS

The study was carried out in 2022 in the "vase life determination room" and "cold storage room" of the Department of Horticulture, Faculty of Agriculture, Ankara University. 'Golden Glory' cultivar belonging to Goldenrod (*Solidago x hybrida*) species was used as plant material in the study (Figure 1).



Figure 1. *Solidago x hybrida* cv 'Golden Glory'.

The flowers were harvested during the commercial harvest maturity (when 5-10% of the flowers were opened) early in the morning (08.00) at 70 cm stem length and flower were packed in plastic sleeve to minimize water loss which were produced in the greenhouse (in Antalya, Turkiye) of the producer company that produces cut goldenrod. Materials were packaged and placed in perforated

cardboard boxes and brought to the laboratory (in Ankara, Turkiye) within 10 hours where the study will be carried out (Figure 2). Before storage, all flowers were pulsed for 6 hours in buckets containing pure water at a temperature of 21 ± 2 °C. Afterwards, the flowers were stored dry in a cardboard box in a cold store at 2-4 °C.



Figure 2. Sorting and packaging of *Solidago x hybrida* cv 'Golden Glory'.

Measured Factors

Vase Life

The vase life of the flowers was measured in a vase life room with $21 \pm 2^\circ\text{C}$ temperature, $65 \pm 5\%$ relative humidity, 1000 lux light and 12 hours

day (Ueyama and Ichimura, 1998; Ferrante et al., 2007; Lü et al., 2010). In the vase life room, the bottoms of the flowers were cut obliquely from 1 cm and placed in vases containing 750 ml of distilled water length (Figure 3).



Figure 3. Vase life determination room

As a visual parameter in determining the vase life of flowers; leaf senescence, inflorescence, leaf senescence + inflorescence were evaluated separately. The vase life of the flowers was terminated if any of the following criteria were met.

- 1- Leaf yellowing in $\geq 50\%$ of leaves
- 2- Fading, browning, closing in $\geq 50\%$ of opened flowers
- 3- Leaf yellowing in $\geq 50\%$ of leaves + wilting, browning, closing in $\geq 50\%$ of opened flowers

Relative Chlorophyll Exchange Rate During Storage (%)

All the stems used in the study were measured before and after storage. The measurement of chlorophyll change

in the leaves was conducted with the Minolta brand SPAD-502 chlorophyll measuring device on the leaves at 30, 40 and 42 cm height from the base on the branch. Chlorophyll determination was calculated with the help of the following formula presented in Equation 1.

$$\text{RCE (\%)} = (C_t / C_{t=0}) \times 100 \quad (1)$$

C_t : Chlorophyll value after storage (0., 5., 10., ve 15. day)

$C_{t=0}$: Chlorophyll value before storage (0. day)

Relative fresh weight (RFW) During Storage (%)

In order to determine the relative fresh weight changes in the stems during storage, relative fresh weight measurements were conducted on all

stems before and after storage. The stem weights used in the experiment were measured on a digital scale sensitive to 0.01 grams. Relative fresh weight measurements were calculated with the help of the following formula presented in Equation 2. (He et al., 2006; Lü et al., 2010).

$$\text{RFW (\%)} = (W_t / W_{t=0}) \times 100 \quad (2)$$

W_t : Weight of stem after storage (0., 5., 10., ve 15. day)

$W_{t=0}$: Weight of stem before storage (0. day)

Statistical analysis

The experiment was established according to the randomised design with 3 replications where 5 flowers in each replication, with a total of 60 flowers.

SAS statistical software was used in the analysis of the obtained data. The differences between the means were evaluated at the $p \leq 0.01$ level using the Duncan's multiple range test and the correlation data using the Pearson's correlation coefficient.

RESULTS and DISCUSSION

Vase Life (Days)

The effects of different storage periods on vase life are presented in Table 1 and Figure 4. According to the findings obtained in the study, the storage periods was found to be statistically significant on the vase life of cut goldenrod ($p \leq 0.01$).

Table 1. The effects of different storage periods on the vase life of *Solidago x hybrida* cv 'Golden Glory'

Storage period (days)	Vase life (days)
0 (Control)	15.27 ± 0.88 ^{a*}
5	12.13 ± 1.80 ^b
10	4.80 ± 1.78 ^c
15	0.0 ± 0.00 ^d

* The difference between the means with different letters is significant ($p \leq 0.01$)

The longest vase life was determined in the flowers at no-storage application (control) (15.27 days), followed by applications that flowers were stored for 5 days (12.13 days), stored for 10 days (7.43 days), and stored for 15 days (0 days). Prolongation of the storage period resulted in a significant shortening of the vase life of the cut goldenrod flowers compared to the control. Goldenrod flowers stored for 5, 10 and 15 days preserved their opening forms similar to first day. However, the leaves of the flowers stored for 10 days turned yellow partially, and the leaves of the flowers stored for 15 days turned completely to yellow. These leaf yellowing resulted with shortened vase life. Since more than 50% damage

occurred on the leaves of goldenrods stored for 15 days, their vase life was accepted as 0 days. According to the control treatment, the vase life of the flowers decreased by 20% in 5 days of storage, 70% in 10 days of storage and 100% in 15 days of storage periods. The results showed that the longest vase life was in flowers that were no-stored. However, although the vase life of flowers stored for 5 days decreased by 20% compared to the control, this showed that goldenrod flowers would be stored for up to 5 days. Similarly, it has been reported that long and dry storage causes damage to cut flowers and shortens their vase life (Jones and Faragher 1991; Ahmad et al., 2012; Schiappacasse et al., 2014).




Figure 4. The effects of different storage periods on the vase life of *Solidago x hybrida* cv 'Golden Glory'.
* The difference between the means of values with different letters is significant ($p \leq 0.01$).

Relative Chlorophyll Exchange Rate During Storage (%)

The effects of different storage periods on leaf yellowing are presented in Figure 5 and Figure 6. Chlorophyll

density in *Solidago x hybrida* cv 'Golden Glory' leaves showed a decreasing during the storage period. This decrease in the content of chlorophyll was found to be statistically significant ($p \leq 0.01$).




Figure 5. Changes in the chlorophyll content in leaf of *Solidago x hybrida* cv 'Golden Glory' at different storage periods
* The difference between the means with different letters is significant ($p \leq 0.01$)

The chlorophyll content measured as 100% on the first day of the flowers was 86.08% on the 5th day, 78.88 on the 10th day and 61.71% on the 15th day. Although the chlorophyll concentration in the leaves of the flowers

with a storage period of 5 days was statistically significant, this was not reflected in the visual quality. Instead, the leaf yellowing with storage period 10 days was reflected on the visual quality, but did not reach a level that would end

the vase life. Leaf yellowing greater than 50% occurred in the leaves of the flowers stored for 15 days, which resulted with termination. This yellowing in the leaves can be explained by the continuation of the metabolic activities of the flowers and the breakdown of chlorophyll in the leaves as a result of aging. Chlorophyll measurements supported this result. It

has been reported that changes in the amount of ethylene synthesis, respiration, transpiration, genetic structure and plant growth regulators in the plant affect the yellowing of the leaves (Thomas and Stoddart, 1980; Thimann, 1985; Philosoph-Hadas, 1996; Hassan et al., 2003; Sağlam, 2015;).



Figure 6. The effects of different storage periods on leaf yellowing *Solidago x hybrida* cv 'Golden Glory' a) 0 (control) days storage, b) 5 days storage, c) 10 days storage, d) 15 days storage

Relative fresh weight (RFW) During Storage (%)

During dry storage, relative fresh weight loss occurred in *Solidago x hybrida* cv 'Golden Glory' flowers

compared to the first day (Figure 7). These relative fresh weight losses were found to be statistically significant ($p \leq 0.01$).




Figure 7. The effects of storage periods on relative fresh weight of *Solidago x hybrida* cv 'Golden Glory'. * The difference between the means with different letters is significant ($p \leq 0.01$).

Relative fresh weight measured as 100% before storage were measured

as 95% on the 5th day, 88% on the 10th day and 84% on the 15th day. As the

storage period increased, the relative fresh weight of the flowers decreased. The resulting relative fresh weight loss can be explained by the flowers losing water through respiration and transpiration. In parallel with this study, it was reported that the relative fresh weight of the flowers decreases as the dry storage period of cut flowers increases (Joyce et al., 2000; Makwana et al., 2015).

Correlation data of the features examined in the study are presented in

Table 2. The relationship between storage periods, vase life, relative fresh weight and chlorophyll content was significant ($p \leq 0.01$). A negative correlation was found between vase life, relative fresh weight and chlorophyll content as the storage periods increased. A positive correlation was found between the decrease in relative fresh weight and chlorophyll content and vase life.

Table 2. Correlation results of storage periods, vase life, relative fresh weight and chlorophyll content in *Solidago x hybrida* cv 'Golden Glory'.

		Storage Period	Vase Life	Relative fresh weight	Chlorophyll Content
Storage Period	Pearson Correlation	1	-,967**	-,934**	-,887**
	Sig. (2-tailed)		,000	,000	,000
	N	60	60	60	60
Vase Life	Pearson Correlation	-,967**	1	,898**	,843**
	Sig. (2-tailed)	,000		,000	,000
	N	60	60	60	60
Relative fresh weight	Pearson Correlation	-,934**	,898**	1	,824**
	Sig. (2-tailed)	,000	,000		,000
	N	60	60	60	60
Chlorophyll Content	Pearson Correlation	-,887**	,843**	,824**	1
	Sig. (2-tailed)	,000	,000	,000	
	N	60	60	60	60

** . The correlation is significant at the $p \leq 0.01$ level.

CONCLUSIONS

In this study, it was aimed to determine the effects of different storage periods on the vase life of *Solidago x hybrida* cv 'Golden Glory'. The longest vase life was determined at no-stored flowers. Extending the storage period shortened the vase life and damaged the leaves. The leaf yellowing rather than the flowers in the spikes was effective in ending the vase life of the cut *Solidago x hybrida* cv 'Golden Glory'. The results showed that cut goldenrod flowers can be stored for up to 5 days. However, for longer vase life and storage, new studies should be performed with applications to

prevent leaf yellowing, different temperatures and storage conditions.

REFERENCES

- Anonymous. 2008. The project of strengthening the vocational education and training system. Solidago breeding publication, 39., Ankara.
- Ahmad, Iftikhar, et al. 2012. Dry storage effects on postharvest performance of selected cut flowers. HortTechnology 22.4 463-469.
- Armitage, A.M., Laushman, J.M. 2003. Specialty cut flowers: The production of annuals, perennials, bulbs, and woody plants for fresh

- and dried cut flowers. Portland, OR: Timber Press.
- Christiansen, M.W. and Gregersen, P.L. 2014. Members of the barley NAC transcription factor gene family show differential co-ration with senescence-associated genes during senescence of flag leaves. *J. Exp. Bot.*, 65, 4009-4022.
- Çelikel, F. G., Reid, M. S. 2002. Storage temperature affects the quality of cut flowers from the Asteraceae. *HortScience*, 37(1), 148-150.
- Çelikel, F. 2013. Süs Bitkilerinin Hasat Sonrası Kaliteleri ve Yeni Teknolojiler. V. Ulusal Süs Bitkileri Kongresi, pp:17-26, 06-09 May 2013, Yalova.
- Çelikel, F. 2020. Postharvest Quality and Technology of Cut Flowers and Ornamental Plants. *Black Sea Journal of Agriculture*, Vol 3, (3): 225-232.
- Ergür, E.G., Kazaz, S., Kılıç, T., 2016. An Indispensable Element of Bouquet and Floral Arrangements: Cut Foliage. VI. Ornamental Plants Congress, 19-22 April 2016, pp:335-346, Antalya.
- Fanourakis, D., Pieruschka, R., Savvides, A., Macnish, A. J., Sarlikioti, V., Woltering, E. J 2013. Sources of vase life variation in cut roses: a review. *Postharvest Biology and Technology*, 78, 1-15.
- Ferrante, A., Vernieri, P., Serra, G., & Tognoni, F. 2004. Changes in abscisic acid during leaf yellowing of cut stock flowers. *Plant growth regulation*, 43(2), 127-134.
- Ferrante, A., Alberici, A., Antonacci, S., Serra, G. 2007. Effect of promoter and inhibitors of phenylalanine ammonia-lyase enzyme on stem bending of cut gerbera flowers. *ActaHorticulturae*, 755, 471-476.
- Hassan, F. A. S., Tar, T and Zs Dorogi. 2003. Extending the vase life of *Solidago canadensis* cut flowers by using different chemical treatments. *International Journal of Horticultural Science* 9.2 83-86.
- Halevy, A.H., Mayak, S. 1981. Senescence and postharvest physiology of cut flowers. Avi Publishing Company Inc. Westport, Connecticut, 3, 59-143.
- He, Y.J. and Gan, S. 2002. A Gene Encoding An Acyl Hydrolase Is Involved in Leaf Senescence in Arabidopsis. *Plant Cell*, 14, 805-815.
- He, S., Joyce, D.C., Irving, D.E., and Faragher, J.D. 2006. Stem end blockage in cut *Grevillea* ‘Crimson Yul-lo’ inflorescences. *Postharvest Biol. Technol.* 41 (1), 78–84
- Jones, R. and Moody, H. 1993. Caring for cut flowers. Agmedia, St. Kilda, Australia.
- Jones, R., & Faragher, J. 1991. Cold storage of selected members of the Proteaceae and Australian native cut flowers. *HortScience*, 26(11), 1395-1397.
- Joyce, D. C., Meara, S. A., Hetherington, S. E., Jones, P. 2000. Effects of cold storage on cut *Grevillea* ‘Sylvia’ inflorescences. *Postharvest Biology and Technology*, 18(1), 49-56.
- Kazaz, S., Askin, M.A., Tekintas, F.E. 2003. Kesme Çiçeklerde Hasat Sonrası Ömrü Arttıran Uygulamalar. IV. Ulusal Bahçe Bitkileri Congress, 8-12 September 2003, p: 519-522, Antalya.
- Kazaz, S., Karagüzel, O. 2010. Influence of Growth Regulators on the Growth and Flowering Characteristics of Goldenrod (*Solidago x hybrida*). *European Journal of Scientific Research* 45(3): 498-507.
- Kazaz, S., 2012. Türkiye'nin Doğal Bitki Zenginliği. *Çiçek Vizyon Magazine*, Issue:55, Year: 6 January-February, 24-25.p.
- Looze, T., van Staaveren, J. 2003. Post harvest treatment of cut flowers. <http://simonbw.lecture.ub.ac.id/files/2010/03/Posth-Cut-Flower-2.pdf>. Date of access: 18.12.2016.

- Lü, P., Cao, J., He, S., Liu, J., Li, H., Cheng G., Ding Y., Joyce, D.C. 2010. Nanosilver pulsetreatments improve water relations of cutrose cv. Movie star flowers. *Postharvest Biology and Technology*, 57, 196-202.
- Mabini, N. Q. ve Acedo, V. Z., 2013. Vase Life of Selected Florist Greens in Different Holding Solutions with Commercial Preservatives. Department of Horticulture, College of Agriculture and Food Science. Visayas State University. Bybay, Leyte-Philippines, 79-82.
- Macnish, A.J., M.S. Reid, and D.C. Joyce. 2009. Ornamentals and cut flowers, In: E.M. Yahia (ed.). *Modified and controlled atmospheres for the storage, transportation, and packaging of horticultural commodities*. Taylor & Francis Group, Boca Raton, FL. 491–506.
- Makwana, R. J., Singh, A., and Neelima, P. 2015. Effect of cold storage techniques on flower quality and vase life of rose var. 'Sun King'. *The Bioscan*, 10(1), 225-227.
- Nowak, J., Rudnicki, R.M. 1990. Postharvest handling and storage of cut flower. *Florist Green and Potted Plants* Timber Press. Inc., Singapore, 29-64.
- Özzambak, E., 2009. Kesme Yeşillik Yetiştiriciliği. *Çiçek Vizyon Magazine*, Year 4, No.34.
- Penfold, C.A. and Buchanan-Wollaston, V. 2014. Modelling transcriptional networks in leaf senescence. *J. Exp. Bot.*, 65, 3859-3873.
- Philosoph Hadas, Sonia, et al. 1996. Benzyladenine pulsing retards leaf yellowing and improves quality of goldenrod (*Solidago canadensis*) cut flowers. *Postharvest Biology and Technology* 9.1 (1996): 65-73.
- Reid, M.S. 2002. Postharvest handling systems: Ornamental crops, p. 315–325. In: A.A. Kader ed.). *Postharvest technology of horticultural crops*. University of California, Oakland.
- Sacalis, J.N. 1993. Cut flowers: Prolonging freshness, p. 52–54. In: J.L. Seals. (ed.). *Post production care and handling*. 2nd ed. Ball Publ., Batavia, IL.
- Sağlam, N. G. 2015. Leaf Senescence: A View of Its Physiological and Molecular Regulation. *Marmara Journal of Science*, 27(3), 83-92.
- Schiappacasse, F., Moggia, C., Contreras, R. 2014. Studies with long term storage of cut flowers of *Hydrangea macrophylla*. *Idesia*, 32(4), 71-76.
- Shiva KN, Bhattacharjee SK. 2006. Effect of pulsing and wet storage in carbohydrates during the course of senescence in cut Rose. *Indian Journal of Horticulture*, 69(4):419-423.
- Teixeira da Silva, J.A. The cut flower: Postharvest considerations. *J. Biol. Sci.* 2003, 3, 406–442.
- Thimann, K. V. 1985. The interaction of hormonal and environmental factors in leaf senescence. *Biologia plantarum*, 27(2), 83-91.
- Thomas, H. and Stoddart, J.L., 1980. Leaf senescence. *Annu. Rev. Plant Physiol.*, 3 I: 83-111
- Ueyama, S., Ichimura, K. 1998. Effects of 2-hydroxy-3-ionene chloride polymer on the vase life of cut rose flowers. *Postharvest Biology and Technology*, 14, 65-70.
- Van Doorn, W.G. 1997. Water relations of cut flowers. *Horticultural Review*, 18, 1-85.
- Woltering, E. 2008. Quality and Logistics of Horticultural Products. <http://www.horticonsult.nl/wp-content/uploads/2011/05/eu-wur-turkey-2008.pdf>. Date of access: 20.1.2020.