

Pakistan Journal of Advances in Medicine and Medical Research

ISSN 2957-5893 (Online)

Vol 01- Issue 01-April-2022



Obstetric Outcomes Of Teenage Pregnancy Vs. Non Teenage Pregnancy Patients. A Multi-Center Study. OPEN ACCESS

Lubna Tahir, Aysha Bibi, Kiran Jamshed, Sumaria

Yasmin, Seema Amin, Nuzhat Amin  OPEN ACCESS

Platform &
workflow by
OJS / PKP

PKP
PUBLIC
KNOWLEDGE
PROJECT

OJS
OPEN JOURNAL SYSTEMS

www.pjammr.com

DOI: [10.213347/PJAMMR.2022.vol-01-.022](https://doi.org/10.213347/PJAMMR.2022.vol-01-.022)

| Received: 26.04.2022 | Accepted: 28.04.2022 | Published: 11.05.2022

Copyright © 2022 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 International License (PJAMMR-CC BY-NC 3.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

ISSN 2957-5893 (Online)

Obstetric Outcomes Of Teenage Pregnancy Vs. Non Teenage Pregnancy Patients. A Multi-Center Study .

¹ Lubna tahir , ² Aysha bibi , ³ Kiran jamshed , ⁴ Sumaria Yasmin , ⁵ Seema Amin ,

⁶ Nuzhat Amin

¹. Senior District Gynecologist D.H.Q Hospital Timarghra Lower Dir , Pakistan

². Senior Registrar Department of Obstetrics and Gynecology, Saidu Teaching Hospital Swat. Pakistan

³. Woman Medical officer saidu teaching hospital swat , Pakistan

⁴. Assistant Professor obstetrics and gynecology MTI LRH hospital Peshawar , Pakistan

⁵ Senior Registrar Department of Obstetrics and Gynaecology, Saidu Teaching Hospital Swat. Pakistan

⁶ Professor of department of obstetrics and gynecology MMC hospital maradan , Pakistan

Corresponding Author: Dr. AYSHA BIBI

Email: drrehmat123@hotmail.com

ABSTRACT

Original Research Article

OBJECTIVE: The Goal Of This Study Was To Examine The Obstetric Outcomes Of Teenage And Non-Teenage Pregnancies.

MATERIAL AND METHODS: A Multi-Center Study Was Conducted In Kpk, Pakistan, From March 2021 To February 2022. All 14-18-Year-Olds Who Gave Birth In The Two Hospitals Had Obstetric Outcomes Compared To Non-Adolescent Women (19-36). Chi-Square And Students' T-Tests Were Used With A 0.05 Significance Level.

RESULTS: Teenage Mothers Were More Likely To Have Severe Anemia (7.5 Vs. 4.1%) And Chorioamnionitis (2.6 Vs. 0.6%) Than Non-Teenage Mothers, And Their Infants Were More Likely To Have Post Maturity (4.6 Vs. 1.8%) And Meconium Aspiration Syndrome (6.5 Vs. 2.4%). Teenagers Were Less Likely To Be Overweight Than Adults. Teens Are More Likely To Have Instrumental Births (7.1% Vs. 2.2%, P 0.01). Preterm Delivery, Low Birth Weight Infant, RDS, And Fetal And Perinatal Death Were Not Significantly Different Between The Two Groups.

CONCLUSION: Anemia And Chorioamnionitis Are More Common In Teenage Mothers. Teenagers Are More Likely To Use An Instrument. Adolescent Moms Often Have Babies With Post-Maturity And Meconium Aspiration Syndrome.

KEYWORDS: Obstetric, Outcomes, Teenage, Non-Teenage, Pregnancy, A Multi-Center Study, Pakistan

INTRODUCTION

Economic Survey of Pakistan 2019- 21 reports 184.35 million people and 2% growth. Pakistan will rank 5th in population by 2050. 39% are kids. Many females marry as young as 7 years old. In 2001, 37% of women married before 18; the same is feared today. In 2012, the media reported 75 child marriages; 43% were 11-15, and 32% were 6-10. 10-14-year-olds and 15-19-year-olds married in 2008- 2009. Minors marry without supervision. In 2006-07, over half of the teenage females in Pakistan were pregnant. Child and early marriage numbers are uncertain due to a lack of data. Since 1998, Pakistan hasn't done a marriage and age census. Pakistan's child marriages have several reasons. Weak rules, lack of enforcement, children being regarded as commodities/enslaved people, a tribal and feudal social structure, a lack of public knowledge about the bad effects of child marriages, severe poverty, internal trafficking, and a lack of government will are prominent causes. Ineffective birth registration contributes to child marriage. Never prioritizing female birth registration permits manipulating marital ages. No centralized, unbiased, or influential child rights organization monitors child marriages.

Pakistan's child brides and Non-

Muslim child marriages plague Pakistan. Religious minorities conduct and record marriages regardless of children's ages. The previous government lobbied coalition partners to pass the Hindu Marriage Bill 2011, which barred non-Muslim under-18 marriages. Kidnapping and exploitation "Parents sell or pressure children into marriage, forced labor, sexual exploitation, or domestic slavery, " except Sindh bans internal trafficking. Human trafficking is illegal under Pakistan's Sindh Child Protection Authority Act 2012. Early marriage causes Multiple factors to trigger Pakistani child marriages. Lack of public information on child marriage's negative effects, extreme poverty, and trafficking are major factors. Unreliable birth records cause child marriage. Never prioritizing birth registration for children, particularly females, allows age manipulation upon marriage. No central, impartial, or strong child rights authority oversees child weddings.

Child marriages in Pakistan: There is no legal structure for weddings or non-Muslim child marriages in Pakistan. Regardless of age, all religious minority groups conduct and register marriages according to their religion. The previous government lobbied coalition partners to pass the Hindu Marriage

Bill 2011, which barred non-Muslim under-18 marriages. Kidnapping, exploitation

Child marriages, forced labor, sexual exploitation, and domestic slavery are common. All except Sindh ban internal trafficking. This law prohibits human trafficking in Sindh. 14,15
Methods

The research was conducted at the hospital timer Gira-lower dir and saidu teaching hospital swat Pakistan. Two ERBs approved the study.

13-19-year-old singletons were included. A control was a 20-35-year-old lady who delivered a singleton on the same day as the Multi-center case study. Hypertensive, diabetic, cardiac, renal, endocrine, or autoimmune

women were eliminated. Anemia, hypertension, gestational diabetes, poly/oligohydramnios, antepartum/postpartum hemorrhage, ruptured uterus, and chorioamnionitis were compared. Low birth

weight, congenital anomalies, meconium aspiration syndrome, RDS, neonatal hospitalization, and perinatal death. Inclusion-eligible women got counseling and approval. Measuring women showed

their ethnicity, education, and prenatal care. Medical data showed maternal and fetal outcomes.

A questionnaire collected mom, baby, and mom data.

When anemia, low birth weight,

malnutrition, obesity, low birth babies, preterm babies, fetal distress, meconium aspiration, illiteracy, antenatal care, etc., were taken in the range of 8 percent to 86 percent, the anticipated odds ratio for teenagers more than 5 percent level of significance. Input by two operators into SPSS (version 2.4). SPSS generated variable frequencies. The survey compared fetus and mother results. Statistical significance was determined using Chi-square and Student's t. 0.05

METHODS: A multi-center study was conducted in Kpk, Pakistan, from March 2021 to February 2022. All 14-18-year-olds who gave birth in the two hospitals had obstetric outcomes compared to non-adolescent women (19-36). Chi-square and students' t-tests were used with a 0.05 significance level.

RESULTS

There were 910 deliveries at the two hospitals during the study period, with n=128 (86%) being adolescent pregnancies. The total number of completed surveys from the two hospitals was 910 Lower dir 460 and STH-Swat 450, with 128 teenage pregnancies and 782 non-teenage pregnancies used as controls. The study only included married women. The mean ages of teen and non-teen mothers were 16.30.6 and 24.43.2 years. Majority (n=60; 48%) of teenage mothers were 17 years old, followed by 18 years (n=48; 32.2%). n=20 (9.7%) were 16 years old, with none younger. Only 52% of teen mothers were first-time moms. More than 28% had a second. Two hospitals' cases and distribution controls are

Table-1: Cases And Distribution Controls Included From Two Hospitals.

	LOWER DIR N (%)	STH-SWAT N (%)	TOTAL N=128 (%)
TEENAGE	60 (48%)	68 (52%)	128 (100)
CONTROLS	350 (42%)	432 (44%)	782 (86%)
TOTAL	460 (42%)	450 (38%)	910 (80%)

Table-2: Maternal Complications In Teenage And Non-Teenage Mothers Are Compared.

VARIABLE	CASES n (%)	CONTROLS n (%)	P- VALU E
ANAEMIA (HB < 11 GM/DL)	90 (58.0)	632 (44%)	NS
SEVERE ANAEMIA (HB < 7GM/DL)	04 (8.0)	39 (4.3)	0.03
PRETERM PREMATURE RUPTURE OF MEMBRANES (PPROM)	6 (1.1)	8 (0.9)	NS
HYPERTENSIVE DISORDERS OF PREGNANCY	18 (10. 2)	64 (7.0)	NS
GESTATIONAL DIABETES	2 (0.6)	1 (0.1)	NS
OLIGOHYDRAMNIOS	1 (0.6)	7 (0.8)	NS
POLYHYDRAMNIOS	0 (0.0)	4 (0.4)	NS
ANTEPARTUM HAEMORRHAGE	1 (0.6)	10 (1.1)	NS
POSTPARTUM HAEMORRHAGE	1 (0.6)	6 (0.7)	NS
CHORIOAMNIONITIS	5 (2.8)	7 (0.8)	0.01
MATERNAL MORTALITY	0	1 (0.1)	NS
HOSPITAL STAY IN DAYS, (MEAN±SD)	1.6±2.2	1.9±2.3	NS

(NS: Non-significant)

Table-3: Fetal and neonatal complications in teenage and non-teenage mothers are compared.

VARIABLE	CASES n (%)	CONTROLS n (%)	P VALUE
LOW BIRTH WEIGHT (< 2500 GM)	28 (20.5)	141 (18.2)	NS
PRETERM	22 (7.7)	47 (5.9)	NS
POST-TERM	7 (4.6)	14 (1.8)	0.02
FOETAL DISTRESS	14 (9.3)	55 (7)	NS
MECONIUM ASPIRATION SYNDROME	10 (6.5)	19 (2.4)	< 0.01
RESPIRATORY DISTRESS SYNDROME	2 (1.3)	18 (2.3)	NS
ADMISSION TO NEONATAL NURSERY	11 (7.1)	57 (7.3)	NS
CONGENITAL ANOMALIES	1 (0.6)	18 (2.3)	NS
STILL BIRTH	10 (6.5)	66 (8.3)	NS
NEONATAL DEATH IN HOSPITAL	1 (0.7)	11 (1.4)	NS

About 17% were having their third or more baby. Teenage moms had a lower mean BMI than non-teenage mothers (20.2 vs. 22.6, p 0.002). While the difference was not statistically significant, teens had greater literacy and prenatal care uptake than adults.

Groups' maternal problems are compared in Table-2. The majority of moms in both categories were lifeless, but severe anemia was twice as likely among pregnant teens. Chorioamnionitis is twice as common in teenagers. The two groups had similar pregnancy difficulties. Normal vaginal birth rates were similar, while cesarean section rates were lower for teens. This change was non-significant. Among teens, instrumental delivery was more common.

Table 3 illustrates fetal and neonatal outcomes for both groups. Post-term infants are more common in teenage pregnancies than non-teenage pregnancies (p<0.05). Meconium aspiration syndrome was twice as common in cases vs. controls. Other newborn problems

did not show a significant difference in incidence.

DISCUSSION

This multi-center research assessed maternal and fetal risks. This multi-center case study found maternal and fetal hazards of adolescent pregnancy. This may be Pakistan's first multi-university research. Our survey revealed 4.6 percent of adolescent moms in lower dir and swat. 9 percent of adolescent pregnancy in Pakistan

According to studies, all parents in both groups were married, so pregnant teenagers had social support. Teen pregnancy is most common in Kpk (26/128, 24%) and Punjab (28%). Women from Pakistan and Sindh made up about 18% of the group. In illiteracy and prenatal care, teens fared better than adults. This contradicts data on teen moms' literacy and prenatal care. 14% of 128 pregnant teens were in their third or more pregnancy, showing contraceptive ignorance. 18

Maybe our women's nutrition. Teenage moms had greater anemia (7 g/dL). Their immaturity may be owing to fetal growth and resource competition. Studies agree.

Chorioamnionitis may affect adolescent moms. Another research agrees. Unfathomable. Anatomical variations may exacerbate chorioamnionitis.

Puberty is unfinished. The alkalinity of the prepubertal vagina may favor bacterial vaginosis, while the perimenarcheal cervix's large squamocolumnar junction and short cervix may facilitate organisms' ascent to the uterus.

All additional obstetric problems, including hypertensive disorders, gestational diabetes, preterm premature membrane rupture, poly/oligohydramnios, antepartum hemorrhage, and post-partum hemorrhage, were identical among groups. According to Pakistani research, teenage moms are more prone to pregnancy-induced hypertension and puerperal sepsis. 20

Adolescents had more instrumental births and fewer caesareans. Both groups' vaginal birth rates were comparable. Teenage moms' delivery technique is controversial. Some studies reveal that teens had higher rates of normal vaginal birth and lower cesarean section and instrumental delivery rates. Israeli research found that teens had more instrumental deliveries and fewer caesareans than adult women. Under-15s had a greater maternal mortality risk. We discovered 0 teen maternal fatalities and one adult maternal death. 21

Our study found no higher risk of unfavorable perinatal outcomes for teen mothers, save meconium aspiration syndrome and post-maturity. Study-to-study perinatal outcomes

vary. Preterm delivery, low birth weight, and fetal or perinatal mortality are linked to adolescent pregnancy. Other studies revealed no dangers. No research shows a greater rate of post-term births among teens. 22

From 2015 through 2021, many teenage age groups were included in the research, causing varying obstetric findings. Younger teens had average results, according to studies. We couldn't compare their pregnancy outcomes since they were all 16-18. 4 under-17 females and none under-16 Multiparas were also studied. Different birthing systems may help. In many prosperous nations, pregnant adolescents are unmarried, poor immigrants. Inadequate sex education, early first intercourse, low contraceptive awareness, use, poor home circumstances, and high school dropout rates cause teen pregnancy. 24 teenage pregnancy studies come from free-treatment countries.

Teenage pregnancy is widespread in Pakistan families with social support. Youth and poverty restrict their decision-making capacity. You grow up, not reproduce in your teens. Forcing a girl out of school and into marriage might be harmful. Lack of contraceptive information causes physical and mental suffering for many young women. Our study indicated no higher risk for young women due to psychological stress.

The sample size was limited, and the women were predominantly 18-19—unanalyzed teen pregnancy findings. Most tertiary hospitals utilize traditional birth attendants, which may lower teen births. These conditions may harm both mother and fetus. Undocumented teen pregnancies. Larger-scale Teen pregnancy research is essential. 25

CONCLUSION

Teen Pregnancy Is More Likely To Develop Severe Anemia And Chorioamnionitis And Have An Instrumental Birth. Teenage Pregnancies Cause Post-Maturity And Meconium Aspiration Syndrome.

REFERENCES

1. UNICEF, "YOUNG PEOPLE AND FAMILY PLANNING: TEENAGE PREGNANCY," [HTTP://WWW.UNICEF.ORG/MALAYSIA/TEENAGE_PREGNANCIES_-_OVERVIEW.PDF](http://www.unicef.org/malaysia/teenage_pregnancies_-_overview.pdf). VIEW AT [GOOGLE SCHOLAR](#)
2. S. NEAL, Z. MATTHEWS, M. FROST, H. FOGSTAD, A. V. CAMACHO, AND L. LASKI, "CHILDBEARING IN ADOLESCENTS AGED 12–15 YEARS IN LOW RESOURCE COUNTRIES: A NEGLECTED ISSUE. NEW ESTIMATES FROM DEMOGRAPHIC AND HOUSEHOLD SURVEYS IN 42 COUNTRIES," *ACTA OBSTETRICIA ET GYNECOLOGICA SCANDINAVICA*, VOL. 91, NO. 9, PP. 1114–1118, 2012. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)
3. M. WHITWORTH AND R. COCKERILL, OWOEYE GANI, "SOCIO-DEMOGRAPHIC DETERMINANTS OF TEENAGE PREGNANCY IN THE NIGER DELTA OF NIGERIA," *OPEN JOURNAL OF OBSTETRICS AND GYNECOLOGY*, VOL. 2, NO. 3, PP. 239–243, 2012. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)
5. H. E. KULIN, "ADOLESCENT PREGNANCY IN AFRICA: A PROGRAMMATIC FOCUS," *SOCIAL SCIENCE & MEDICINE*, VOL. 26, NO. 7, PP. 727–735, 1988. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)
6. M. IMAMURA, J. TUCKER, P. HANNAFORD, ET AL., "FACTORS ASSOCIATED WITH TEENAGE PREGNANCY IN THE EUROPEAN UNION COUNTRIES: A SYSTEMATIC REVIEW," 2007. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)
7. M. WERE, "DETERMINANTS OF TEENAGE PREGNANCIES: THE CASE OF BUSIA DISTRICT IN KENYA," *ECONOMICS & HUMAN BIOLOGY*, VOL. 5, NO. 2, PP. 322–339, 2007. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)
8. M. SHAW, D. A. LAWLOR, AND J. M.

“ANTENATAL MANAGEMENT OF TEENAGE PREGNANCY,” *OBSTETRICS, GYNAECOLOGY & REPRODUCTIVE MEDICINE*, VOL. 20, NO.11, PP. 323–328, 2010.VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

4. A. IBRAHIM ISA AND I. OLUGBENGANAJMAN, “TEENAGE CHILDREN OF TEENAGE MOTHERS: PSYCHOLOGICAL, BEHAVIOURAL AND HEALTH OUTCOMES FROM AN AUSTRALIAN PROSPECTIVE LONGITUDINAL STUDY,” *SOCIAL SCIENCE & MEDICINE*, VOL. 62, NO. 10, PP. 2526–2539, 2006.VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

9. P. KESKINOGLU, N. BILGIC, M. PICA KCIEFE, H. GIRAY, N. KARAKUS, AND T. GUNAY, “PERINATAL OUTCOMES AND RISK FACTORS OF TURKISH ADOLESCENT MOTHERS,” *JOURNAL OF PEDIATRIC AND ADOLESCENT GYNECOLOGY*, VOL. 20, NO. 1, PP. 19–24, 2007.VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

10. B. S. ZUCKERMAN, D. K. WALLER, D. A. FRANK, C. CHASE, AND B. HAMBURG, “ADOLESCENT PREGNANCY: BIOBEHAVIORAL DETERMINANTS OF OUTCOME,” *THE JOURNAL OF PEDIATRICS*, VOL. 105, NO. 6, PP. 857–863, 1984.VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

11. N. GUPTA, U. KIRAN, AND K. BHAL, “TEENAGE PREGNANCIES: OBSTETRIC CHARACTERISTICS AND OUTCOME,” *EUROPEAN JOURNAL OF OBSTETRICS & GYNECOLOGY AND REPRODUCTIVE BIOLOGY*, VOL. 137, NO.2, PP. 165–171, 2008.VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

12. A. KURUP, O. VIEGAS, K. SINGH, AND S. S.

AMERICAN JOURNAL OF OBSTETRICS AND GYNECOLOGY, VOL. 194, NO. 3, PP. 840–845, 2006.VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

14. A. BACCI, G. M. MANHICA, F. MACHUNGO, A. BUGALHO, AND M. CUTTING, “OUTCOME OF TEENAGE PREGNANCY IN MAPUTO, MOZAMBIQUE,” *INTERNATIONAL JOURNAL OF GYNECOLOGY & OBSTETRICS*, VOL. 40, NO. 1, PP. 19–23, 1993.VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

15. O. BUKULMEZ AND O. DEREN, “PERINATAL OUTCOME IN ADOLESCENT PREGNANCIES: A CASE-CONTROL STUDY FROM A TURKISH UNIVERSITY HOSPITAL,” *EUROPEAN JOURNAL OF OBSTETRICS & GYNECOLOGY AND REPRODUCTIVE BIOLOGY*, VOL. 88, NO. 2, PP. 207–212, 2000.VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

16. D. UKIL AND U. I. ESEN, “EARLY TEENAGE PREGNANCY OUTCOME: A COMPARISON BETWEEN A STANDARD AND A DEDICATED TEENAGE

RATNAM, "PREGNANCY OUTCOME IN UNMARRIED TEENAGE NULLIGRAVIDAE IN SINGAPORE," *INTERNATIONAL JOURNAL OF GYNECOLOGY & OBSTETRICS*, VOL. 30, NO. 4, PP. 305– 311, 1989. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

13. B. T. BATEMAN AND L. L. SIMPSON, "HIGHER RATE OF STILLBIRTH AT THE EXTREMES OF REPRODUCTIVE AGE: A LARGE NATIONWIDE SAMPLE OF DELIVERIES IN THE UNITED STATES," *ANTENATAL CLINIC*," *JOURNAL OF OBSTETRICS AND GYNAECOLOGY*, VOL. 22, NO. 3, PP. 270– 272, 2002. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

17. R. T. MIKOLAJCZYK, J. ZHANG, A. P. BETRAN, ET AL.,

18. "A GLOBAL REFERENCE FOR FETAL- WEIGHT AND BIRTHWEIGHT PERCENTILES," *THE LANCET*, VOL. 377,

NO. 9780, PP.1855–1861, 2011. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

19. D. J. MURPHY, "EPIDEMIOLOGY AND ENVIRONMENTAL FACTORS IN PRETERM LABOUR," *BEST PRACTICE & RESEARCH CLINICAL OBSTETRICS & GYNAECOLOGY*, VOL. 21, NO. 5, PP. 773– 789, 2007. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

20.-C. LO, J.-J. HSU, C.-C. HSIEH, T. T. HSIEH, AND T.-H. HUNG, "RISK FACTORS FOR 24 *TAIWANESE JOURNAL OF OBSTETRICS AND GYNECOLOGY*, VOL. 46, NO. 4, PP. 389– 394, 2007. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLARS](#). W. WEN, G. SMITH, Q. YANG, AND

21. P. CHANTRAPANICHKUL AND S. CHAWANPAIBOON, "ADVERSE PREGNANCY OUTCOMES IN CASES INVOLVING EXTREMELY YOUNG MATERNAL AGE," *INTERNATIONAL JOURNAL OF GYNECOLOGY & OBSTETRICS*, VOL. 120, NO. 2, PP. 160–164, 2013. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

M. WALKER, "EPIDEMIOLOGY OF PRETERM BIRTH AND NEONATAL OUTCOME," *SEMINARS IN FETAL AND NEONATAL MEDICINE*, VOL. 9, NO. 6, PP. 429– 435, 2004. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

24. G. C. DI RENZO, I. GIARDINA, A. ROSATI, ET AL., "MATERNAL RISK FACTORS FOR PRETERM BIRTH: A COUNTRY- BASED POPULATION ANALYSIS," *EUROPEAN JOURNAL OF OBSTETRICS & GYNECOLOGY AND REPRODUCTIVE BIOLOGY*, VOL. 159, NO. 2, PP. 342–346, 2011. VIEW AT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

25.C.-C. LO, J.-J. HSU, C.-C. HSIEH, T. T. HSIEH, AND T.-H. HUNG, "RISK FACTORS FOR SPONTANEOUS PRETERM DELIVERY BEFORE 34 WEEKS OF GESTATION AMONG TAIWANESE

WOMEN," *TAIWANESE JOURNAL OF OBSTETRICS AND GYNECOLOGY*, VOL. 46, NO. 4, PP. 389–394, 2007. VIEWAT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)

22.S. THAITHAE AND R. THATO, "OBSTETRIC AND PERINATAL OUTCOMES OF TEENAGE

PREGNANCIES IN THAILAND," *JOURNAL OF PEDIATRIC AND ADOLESCENT GYNECOLOGY*, VOL. 24, NO. 6, PP. 342–346, 2011. VIEWAT: [PUBLISHER SITE](#) | [GOOGLE SCHOLAR](#)