



Antibody Characterization Report for TYRO protein tyrosine kinase-binding protein (TYROBP)

YCharOS Antibody Characterization Report

Author(s): Walaa Alshafie¹, Maryam Fotouhi¹, Riham Ayoubi¹, Peter S. McPherson^{1*} and Carl Laflamme^{1*}

¹ Tanenbaum Open Science Institute, Structural Genomics Consortium, Montreal Neurological Institute, McGill University, Montreal, Canada

* Corresponding authors: carl.laflamme@mcgill.ca, peter.mcpherson@mcgill.ca

Target:

Recommended protein name: TYRO protein tyrosine kinase-binding protein

Alternative protein name: DNAX-activation protein 12, DAP12

Gene name: TYROBP

Uniprot: O43914

We are a third-party organization with the mission to characterize commercial antibodies for all human protein through open science [1]. This report guides researchers to select the most appropriate antibodies for TYRO protein tyrosine kinase-binding protein. We used an antibody characterization pipeline [2] based on knockout (KO) cells to perform head-to-head comparisons of commercial antibodies for TYRO protein tyrosine kinase-binding protein by immunoblot (Western blot), immunoprecipitation and immunofluorescence. THP-1 was selected based on evidence of appropriate TYRO protein tyrosine kinase-binding protein expression [3, 4]. A THP-1 *TYROBP* KO line is available at Abcam and was used in this study.

The authors do not provide an assessment of the quality of the tested antibodies as their respective performances are limited to our finite experimental conditions. The readers should interpret the present findings based on their own scientific expertise. The authors acknowledge that an antibody that demonstrates specificity in the stated test conditions can be suboptimal in a different experimental format or in cell lines that differ from those directly tested here.

Table 1: Summary of the TYRO protein tyrosine kinase-binding protein antibodies tested

Company	Catalog number	Lot number	RRID (Antibody Registry)	Clonality	Clone ID	Host	Concentration (µg/µl)	Vendors recommended applications
Bio-Techne	MAB5240*	CARS0620011	AB_95182	monoclonal	406288	mouse	0.50	Wb
Abcam	ab93846	GR3257775-1	AB_10865289	polyclonal	-	rabbit	1.00	Wb, IF
Abcam	ab155779	GR131013-13	AB_2885095	polyclonal	-	rabbit	0.65	Wb
Abcam	ab124834**	GR104841-12	AB_10971363	recombinant-mono	EPR5173	rabbit	0.72	Wb, IF
Thermo Fisher Scientific	MA5-34787**	WJ3417748D	AB_2848695	recombinant-mono	JG38-70	rabbit	1.00	Wb
Thermo Fisher Scientific	PA5-26861	WJ3414154	AB_2544361	polyclonal	-	rabbit	0.40	Wb
Cell Signaling Technology	97415**	1	AB_2916164	recombinant-mono	E7U7T	rabbit	0.10	Wb, IP, IF
Cell Signaling Technology	46088**	1	AB_2916165	recombinant-mono	E8P9U	rabbit	0.48	other
Cell Signaling Technology	12492**	1	AB_2721120	recombinant-mono	D7G1X	rabbit	0.048	Wb, IP

Wb=Western blot, IP= immunoprecipitation, IF=immunofluorescence, *=monoclonal antibody, **=recombinant antibody

Table 2: Summary of the cell lines used

Institution	Catalog number	RRID (Cellosaurus)	Cell line	genotype
Abcam	ab271147	CVCL_0006	THP-1	WT
Abcam	ab273726	CVCL_B7TG	THP-1	TYROBP KO

Figure 1: TYRO protein tyrosine kinase-binding protein antibody screening by immunoblot.

Lysates of THP-1 WT and *TYROBP* KO were prepared, and 100 µg of protein were processed for immunoblot with the indicated TYRO protein tyrosine kinase-binding protein antibodies. The Ponceau stained transfers of each blot are shown. Antibody dilution used: MAB5240* at 1/250, ab93846 at 1/1000, ab155779 at 1/500, ab124834** at 1/2000, MA5-34787** at 1/500, PA5-26861 at 1/1000, 97415** at 1/1000, 46088** at 1/500, 12492** at 1/1000. Predicted band size: 12 kDa. *=monoclonal antibody, **=recombinant antibody

Figure 2: TYRO protein tyrosine kinase-binding protein antibody screening by immunoprecipitation.

A) THP-1 lysates were prepared, and immunoprecipitation was performed using 1.0 µg of the indicated TYRO protein tyrosine kinase-binding protein antibodies pre-coupled to Dynabeads protein G or protein A. Ability of the antibodies to capture TYRO protein tyrosine kinase-binding protein was first assessed by comparing the level of TYRO protein tyrosine kinase-binding protein from the starting material to the unbound fractions. **B)** Analysis of the immunoprecipitate for the antibody that showed depletion of TYRO protein tyrosine kinase-binding protein in (A). For immunoblot, 12492** was used at 1/1000. The Ponceau stained transfers of each blot are shown. SM=10% starting material; UB=10% unbound fraction; IP=immunoprecipitate. *=monoclonal antibody, **=recombinant antibody

Figure 3: TYRO protein tyrosine kinase-binding protein antibody screening by immunofluorescence.

THP-1 WT and *TYROBP* KO cells were labelled with a green or a far-red fluorescent dye, respectively. WT and KO cells were mixed and plated to a 1:1 ratio on coverslips. Cells were stained with the indicated TYRO protein tyrosine kinase-binding protein antibodies and with the corresponding Alexa-fluor 555 coupled secondary antibody including DAPI. Acquisition of the blue (nucleus-DAPI), green (identification of WT cells), red (antibody staining) and far-red (identification of KO cells) channels was performed. Representative images of the merged blue and red (grayscale) channels are shown. WT and KO cells are outlined with yellow and magenta dashed line, respectively. Antibody dilution used: MAB5240* at 1/500, ab93846 at 1/1000, ab155779 at 1/600, ab124834** at 1/700, MA5-34787** at 1/1000, PA5-26861 at 1/400, 97415** at 1/400, 46088** at 1/500, 12492** at 1/40. Bars = 10 µm. *=monoclonal antibody, **=recombinant antibody




Figure 1: TYRO protein tyrosine kinase-binding protein antibody screening by immunoblot




Figure 2: TYRO protein tyrosine kinase-binding protein antibody screening by immunoprecipitation




Figure 3 : TYRO protein tyrosine kinase-binding protein antibody screening by immunofluorescence

Materials and methods

Antibodies

All TYRO protein tyrosine kinase-binding protein antibodies are listed in Table 1. Peroxidase-conjugated goat anti-mouse and anti-rabbit antibodies are from Thermo Fisher Scientific (cat. number 62-6520 and 65-6120). Alexa-555-conjugated goat anti-mouse and anti-rabbit secondary antibodies are from Thermo Fisher Scientific (cat. number A21424 and A21429).

Cell culture

Cells were cultured in DMEM high glucose (GE Healthcare cat. number SH30081.01) containing 10% fetal bovine serum (Wisent, cat. number 080450), 2 mM L-glutamate (Wisent cat. number 609065, 100 IU penicillin and 100 µg/ml streptomycin (Wisent cat. number 450201).

Antibody screening by immunoblot

Immunoblots were performed as described in our standard operating procedure [5]. THP-1 WT and *TYROBP* KO were collected in RIPA buffer (25mM Tris-HCl pH 7.6, 150mM NaCl, 1% NP-40, 1% sodium deoxycholate, 0.1% SDS) supplemented with protease inhibitor. Lysates were sonicated briefly and incubated 30 min on ice. Lysates were spun at ~110,000xg for 15 min at 4°C and equal protein aliquots of the supernatants were analyzed by SDS-PAGE and immunoblot. BLUelf prestained protein ladder from GeneDireX (cat. number PM008-0500) was used.

Immunoblots were performed with large 10-20% gradient polyacrylamide gels and transferred on nitrocellulose membranes. Proteins on the blots were visualized with Ponceau staining which is scanned to show together with individual immunoblot. Blots were blocked with 5% milk for 1 hr, and antibodies were incubated O/N at 4°C with 5% bovine serum albumin in TBS with 0,1% Tween 20 (TBST). Following three washes with TBST, the peroxidase conjugated secondary antibody was incubated at a dilution of ~0.2 µg/ml in TBST with 5% milk for 1 hr at room temperature followed by three washes with TBST. Membranes are incubated with ECL from Pierce (cat. number 32106) prior to detection with HyBlot CL autoradiography films from Denville (cat. number 1159T41).

Antibody screening by immunoprecipitation

Immunoprecipitation was performed as described in our standard operating procedure [6]. Antibody-bead conjugates were prepared by adding 1 µg to 500 ul of Pierce IP Lysis Buffer from

Thermo Fisher Scientific (cat. number 87788) in a microcentrifuge tube, together with with 30µl of Dynabeads protein A- (for rabbit antibodies) or protein G- (for mouse antibodies) from Thermo Fisher Scientific (cat. number 10002D and 10004D, respectively). Pierce IP Lysis Buffer was supplemented with the Halt Protease Inhibitor Cocktail 100X from Thermo Fisher Scientific (cat. number 78446) at a final concentration of 1x. Tubes were rocked for ~2 hrs at 4°C followed by several washes to remove unbound antibodies.

THP-1 WT were collected in Pierce IP buffer (25 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% NP-40 and 5% glycerol) supplemented with protease inhibitor. Lysates are rocked 30 min at 4°C and spun at 110,000xg for 15 min at 4°C. One ml aliquots at 1.0 mg/ml of lysate were incubated with an antibody-bead conjugate for ~2 hrs at 4°C. Following centrifugation, the unbound fractions were collected, and beads were subsequently washed three times with 1.0 ml of IP lysis buffer and processed for SDS-PAGE and immunoblot on 10-20% polyacrylamide gels. Prot-A:HRP (MilliporeSigma, cat. number P8651) was used as a secondary detection system at a dilution of 0.4 µg/ml for an experiment where a rabbit antibody was used for both immunoprecipitation and its corresponding immunoblot. both immunoprecipitation and its corresponding immunoblot.

Antibody screening by immunofluorescence

Immunofluorescence was performed as described in our standard operating procedure [7]. THP-1 WT and *TYROBP* KO were labelled with a green and a far-red fluorescence dye, respectively. The fluorescent dyes used are from Thermo Fisher Scientific (cat. number C2925 and C34565). WT and KO cells were plated on glass coverslips as a mosaic and incubated for 24 hrs in a cell culture incubator. Cells were fixed in 4% PFA (in PBS) for 15 min at room temperature and then washed 3 times with PBS. Cells were permeabilized in PBS with 0,1% Triton X-100 for 10 min at room temperature and blocked with PBS with 5% BSA, 5% goat serum and 0.01% Triton X-100 for 30 min at room temperature. Cells were incubated with IF buffer (PBS, 5% BSA, 0,01% Triton X-100) containing the primary TYRO protein tyrosine kinase-binding protein antibodies O/N at 4°C. Cells were then washed 3 × 10 min with IF buffer and incubated with corresponding Alexa Fluor 555-conjugated secondary antibodies in IF buffer at a dilution of 1.0 µg/ml for 1 hr at room temperature with DAPI. Cells were washed 3 × 10 min with IF buffer and once with PBS. Coverslips were mounted on a microscopic slide using fluorescence mounting media (DAKO).

Imaging was performed using a Zeiss LSM 700 laser scanning confocal microscope equipped with a Plan-Apo 20x air objective (NA = 0.8). Analysis was done using the Zen navigation software (Zeiss). All cell images represent a single focal plane. Figures were assembled with Adobe Illustrator.

References

1. Laflamme, C., et al., *Opinion: Independent third-party entities as a model for validation of commercial antibodies*. N Biotechnol, 2021. **65**: p. 1-8 DOI: 10.1016/j.nbt.2021.07.001.
2. Laflamme, C., et al., *Implementation of an antibody characterization procedure and application to the major ALS/FTD disease gene C9ORF72*. Elife, 2019. **8** DOI: 10.7554/eLife.48363.
3. Nusinow, D.P., et al., *Quantitative Proteomics of the Cancer Cell Line Encyclopedia*. Cell, 2020. **180**(2): p. 387-402 e16 DOI: 10.1016/j.cell.2019.12.023.
4. *DepMap, Broad*. 2019.
5. Ayoubi, R., P.S. McPherson, and C. Laflamme, *Antibody Screening by Immunoblot*. 2021 DOI: <https://doi.org/10.5281/zenodo.5717510>.
6. Ayoubi, R., et al., *Antibody screening by Immunoprecipitation*. 2021 DOI: <https://doi.org/10.5281/zenodo.5717516>.
7. Alshafie, W., P. McPherson, and C. Laflamme, *Antibody screening by Immunofluorescence*. 2021 DOI: <https://doi.org/10.5281/zenodo.5717498>.