

Centres of endemism of Noctuidae (Lepidoptera) in the Palaearctic arid mountains: biogeographical and phylogenetic implications

With 21 figures, 4 tables and 2 Appendix

ZOLTÁN VARGA¹

¹ Dept. Evolutionary Zoology & Human Biology, Faculty of Science, University of Debrecen, H-4010 Debrecen, Hungary. – varga.zoltan@science.unideb.hu

Published on 2022-07-31

DOI: 10.3897/contrib.entomol.72.e87196

Abstract

The oreal fauna is connected with orographically limited non-arboreal habitats. Its chorological centres can be recognised by the high species-diversity of numerous typical genera, and by the accumulated occurrence of endemic species and/or subspecies of disjunct species. The oreal fauna is partitioned to the alpine type, as the faunal type of humid high-mountains with strong connections to the tundral zonobiome, and the xeromontane type, as the faunal type of arid high-mountains with close connections to the eremic zonobiomes. As the results of revisions of several Noctuinae genera, species groups and/or sister species were recognised and their distributions were mapped. The restricted areas of allopatric sister species, often described by us as new for science, fulfil the criteria of the “areas of endemism”. Core areas of the Palaearctic xeromontane Noctuidae, outlined by the distribution of endemic species, have been proven by the occurrence of allopatric subspecies of polytypic species, and/or by the presence of allopatric sister species. In the revised genera of Noctuidae several types of allopatric speciation have been identified based on the analysis of the areas of endemism and of vicariance patterns. As a result of these analyses, it is proved that allopatric sister species, as elementary monophyletic supraspecific units, are suitable for phylogenetic biogeographical surveys. Although the major part of the xeromontane fauna appears to be range-restricted, a considerable fraction of the species could have expanded into the steppic zonobiome due to adaptive changes of their life cycles. High diversity of cold-adapted species originated from the Sino-Himalayan mountains by passing two main filter-corridors. One track of this bifurcation was directed across the “Rhododendron-corridor” to the Holarctic taiga zone while the other one, across the “Xeromontane filter-corridor” to the mountain systems of Central and Inner Asia. This bifurcation becomes apparent from the taxonomic division of the genera, composing both of these main faunal types. Supposedly, the faunal movements of the xeromontane species in the West Palaearctic had been shaped by the Messinian salinity crisis and, additionally, significantly influenced by the Mid-Pleistocene climatic transition which deeply transformed the zonality of the vegetation by cooling and aridisation of vast areas.

Key words

Noctuidae, oreal fauna, xeromontane fauna, areas of endemism, monophyletic species groups, allopatric speciation, climatic constraints, filter-corridors

Zusammenfassung

Die Orealfauna ist mit den orographisch geprägten non-arborealen Lebensräumen eng verbunden. Ihre chorologischen Zentren lassen sich durch die Artenmannigkeit bestimmter Genera und ein gehäuftes Vorkommen von endemischen Arten bzw. Subspezies disjunkter Arten erkennen. Die Orealfauna unterteilt sich einerseits in den alpinen Faunentyp, d. h. den Faunentyp der humiden Hochgebirge, mit engen Beziehungen zum tundralen Zonobiom, andererseits in den xeromontanen Faunentyp, d. h. den Faunentyp der ariden Hochgebirge, mit engen Beziehungen zum eremialen Zonobiom. Als ein Ergebnis der Revisionen mehrerer Noctuiden-Genera, Artengruppen bzw. Geschwisterarten konnte dies nachgewiesen werden; ihre Verbreitungen wurden kartiert. Die engbegrenzten Areale der allopatrischen Geschwisterarten, die oft als neu für die Wissenschaft beschrieben wurden, entsprechen den Kriterien der „*areas of endemism*“. Die Kerngebiete der palaearktischen xeromontanen Noctuidae, die aufgrund der Verbreitung der endemischen Arten postuliert wurden, wurden durch das Vorkommen der allopatrischen Subspezies der polytypischen Arten, bzw. der allopatrischen Geschwisterarten bestätigt. In den revidierten Noctuidengattungen wurden mehrere Typen der allopatrischen Artbildung durch die Analyse von Endemismusarealen bzw. Vikarianzmustern nachgewiesen. Als Ergebnis dieser Untersuchungen wurden die allopatrischen Geschwisterarten als elementare monophyletische supraspezifische Einheiten bestätigt, die sich als geeignet für phylogenetisch-biogeographische Untersuchungen erwiesen. Obwohl der größte Teil der xeromontanen Fauna engbegrenzte Areale aufzuweisen scheint, konnte sich ein beträchtlicher Anteil der Arten weit in die zonalen Steppengebiete durch die adaptiven Veränderungen der Lebenszyklen ausbreiten. Die große Mannigfaltigkeit der im sino-himalayischen Hochgebirge entstandenen kälteangepassten Arten lässt sich durch das Durchdringen von zwei wichtigen Filter-Korridoren ableiten. Ein Zweig dieser Bifurcation verläuft durch den „Rhododendron-Korridor“ in den holarktischen Taiga-Gürtel, während ein anderer Zweig durch den xeromontanen Filter-Korridor in die Gebirgssysteme von Zentral- und Innerasien verläuft. Diese Unterteilung lässt sich durch die unterschiedliche taxonomische Verteilung jener Genera verstehen, aus denen diese beiden grundsätzlichen Faunentypen bestehen. Es ist wahrscheinlich, dass die Ausbreitungsvorgänge der xeromontanen Elemente in der Westpaläarktis durch die Messinischen Salinitätskrise geformt wurden; einen weiteren wichtigen Einfluss besaß die mittelpaläozäne klimatische Transition, während derer durch großräumige Abkühlung und Austrocknung die zonale Anordnung der Vegetation umgeordnet wurde.

Introduction - Oreal fauna, definition and subdivision

The composition of terrestrial animal assemblages strongly depends on the level of primary production, limited by the available solar energy and water circulation (BEER et al. 2010, PAPPAS et al. 2016, SIMOVÁ & STORCH 2017). Satellite images clearly reveal that both the Net Primary Production (NPP) and the Gross Primary Production (GPP) are the lowest in the tundral and desert zonobiomes. They are, however, similarly low in orobiomes, i.e. in different high mountains, surrounded by belts with a higher level of primary production (e.g. RODIN & BAZILEVICH 1966, PRINCE & GOWARD 1995, RUNNING et al. 2004). If we simply specify the arboreal biomes (see: DE LATTIN 1967) as terrestrial habitats characterised by medium or higher level of primary production, the oreal fauna can be identified as the faunal type of the orographically defined non-arboreal biomes (VARGA 1997). Its members, as a rule, have insular, endemic or disjunct areas in the Eurasian high mountains (Fig. 1). Their core areas can be recognised by the high species-diversity of numerous typical genera and, thereby, the accumulated occurrence of endemic species and/or subspecies of disjunct species.

Parallel with the non-arboreal zonobiomes, the oreal fauna can be subdivided into the *alpine* type, as the faunal type of humid high-mountains with prevailing glacial

weathering and morphology, and with strong connections to the *tundral* zonobiome, and the *xeromontane* type, as the faunal type of arid high-mountains with prevailing physical weathering and manifold connections to the *eremic* zonobiomes. As the dynamics of the *alpine* faunal type is closely connected with the Quaternary glaciations, its history is characterised by long-distance range translocations and disjunctions, resulting in the diversification of a great number of *arctic-alpine* species (VARGA & SCHMITT 2008, SCHMITT 2009). On the contrary, the evolutionary history of the *xeromontane* faunal type was supposedly less disturbed by glaciation-interglaciation cycles. Thus, it has a high potential for speciation in several groups adapted to the seasonally arid conditions as many Noctuinae moths belonging to the genera *Euxoa*, *Dichagyris*, *Actebia* s.l., *Chersotis*, *Rhyacia*, *Standfussiana*, *Eugnorisma*, *Spaelotis*, *Xenophysa*, etc. (VARGA 1995, 1997, 2010b), Polyommatus subg. *Agrodiaetus*, ECKWEILER & HÄUSER 1997, KANDUL et al. 2004, 2007, LUKHTANOV et al. 2005), and also in some flightless grasshopper groups (e.g. *Conophyma*, *Nocaracris*, *Nocarodes*, see: LI et al. 2011, ÜNAL 2016). Considering the composition of the European alpine Lepidoptera, high species diversity was found in several “Microlepidoptera” families and genera (Gelechiidae:

Caryocolum, HUEMER et al. 2014, HUEMER & KARSHOLT 2020; *Sattleria*, HUEMER & HEBERT 2011, HUEMER & TIMOSSI 2014; Yponomeutidae: *Kessleria*, HUEMER & MUTANEN 2015), certain genera of Geometridae (e.g. *Entephria*, *Psodos*, *Charissa*, *Elophos*; HUEMER 1998, MÜLLER et al. 2019) and Nymphalidae (*Boloria*, *Erebia*; VARGA 1996, 2003, VARGA & SCHMITT 2008), however, surprisingly much less in the otherwise highly diverse Noctuidae (VARGA 2003). This situation is contrasting with the outstanding taxonomic diversity of the xeromontane faunal assemblages in the Central and Inner Asiatic¹ mountain systems (Fig. 1, VARGA 2010b) where numerous genera of Noctuidae (mostly of Noctuinae, Hadenini, Xylenini, Apameini, and Oncocnemidinae) feeding on grasses and herbaceous plants predominate not only in the species composition but also in the density of individuals.

Following these preliminary considerations, I developed the hypothesis that we can recognise some typically repeated patterns of vicariance in range-restricted sister species and, in parallel, essentially similar disjunct distributions in polytypic species in numerous genera of the xeromontane Noctuidae. Based on the taxonomic revisions of several diverse genera, mostly from Noctuinae, I also hypothesised that these patterns reveal important general trends of allopatric speciation which can be uncovered by phylogenetic biogeographical methods. Furthermore, I found it plausible to search for certain repeatedly appearing biogeographical and phylogenetic connections between the range-restricted xeromontane species and the zonally distributed congeneric steppic species. Finally, I hypothesised that the core areas of the Palaearctic xeromontane fauna are connected with the Central and Inner Asiatic high mountains as a consequence of the cooling and aridisation of these regions following the emergence of the Sino-Himalayan mountain systems and these core areas were modulated by the Mid-Pleistocene climatic transition.

Materials and methods

This biogeographical review is based on generic and suprageneric taxonomic revisions of several Noctuidae genera (mostly Noctuinae: *Dichagyris*, *Actebia*, *Euxoa*, *Chersotis*, *Rhyacia*, *Standfussiana*, *Eugnorisma*, *Goniographa*, *Spaelotis*, *Xenophysa*). Most of the type specimens of the revised taxa were studied; thousands of genitalia slides were prepared, photographically and graphically documented and electronically stored. In a book series (VARGA et al. 2013, 2015, 2019) and in generic revisions (VARGA & RONKAY 1987, 1996, 2002; RONKAY & VARGA 1999; VARGA & GYULAI 1999; VARGA 1998, 2011a, b, 2014; VARGA et al. 1989, 2018) numerous

new genera, subgenera, species and subspecies were described, subgenera and species groups were separated, pairs or triplets of sister species were stated, and new taxonomical statuses were established.

Based on the data of large European public and private collections, and of relevant literature, distribution maps of Noctuinae species were prepared, with special respect to allopatric sister species and disjunct polytypic species. Distributional data matrices of the genera *Dichagyris*, *Chersotis*, *Rhyacia*, *Standfussiana*, *Eugnorisma*, *Goniographa*, *Xenophysa* were prepared and the species were ordered to some types of distributions (Supplement 1), and they also grouped according to the types of geographical variations (Results 1). In the revised genera the species have been arranged into subgeneric groups, the allopatric sister species/subspecies relations were established, and based on them, area-dendograms of species groups were constructed (Figs 4, 5, 13).

In the first step, following the principle of the “areas of endemism” (HAROLD & MOOI 1994), I outlined some core areas by the accumulated occurrence of strictly endemic species (e.g. Kopet-Dagh Mts, parts of Hindukush Mts, the Pamirs and the Tien Shan system, etc. (Results 2). In the next step, these core areas were confirmed by the repeated patterns of disjunct polytypic species, and pairs (eventually triplets) of allopatric sister species. These species-group taxa were considered as elementary monophyletic supraspecific units (Results 3). Principally, this procedure is followed by the analysis of monophyletic species groups within a genus, then by phylogenetic analyses of subgenera and genera etc. It is essential that the whole multi-step procedure runs from the taxonomically lower levels to the higher ones. As the last step, according to the available life cycle data (aestivation, hibernation), some eco-physiological backgrounds of restricted distribution vs expansive area dynamics were uncovered (Figs 7, 14, 15, 18; Results 3 and Discussion 1).

Results 1: Types of geographical and taxonomical subdivisions in xeromontane Noctuidae

Based on the geographical variation (in widely distributed, mostly polytypic species) and on the patterns of vicariance (in disjunct species and allopatric sister species), several types of differentiation were recognised in Palaearctic xeromontane Noctuidae.

The following types of geographical variation have been separated (Tables 1–4, Supplement I).

- Widely distributed, often polytypic species with parallel polymorphism of distinctly marked vs. unicolorous forms in different populations² (e.g. in some

¹ On the meaning of the terms Central vs Inner Asia see: Fig. 1 and Discussion 2.

² The full names of the species (with authors and years) mentioned as examples in the text are listed in the Tables 1–4 and Appendix I.

- Euxoa* spp.: the “huebneroides” types vs unicolorous forms, *Dichagyris terminicincta*, *D. psammochroa*, *Eugnorisma* (E.) *insignata*, *E. (Metagnorisma) pontica*). In these species the relative frequency of parallel marking and/or colouration variations often show some geographical trends (e.g. *Euxoa christophi* & f. *lugens*, *Euxoa sigmata* & f. *designata*, *D. terminicincta* f. *phaeotaenia* & f. *capnista*; *D. psammochroa* & f. *dichroa*, *E. insignata* f. *pallescens* & f. *leuconeura*; *E. (M.) pontica* & f. *consenescens*), however, without any clear subspecific differentiation.
- Parallel polymorphism and taxonomic differentiation are combined in the polytypic *Euxoa adumbrata*. As an extreme case, a contrasting vs. unicolorous dark bilateral somatic mosaic male specimen (Mongolia, Govi Altai aimak, Adž Bogd Mts, leg. Peregovits & Varga, figured in Fibiger 1997: 35) was found as evidence of co-specificity of infraspecific colour morphs. Incipient allopatric speciation was shown in the widely distributed, disjunct West-Central Asiatic sister species *Euxoa homicida* – *E. transcaspica* (Fig. 2). These cases may be interpreted as intermediate stages of differentiation of a polytypic species into a complex of sister species.
 - Some further species are also polytypic (*polytypic species) with disjunctions and peripheral subspeciation at the western boundary of the range, e.g.: *Euxoa heringi**, *Dichagyris melanura**³, *D. celsicola**, *Chersotis capnistis**, *Ch. zukowskyi**, *Ch. laeta**, *Rhyacia psammia** (= *nyctymerides*). These species are distributed in Asia Minor and/or in Western and Central Asiatic mountains. The marginal populations of most species are fragmented in high mountains of the southern part of the Balkan Peninsula by the breakdown of the Aegean Arc (Fig. 3). This process has already resulted in multiple cases of insular speciation in Crete, e.g. *Euxoa malickyi* (sister species of *E. heringi*), *Dichagyris rhadamanthys* (sister species of *D. melanura*).
 - Sibling species with peripatric isolation in West and/or Central Asia: These are widely distributed species with an allo- or parapatric sister species at the eastern or south-eastern periphery of the range of the more expansive, often polytypic* species, e.g. *Dichagyris vallesiaca** – *D. fuscashmiriana*, *D. verecunda** – *D. karakorealis*, *D. singularis** – *D. melanofusca*, *Eugnorisma insignata** – *E. conformis* (see: VARGA, RONKAY & YELA 1990: Figs 73, 75); *Chersotis elegans* – *Ch. eberti* (see: DUFAY & VARGA 1995, VARGA 1997: Fig. 7), *Ch. juvenis* – *Ch. kouros* (Fig. 4), *Ch. sordescens* – *Ch. hercizi* (Fig. 5). These cases are considered as results of allopatric speciation by peripheral isolation.
 - Allopatric pairs of West- and Central/Inner Asiatic sibling species (Tables 1–3): *Euxoa aneucta* – *E. subeucta*, *E. sigmata* – *E. metasigmata* (Fig. 6), *Dichagyris striata* – *D. tyrannus*, *D. humilis* – *D. hypotacta*, *D. darius* – *D. argentea*, *D. taftana* – *D. guentereberti*, *D. strenua* – *D. vietteana*; *Hemixarnis iuguma* – *H. berezskii*; *Chersotis firdusi* – *Ch. fidahusseini* (Fig. 5); *Ch. capnistis* – *Ch. leucostola* (Fig. 7), *Rhyacia gabori* – *R. evartianae* (Fig. 8); *Goniographa marcida* – *G. gyulaipeteri*; *Xenophysa cacumena* – *X. afghanorea* (MIKKOLA & et al. 1987; VARGA 1993, 1998, 2002, 2014; VARGA & GYULAI 2000; VARGA et al. 2011). In these species the populations of Asia Minor, West and North Iran are taxonomically differentiated from the populations distributed in Central Asia (Pamirs, Hindu Kush and Tien-Shan Mts). In other cases the allopatric species occur in Iran + Transcaspia, and in Central Asia (Hissar-Darwaz Mts, Pamirs, Hindu Kush Mts, Tien-Shan Mts). The allopatric species are usually separated by the huge arid belt East of Transcaspia (see: Figs 5, 7, 8) as a consequence of range-restriction by aridisation of Central Asia.
 - Allopatric pairs (or triplets) of Central Asiatic sibling species, differentiated either between different parts of the Tien Shan system, or between the Tien Shan system and other Central Asiatic massifs as Hissar Mts, Pamirs and/or Hindu Kush (Table 3): *Chersotis vicina* – *Ch. petermarci*; *Ch. calorica* – *Ch. shandur* (Fig. 4), *Ch. delear* – *Ch. electrographa* – *Ch. vargai*, *Rhyacia diplogramma* – *R. oromys*; *Goniographa decussa* – *G. discussa*; *G. funkei* – *G. metafunkei* – *G. naumannii* (Fig. 9).
 - Allo- or parapatric groups of species, similarly to the former groups, with long distance disjunctions between the Anatolian-Iranian and Central Asiatic mountains (Table 1, 3): *Dichagyris catalaipa* – *D. psammochroa* – *D. afghana* – *D. kurbatskyi* – *D. apochora*; *Chersotis binaloudi* – *Ch. antigrapha* – *Ch. argylographa* (Fig. 10); *Chersotis ronkayorum* – *Ch. sterilis* – *Ch. nitens* – *Ch. metagrapha* (Fig. 7); *Chersotis sarhada* – *Ch. hoppei* – *Ch. lehmanni* – *Ch. lukhtanovi*.
 - Pairs or groups of partially sympatric sibling species, evolved by former geographical isolation and secondary overlap: *Chersotis andereggi* – *Ch. acutangula*; *Ch. juvenis* – *Ch. kouros*; *Ch. ocellina* – *Ch. alpestris* – *Ch. oreina* – *Ch. stridula* – *Ch. transiens*; *Chersotis hahni* – *Ch. curvispina* – *Ch. cryptographa*; *Rhyacia subdecora* – *R. scythropa* – *R. oxytheca* (Fig. 8); *Bryopolia virescens* – *B. tsvetaevi* – *B. chamaeleon* – *B. chrysospora*; *Bryopolia monotona* – *B. thomasi* – *B. ronkayorum* (VARGA et al. 1990).
 - Species groups evolved by vicariance and, additionally, by the change of the life cycle. In these cases some allopatric endemic species are opposed to a widely distributed sister clade in which the increased dispersal capacity is the consequence of the adaptive change of the life-cycle (*Chersotis capnistis*-group, VARGA & RONKAY 1996). Detailed description of these life-cycle changes and biogeographical consequences is provided in the Results 3.

³ The western isolated populations represent the nominotypical subspecies.




Fig. 1: Important parts and mountain systems of Eurasia.




Fig. 2: Sister species with incipient allopatric speciation, with slight differentiation in the genitalia: *Euxoa homicida* (Asia minor and Iran, exception: Khorasan and Baluchistan) and *E. transcaspica* (Transcaspia, Afghanistan, Pakistan; subspecies are here not distinguished).




Fig. 3: Distribution of the species: *Dichagyris celsicola gracilis* (1), *Chersotis zukowskyi* (2), and *Rhyacia psammia* (3). They are distributed in Western and/or Central Asiatic mountains; some marginal populations have been fragmented in the mountains of the southern part of the Balkan Peninsula by the breakdown of the Aegean bridge. *Rhyacia psammia* is subdivided to six subspecies (VARGA 2011).

Results 2: The xeromontane fauna: areas of endemism and phylogenetic implications

A major part of the xeromontane fauna appears to be range-restricted with numerous relict-like species, especially in certain mountain systems of Central and Inner Asia. The restricted areas of the endemic species and of the allopatric sister-species correspond to the criteria of the “*areas of endemism*” (HAROLD & MOOI 1994), therefore, as monophyletic species groups, they are suitable for multi-step phylogenetic-biogeographical studies (see: Methods).

As a result of taxonomic revisions, those genera proved to be most suitable for phylogenetic-biogeographical analyses in which (i) there are numerous strictly endemic species, (ii) there are also some polytypic species with disjunct ranges, and (iii) also some expansive species occur with large, seemingly continuous ranges. These highly diverse genera, as e.g. *Euxoa*, *Dichagyris* s.l., *Actebia* s.l. (subg.: *Parexarnis*, *Protexarnis*, *Hemixexarnis*), *Chersotis* (VARGA 1998), *Rhyacia*, *Eugnorisma* (VARGA, RONKAY & YELA 1990), *Xenophysa* (VARGA 1989a, 2011) etc. can often be subdivided into some subgenera and/or several groups of species with closer phylogenetic relationships (Supplement 2).

Eco-physiological responses to climatic challenges proved to be significant drivers of the ranges. The phylogenetic bifurcation of the Noctuinae genus *Xenophysa* (Figs 11–13) reflects a close connection with some essentially different climatic belts of the arid high mountains of Eurasia. The range of distribution of the western lineage (“*X. junctimacula*”-species group) reaches the central part of the Hindu Kush Mts, and its limitation clearly coincides with the eastern boundary of the sub-Mediterranean equinoctial type of precipitation (AGAKHANJANTS 1981, AGAKHANJANTS & BRECKLE 1995). The only widespread species pair of this group, *X. cacumena* and *X. afghanoreia* shows a typical long-distance disjunction (Elburz and Kopet-Dagh Mts vs. Hindu Kush Mts) with occurrences in the western Hindu Kush massif (Koh-i-Baba) and the Paghman Mts. The presence of numerous range-restricted species of this group (*X. argyrogramma*, *X. xenogramma*, *X. monastica*, *X. poecilogramma*) seems to be confined to the central and eastern-south-eastern flank of the Hindu Kush mountain range (Fig. 11). The range of the eastern lineage (“*X. agnoscita*”-group: *X. agnoscita*, *X. naumannii*, *X. pseudopoecila* and the more isolated *X. sharhu*) extends from the Saravshan and Hissaro-Darwaz Mts to the western part of Mongolia (Fig. 12), reaching also to the western Tien-Shan and Karakoram Mts. They have a narrow belt of overlapping with the western species group, mostly in the Hissaro-Darwaz range and in the eastern part of the Hindu Kush Mts. The most closely related sister species, the widespread *X. agnoscita* and the restricted *X. naumannii* seem to be parapatric, while the distribution of the widespread *X. pseudopoecila* is partly sympatric with both species. However, *X. poecilogramma*, which belongs to the western group, locally co-occurs

with both latter species in the eastern flank of the Hindu Kush range (Shandur pass, Figs 11–13).

The range of distribution of some genera or subgenera of Noctuidae, typical for xerothalmic scrub-forests of western Asia (e.g. *Ostheldera*, see: RONKAY & VARGA 1993; species groups of the large and heterogeneous genera *Polymixis*, *Mniotype*, *Anchoscelis* etc.) also extends to the same eastern boundary of the sub-Mediterranean type of precipitation in the Hindu Kush and western Tien-Shan ranges as the western clade of *Xenophysa*.

The variety of allo- and sympatric distributions of sister species, combined with disjunctions of the ranges, has been observed in the *Eugnorisma chaldaica* – *E. spodia* species group. The widespread *E. chaldaica* occurs from Central Anatolia and the south-eastern Russian steppes to southern Siberia, with subspeciation from south-eastern Turkey to the adjacent areas of North Iran and also in the western Altai Mts (Fig. 14). Between the two major population groups of *E. chaldaica* a wide disjunction is gapping, from North Iran to the western Altai Mts in north-eastern part of Kazakhstan, which is „filled“ by further taxa: *E. kristenseni* and the stenotopic *E. spodia spodia* in the Kopet-Dagh Mts, and the widely distributed *E. kristenseni* and *E. spodia psammochrea* in southern Kazakhstan and Uzbekistan, respectively (VARGA & RONKAY 1987; Varga & et al. 1990, 2015). In the *Eugnorisma ignoratum* species group (Fig. 15) there are also two widespread species: *Eugnorisma ignoratum* and *E. mikkolai* occurring in the hilly and zonal steppe areas, as opposed to the montane *E. puengeleri* (sister species of *E. mikkolai*, eastern Hindu Kush in Afghanistan and North-West Pakistan), *E. tamerlana*, a Central Asiatic semi-desert species and the endemic *E. cuneiferum*, which only occurs in the mountains of Transcaspia (Iran, Prov. Khorasan and Turkmenistan: Kopet-Dagh Mts, VARGA et al. 1990, VARGA & RONKAY 1994).

Results 3: Allopatric speciation and species groups in the genera *Chersotis* BOISDUVAL, 1840 and *Rhyacia* HÜBNER, [1821]

The genus *Chersotis* is rich in pairs and/or groups of closely related species (Tables 3–4, Appendix II). Here those allopatric species-groups are treated which are distinguished by recognised synapomorphies of genitalia. Significant characters of these species-groups are already described and illustrated in taxonomic revisions (VARGA 1996b, 1998; VARGA et al. 2013).

We have selected two species groups which nearly exclusively consist of range-restricted species. The sibling pair *Chersotis firdusii* (Elburz Mts; Kopet-Dagh Mts) and *Ch. fidahusseini* (West Tien Shan, Hindu Kush, Karakoram, North-West Himalaya) are considered as sister group of the species pair consisting of the more widespread *Ch. sordescens* and the West Himalayan endemic




Fig. 4: Distribution of the pairs of sister species: *Ch. juvenis* (1. Asia Minor, Iran, exception: Khorasan) – *Ch. kouros* (2. Iran: Khorasan, Turkmenistan) and *Ch. calorica* (3. Kazakhstan, China: Xinjiang) – *Ch. shandur* (4. Afghanistan, Pakistan). Taxonomic details: VARGA et al. 2013.

Ch. herczi (Himalayas: Pakistan, Kaghan valley (Fig. 5)). The *Ch. rectangula-andereggii* species group can be regarded as outgroup of this species-group. The next „triplet“ of species uniformly shows some characteristic features of genital structures. The westernmost species, *Ch. binaloudi* occurs in Khorassan (NE Iran), and its sister species are the ‐twins‐: *Ch. antigrapha* (Hindukush Mts) and *Ch. argylographa* (W Pamirs range, Fig. 10). We consider the set of the *Ch. vicina* group and *Ch. juvenis* group as their sister-group (the taxonomic details and species descriptions are published by VARGA 1996b, 1998; VARGA & RONKAY 1996; VARGA et al. 2013). Both latter species groups consist of range-restricted allopatric species.

Nearly the same pattern of allopatric distribution was observed in a species group (subg. *Anchorhyacia* VARGA, 2011) of the closely related genus *Rhyacia*. The allopatric *R. gabori* (Kopet-Dagh Mts) and *R. evarthanae* (Hindukush and Hissaro-Darwaz range) are forming together the sister-group of *R. scythropa* (Hindukush: Koh-i-Baba and Paghman Mts) and *R. oxytheca* (West Pamirs, East Hindukush, Karakoram, North-West Himalayas, Fig. 8). All these species share the acute process of the *juxta* as synapomorphy. Their probable sister-group is the widely distributed, polytypic *R. subdecora*. Two further species pairs also belong to

this subgenus, the partly sympatric *R. psammia** (polytypic, widely distributed) – *R. nyctymerina* (Tien Shan) and the allopatric twin species *R. diplogramma* (Tien Shan system) and *R. oromys* (Pamirs, Hindukush, West Himalayas). All of these species groups share the synapomorphies as follows: the modified clavus densely covered by small spines, the hook-shaped carina of the aedeagus, and the large subbasal and the conical, sclerotised medial diverticula of the vesica (VARGA 1990, 2011).

In some other cases, there is a partial overlap in the distribution of the closely related species. These overlapping species have a fairly large extended range with subspecific subdivision (e.g. the species pairs *Chersotis rectangula* and *Ch. andereggii*, *Ch. elegans** and *Ch. anatolica**, *Ch. fimbriola** and *Ch. laeta**, Fig. 16). As all of these species are widely distributed and mostly polytypic*, their distribution patterns should be considered as cases of secondary overlaps, subsequently to the previous allopatric process of speciation.

The geographical extension of the allopatric areas of sister species may be rather different. The allopatric *Chersotis andereggii* and *Ch. juncta*, *Ch. firdusii* and *Ch. fidahusseini*, *Ch. capnistis** and *Ch. leucostola*, *Ch. sarhana* and *Ch. lehmanni* have relatively extended ranges, while in other cases the large area of an expansive species is combined with the peripherally isolated ranges




Fig. 5: Distribution of the pairs of sister species: *Chersotis firdusii* (1. Iran) – *Ch. fidahusseini* (2. Kirghisia, Tadzhikistan, Afghanistan, Pakistan, two subspecies) and *Ch. hercizi* (3. Pakistan: Kashmir) – *Ch. sordescens* (4. Central Asia). Taxonomic details: VARGA et al. 2013.




Fig. 6: Distribution of the pairs of sister species: *Euxoa sigmata* (1. Iran: Khorasan, Turkmenistan) – *E. metasigmata* (2. Afghanistan, Pakistan) and *E. aneucta* (3. Iran: Khorasan) – *E. subeucta* (4. Afghanistan, Pakistan). Taxonomic details: VARGA 2014.

of some range-restricted species, obviously originating from marginal splitting: *Ch. semna* vs. *Ch. pachnosa*; *Ch. juvenis* vs. *Ch. kouros*, *Ch. elegans** vs. *Ch. kacem* and *Ch. eberti*. In some other cases both members of the allopatric pair are strictly endemic, e.g.: *Chersotis*

ronkayorum – *Ch. sterilis*, *Ch. nitens* – *Ch. metrapha*, *Ch. petermarci* – *Ch. vicina*, *Ch. calorica* – *Ch. shandur*, *Ch. binaloudi* – *Ch. antigrapha*, *Ch. gratissima* – *Ch. zaghros*, *Ch. nuppenenorium* – *Ch. nekrasovi* (Fig. 17).




Fig. 7: The *Chersotis capnista* group is subdivided into four allopatric, endemic species (1. *Ch. ronkayorum*, 2. *Ch. sterilis*, 3. *Ch. nitens*, 4. *Ch. metrapha*) and to the *Ch. capnista* (5.) – *Ch. leucostola* (6.) pair of species. The latter have large fat-bodies and a life-cycle with aestivation of imagos with a relatively large range of distribution. Taxonomy and phylogeny: VARGA 1998.




Fig. 8: Distribution of the species: *Rhyacia gabori* (1) – *Rh. evartianae* (2) – *Rh. scythropa* (3) – *Rh. oxytheca* (4) – *Rh. subdecora* (5). The range of *Rh. psammia* (6) is on Fig. 1. Taxonomy and phylogeny: VARGA 2011.




Fig. 9: Distribution of the *Goniographa funkei*-group; one more distant species, *G. marcida* (1), and three closely related ones: *G. naumannii* (1), *G. metafunkei* (2), *G. funkei* (3). The sister species (2) and (3) are strictly allopatric, (1) and (2) are overlapping in the Hissar Mts. Taxonomy and phylogeny: VARGA & RONKAY 2002.




Fig. 10: Distribution of the *Dichagyris psammochroa*-group: *Dichagyris catalaipa* (1) – *D. psammochroa* (2a-2b) – *D. afghana* (3) – *D. apchora* (4) – *D. kurbatskyi* (5); Distribution of the *Chersotis binaloudi*-group: *Ch. binaloudi* (6) – *Ch. antigrapha* (7) – *Ch. argyllographa* (8). Taxonomy and phylogeny: VARGA 1993, 1998; VARGA & RONKAY 1996; VARGA et al. 2021.

However, the existence of a few, phylogenetically strongly separated species (e.g. *Ch. sjuntensis*, *Ch. glebosa*, *Ch. illauta*) supports the hypothesis, that in this genus also an older wave of speciation should have proceeded, from which most of the species became extinct. This hypothesis is strongly supported by such cases in which the species-groups consist of one strongly differentiated

species, which can be regarded as outlier of all other members of the species group, e.g. *Ch. larixia** (with extremely fragmented range of distribution) for the *Ch. elegans-kacem-eberti-anatolica* group. All these cases refer to an ancient and a subsequent, more recent wave of the allopatric speciation in the genus *Chersotis*.




Fig. 11: The West Asiatic species group of the genus *Xenophysa*: *X. junctimacula huberi* (1) – *X. junctimacula junctimacula* (2) – *X. cacumena* (3) – *X. afghanoreia* (4) – *X. argyrogramma* (5) – *X. xenogramma* (6) – *X. monastica* (7). Outgroup: *X. paradoxa* (8). Taxonomy and phylogeny: VARGA 2011.




Fig. 12: The Inner Asiatic species of the genus *Xenophysa*: *X. agnosta* (1) – *X. naumannii* (2) – *X. pseudopoecila* (3) – *X. poecilogramma* (4) – *X. sharhu* (5). Taxonomy and phylogeny: VARGA 2011.

Discussion 1: Climatic constraints in the life cycle evolution of Noctuidae

In *Noctuidae* there are several types of univoltine life cycles, adapted to the seasonally cold and/or arid, temperate climate by different forms of larval and/or imago diapause (RYABOV 1956; SUKHAREVA 1999; SAULICH et al. 2017). Such univoltine life cycles probably can be derived from ancestral polycyclic types due to the insertion of diapause periods as hibernation and/or aestivation (Fig. 19). In these cases the larvae can use some resources of short duration by the right timing of active periods, and they can also be adapted to the seasonal insufficiency of resources by the diapause. There are several, in parallel developed different combinations of life cycles, connected to the centres of species diversity,

mentioned above. One of the typical responses of noctuid moths to the aridity is the aestivation of the larvae (most often in the prepupal stage, SPECHT et al. 2013), and/or of the adults, which has evolved in parallel in many different taxonomical groups (Fig. 19), mostly in the subfamily *Noctuinae* (*Euxoa*, *Dichagyris* s.l., *Actebia* s.l., *Chersotis*, *Rhyacia* s.l., *Standfussiana*, *Noctua*, *Spaelotis* etc.). The well-developed abdominal fat bodies obligatorily serve as pre-requisite of the successful aestivation and reproduction after the aestivation of the adults.

In the “*capnistis*” group of the genus *Chersotis* there are four stenotopic endemic species (Fig. 7) in high altitudes of arid high mountains of West and Central Asia, without well-developed fat bodies, while the sister species *Ch. capnistis** and *Ch. leucostola* have large fat-bodies and a life cycle with imaginal aestivation, combined with




Fig. 13: The phylogenetic bifurcation of the Noctuinae genus *Xenophysa* clearly shows a close connection to the two main different climatic provinces of the arid high mountains of Eurasia. 1. *X. cacumena*, 1a. *X. afghanoreia*, 2h. *X. junctimacula huberi*, 2. *X. junctimacula*, 3. *X. argyrogramma*, 4. *X. xenogramma*, 5. *X. poecilogramma*, 6. *X. pseudopoecila*, 7. *X. naumannii*, 8. *X. agnistica*, 9. *X. sharhu*. p *X. paradoxa* (outgroup). Taxonomy and phylogeny: VARGA 2011.

a relatively large range of distribution (VARGA & RONKAY 1997: 115), extending into moderate altitudes in steppe areas of the southern Ural Mts (*Ch. capnistis*) or the Dzhungarian Ala-Too (*Ch. leucostola*).

The possibility to gain a “zonal” expansion into the steppe biome seems to be linked to those noctuid moths (mostly Noctuinae) which either emerge at early summer and subsequently have a post-aestivation dispersal period in the late summer or early autumn (Fig. 19) as the above-mentioned *Chersotis* species, or numerous species of *Noctua*, *Spaelotis* and *Standfussiana*, or they have their whole, much shorter adult period in this season of the year. See examples: *Euxoa basigramma**, *E. christophi* (Fig. 18), *E. deserta**, *E. diaphora*, *E. distinguenda**, *E. fallax*, *E. foeda*, *E. hastifera**, *E. sabuletorum*, *E. segnialis**; *Agrotis bifurca*, *A. psammocharis*, *A. ruta*, *A. trifurca*; *Chersotis elegans**, *Ch. anatolica**, *Eugnorisma chaldaica**, *E. eminens**, *E. ignoratum*, *E. insignata**, *E. mikkolai* etc. (Figs 14–15).

As one of main results of our preliminary taxonomic surveys (e.g. RONKAY & VARGA 1999; VARGA 1996b, 1998, 2011; VARGA & RONKAY 1997; VARGA et al. 2013), taxonomic groups of the Mediterranean xeromontane fauna appear to have two different main sources of origin. The bulk of genera probably can be derived from the primary bifurcation of the Palaearctic xeromontane

faunal complex (Fig. 20). The other group had evolved from diverse Mediterranean xerophilous arboreal groups by adaptation to the aridity in connection with the late Tertiary Messinian Saline Crisis (VARGA 1997, 2010b). This hypothesis is strongly supported by the “macro”-taxonomic duality of the Mediterranean xeromontane Noctuidae. Those genera which belong to the subfamily Noctuinae and have “cutworm”-type, terricolous larvae, obviously originated in the continental, western and central Asiatic orobomes (e.g. *Euxoa*, *Dichagyris*, *Chersotis*, *Rhyacia*, *Standfussiana*, etc.). Their Mediterranean representatives regularly belong to some different derived phyletic lines (subgenera) of these polytypic genera. Their East Mediterranean-Anatolian taxa often display western and Central Asiatic connections but only in a few cases a marginal speciation in the Mediterranean ranges.

This group of genera correspond to the “*saxobiotic ecofaunal complex*” of the Palaearctic Orthoptera, described by BEY-BIENKO (1948; see also SHUMAKOV 1963, PRAVDIN & MISHTSHENKO 1980). The diversity centres of *saxobiotic* Orthoptera genera (“kamenistiy tip fauni”, e.g. Catantopinae: *Conophyma* ZUBOVSKY, 1898; Pamphaginae: *Saxetania* MISHTSHENKO, 1951; *Nocarodes* FISCHER-WALDHEIM, 1846; *Nocaracris* UVAROV, 1928; *Paranocarodes* BOLIVAR, 1916; *Paranothrotes*




Fig. 14: Distribution of the species of the *Eugnorisma chaldaica* group: *E. chaldaica chaldaica* (1), *E. chaldaica kordestana* (2), *E. chaldaica rubicunda* (3), *E. spodia spodia* (4), *E. spodia psammochroa* (5), *E. kristensenii* (6). Taxonomy and phylogeny: VARGA et al. 2015.




Fig. 15: Distribution of the species: *Eugnorisma ignoratum* (1), *E. mikkolai* (2), *E. puengeleri* (3), *E. tamerlana* (4), and *E. cuneiferum* (5). Taxonomy and phylogeny: VARGA et al. 2015.

MISHTSHENKO, 1951, etc.) are to be found in the Western- and Central Asiatic arid-semiarid high mountains (Elburz, Kopet-Dagh, Hindukush, Hissaro-Darwaz and western

Tien-Shan ranges). Their ranges greatly overlap with the core areas of the Noctuidae genera mentioned above.




Fig. 16: Distribution of the species pair *Chersotis fimbriola* – *Ch. laeta*: *Ch. fimbriola iminenia* (1), *Ch. fimbriola iberica* (2), *Ch. fimbriola hackeri* (3), *Ch. fimbriola vallensis* (4), *Ch. fimbriola fimbriola* (5), *Ch. fimbriola baloghi* (6), *Ch. fimbriola maravignae* (7), *Ch. fimbriola thurneri* (8), *Ch. fimbriola niculescui* (9), *Ch. fimbriola* ssp. (10), *Ch. fimbriola zernyi* (11), *Ch. fimbriola bohatschi* (6); *Ch. laeta leonhardi* (a), *Ch. laeta achaiana* (b), *Ch. laeta cretica* (c), *Ch. laeta macini* (d), *Ch. laeta laeta* (e), *Ch. laeta euxinia* (f). Taxonomic details: HACKER & VARGA 1990.




Fig. 17: Distribution of the species related to *Chersotis friedeli*: *Ch. friedeli* (1), *Ch. stenographa* (2), *Ch. gratissima* (3), *Ch. zagrosi* (4), *Ch. nupponenorum* (5), *Ch. nekrasovi* (6). (3) + (4) and (5) + (6) are allopatric sister species. Taxonomy and phylogeny: VARGA et al. 2013.

Another group of xeromontane genera probably has been originally connected to some xerophilous scrub formations as e.g. *Eugnorisma* (with its subgenus *Metagnorisma*), *Auchmis* (connected to *Berberis*), or *Lophoterges* (connected to *Lonicera*; RONKAY & VARGA

1997; RONKAY 2000, 2005). Their Mediterranean-Anatolian taxa display western and Central Asiatic connections (VARGA 2010b), and only a marginal speciation in the Mediterranean ranges. Oppositely, some other Mediterranean genera display an essen-

tially autochthonous evolution, influenced by the younger Tertiary aridisation of the Mediterranean Basin in connection with the Messinian Saline Crisis. The Ponto-Mediterranean (including parts of Asia Minor), the Atlanto-Mediterranean and Maghreb areas usually display a high level of species diversity in these genera. Examples can be mentioned from the tribe Oncocnemidini: *Omphalophana*, *Teinoptera*, *Cleonymia*, *Amephana*, *Metopoceras*, *Harpagophana*, etc. and from genera of Xylenini (*Leucochlaena*, *Aporophila*, *Antitype*, *Ammoconia*, subgenera of *Mniotype*, *Polymixis*, *Anchoscelis*, *Conistra*, etc.). Due to this paleo-ecological background this biogeographical group can be considered as equivalent of the “Paleo-Mediterranean-Xeromontane” faunal type of ornithologists (STEGMANN 1938, VOOUS 1960, 1963) and with the faunal type of the “ancient Mediterraneanum” of the Russian biogeographical school (BEY-BIENKO 1948, SHUMAKOV 1963, KRYZHANOVSKY 1965).

Discussion 2: Subdivision and history of the Palaearctic xeromontane fauna

In Europe and adjacent areas the majority of xeromontane species occur in the summer-dry Mediterranean mountain systems, from the Atlas Mts to Asia Minor (STEGMANN 1938, VOOUS 1960, KRYZHANOVSKY 1965, VARGA 1997). To the contrary, Central and Northern Europe just holds a few species of this faunal type and were only colonised by xeromontane species possibly during the cold-continental younger glacial and early post-glacial phases only (MIKKOLA et al. 1987; VARGA 1996, 2010a, 2010b). A much higher diversity of xeromontane species was observed from the arid high mountains of East Anatolia to the mountain systems of Central and Inner Asia, mostly due to the high species numbers of some typical genera e.g. in Noctuidae (subfamily Noctuinae), Lycaenidae: Polyommatus, Orthoptera: Catantopidae (VARGA 1996, 2010b; ECKWEILER & HÄUSER 1997; PRAVDIN & MISHTSHENKO 1980; ÜNAL 2016).

Thus, the composition of the xeromontane fauna is subdivided into a West Palaearctic (Mediterranean-xeromontane) and a Central- and Inner-Asiatic (Continental) sub-type. The protracted biogeographical confusion in the use of terms Central Asia vs. Middle Asia, or Inner Asia (used historically as “Innermost Asia”, or “Died Heart of Asia”, see: M. A. Stein 1928) has been clarified by the recent paleo-ecological surveys. They have shown a deep historical split between the more western Central Asiatic and the Inner Asiatic (Mongolian-Tibetan) steppe biota, both in the composition of vegetation, and in the mammalian assemblages (BARBOLINI et al. 2020). This split is clearly reflected by the differentiation of the faunal composition of the more western Central Asiatic vs. the cold-continental Inner Asiatic mountain systems. While the former

regions are significantly influenced by the sub-Mediterranean, equinoctial type of precipitation, the latter show an impoverished version of the continental type with scarce summer precipitation maxima (AGAKHANJANTS 1981, AGAKHANJANTS & BRECKLE 1995). This subdivision was also shown in the area-cladograms of the genus *Xenophysa* (VARGA 1989a, 2011). In the xeromontane fauna of the „western“ Central Asiatic mountains, some butterfly genera (e.g. *Parnassius*, *Karanasa*, *Paralasa*, subgenera of *Polyommatus*) and also typical genera of Noctuidae (e.g. *Euxoa*, *Dichagyris*, *Chersotis*, *Rhyacia*, *Eugnorisma*, *Goniographa*; and some oligotypic genera, e.g. *Hypsophila*, *Fergana*) predominate. The highest diversity of these genera mostly overlap with the core areas of the “saxobiotic” Orthoptera genera, see: Discussion 1.

Contrarily, in the eastern, Inner Asiatic group of the mountains, the typical butterflies belong to the genera *Colias*, *Oeneis* and *Boloria*, which also have xeromontane connections but penetrated into the tundral zonobiome. The Noctuidae of the Inner Asiatic xeromontane faunal type consist of several “Mongolian-Tibetan” cold-continental genera, e.g. *Trichosilia*, subgenera of *Xestia* (*Anomogyna*, *Pachnobia*, *Raddea*, *Estimata*, *Schoeyenia*), subgenera of *Anarta* (*Calocestra*, *Anarta* s. str.), *Lasionycta*, *Oncocnemis* and *Sympistis*, etc. Supposedly, the connections of this faunal type with the cold steppic and tundral zonobiomes have been mostly influenced by the recently uncovered Mid-Pleistocene climatic transition (DIESTER-HAAS et al. 2018; SUN et al. 2019; WILLEIT et al. 2019) which had deeply transformed the general zonality of the vegetation by the global decline of the primary productivity.

Hence, the connections of the xeromontane fauna must be also historically more manifold and partially more ancient than the oreotundral connections of the alpine faunal type, resulted mostly in the Quaternary climatic fluctuations and area dislocations (VARGA 1995, 1996). Thus, the fauna of the subtropical, monsoonic orobomes (e.g. in Southern China and the Himalayan region) displays a somewhat less differentiated, hypothetically more ancestral character. This regularity is clearly unfolding from the ranges of Poliina genera/subgenera *Tricheurois*, *Haderonia* and *Metallopolia* displaying the species diversity restricted to the Sino-Himalayan area, as opposed to the more derived genera *Polia* s. str. and *Ctenoceratoda* (VARGA et al. 2019, Fig. 20).

We suppose that a bulk of basal groups of the fauna have evolved, jointly with some groups of *Angiospermae*, in the Eastern Gondwana (see: theory of AXELROD 1960). They could expand northwards after the earliest collision of the Gondwanian plates with south-eastern Asia (e.g. Southern Tibet and parts of south-western China) forming an important source of the Sino-Himalayan core area of biodiversity as was also shown e.g. in the Passerine birds (FJELDSÅ 2013; CAI et al. 2019). A high diversity of cold-adapted ancestral species originating from the seasonally humid, monsoonic

southeastern Asiatic mountains have been constrained by passing two main filter-corridors (Fig. 21, modified from VARGA 1995).

- The „*Rhododendron*-corridor”, being characteristic for several evergreen *Angiospermae*, e.g. *Vaccinium*, *Empetrum*, *Rhododendron* incl. *Ledum*, etc., which now compose a major part of the undergrowth of the Siberian (especially southern Siberian mountain) taiga, typical for a number of “taiga-birds” (e.g. *Ficedula*, *Phylloscopus*, *Luscinia calliope*, *Tarsiger cyanurus*, *Dumetica thoracica* etc.) and also for numerous taiga-inhabiting Lepidoptera (Geometridae: *Dysstroma*, *Chlorochlysta*; Noctuidae: *Xestia* subg. *Pachnobia*, *Anomogyna*; *Polia*, *Lasionycta*, species-groups of *Apamea* etc., Figs 20–21). The endemic species of these groups mostly inhabit the southern Siberian mountain taiga, while some of their relatives are widely dispersed, often having a Trans-Palaearctic or even Holarctic range of distribution.
- The „xeromontane” route, leading from the Trans-Himalayan mountains to the Karakoram, Pamir, Transalai, or to the Hissar, Seravshan, West Tien-Shan and Hindukush ranges, on one hand, or via East-Turkestan to the Eastern Tien-Shan and Altai-Sayan systems, on the other, resulting in a radiation of a rich continental xeromontane fauna. This bifurcation becomes apparent from the taxonomic division of the genera typical for these areas. In the fauna of the first group of the Inner Asiatic mountains, some butterflies (*Parnassius*, *Karanasa*, *Paralasa*, numerous Polyommatus genera) and oligotypic genera of Noctuidae (*Hypsophila*, *Fergana*) predominate. In the second group of the mountains the typical butterflies belong to *Colias*, *Oeneis* and *Boloria*, which supposedly have a xeromontane origin but penetrated into the tundral zonobiomes. The typical genera of this second, more “Siberian”, group of (originally) xeromontane Noctuidae are e.g. *Trichosilia*, *Lasionycta*, *Discestra*, *Anarta* and some species groups of the *Oncocnemis*-*Sympistis* generic complex.

Based on these biogeographical facts, the connections of the Continental-Inner-Asiatic xeromontane fauna should be historically more manifold and ancient than the “typical” oreotundral connections of the alpine faunal type which can be mostly regarded as a consequence of the younger Quaternary climatic fluctuations and area dislocations.

Acknowledgements

Author is highly indebted to L. Ronkay, L. Rákosi and Th. Schmitt for the long-year cooperation, for critical comments and suggestions to the earlier version of the manuscript. I am especially grateful to G. Ronkay for the excellent pictures of the mapped Noctuidae species.

The research was supported by a senior fellowship in the Collegium Budapest, Institute of Advanced Study (2001), by the Re-Invitation Grant of the Alexander von Humboldt Foundation (Zoological State Collection in Munich, 2007), by the OTKA Foundation Grant No. 16465, by the Grant No. K116694 of the National Foundation of Research, Development and Innovation (NKFI-OTKA) and by the SYNTHESYS Project DE-TAF 6708.

References

- AGAKHANJANTS, O. E. 1981: The Arid High Mountains of the URSS (in Russian). – Moscow, Izd. Mysl': pp. 170.
- AGAKHANJANTS, O. E. & BRECKLE, S.-W. 1995: Origin and evolution of the mountain flora in Middle Asia and neighbouring mountain regions. – In: CHAPIN, F. S. & KÖRNER, C. (eds.) Arctic and Alpine Biodiversity: Patterns, Causes and Ecosystem Consequences. – Springer, Berlin-New York, pp. 63–80. – https://doi.org/10.1007/978-3-642-78966-3_5.
- AXELROD, D. I. 1970: Mesozoic paleogeography and early angiosperm history. – The Botanical Review **36**: 277–319.
- BARBOLINI, N.; WOUTERSEN, A.; DUPONT-NIVET, G. et al. 2020: Cenozoic evolution of the steppe-desert biome in Central Asia. – Science Advances 2020; **6**: eabb8227 9 October 2020.
- BEER, C.; REICHSTEIN, M.; TOMEMLERI, E.; CIAIS, P.; JUNG, M.; CARVALHAIS, N.; RÖDENBECK, C.; ARAIN, M. A.; BALDOCCHI, D. & BONAN, G. B. 2010: Terrestrial gross carbon dioxide uptake: Global distribution and covariation with climate. – Science **329**: 834–838.
- CAI, T.; SHAO, S.; KENNEDY, J. D.; ALSTRÖM, P.; MOYLE, R. G.; QU, Y.; LEI, F. & FJELDSÅ, J. 2020: The role of evolutionary time, diversification rates and dispersal in determining the global diversity of a large radiation of passerine birds. – Journal of Biogeography **47**: 1612–1625.
- DIESTER-HAAS, L.; BILLUPS, K. & LEAR, C. 2018: Productivity changes across the mid-Pleistocene climate transition. – Earth-Science Reviews **179** (2018): 372–391.
- DUFAY, C. & VARGA, Z. 1995: A new noctuid species from Iran, *Chersotis eberti* sp. n. (Lep.: Noctuidae). – Acta Zoologica Academiae Scientiarum Hungaricae **41**: 35–45.
- ECKWEILER, W. & HÄUSER, C. L. 1997: An illustrated checklist of *Agrodiaetus* HÜBNER, 1822, a subgenus of *Polyommatus* LATREILLE, 1804 (Lepidoptera, Lycaenidae). – Nachrichten des entomologischen Vereins Apollo 1997: 113–166.
- ENGHOFF, H. 1995: Historical biogeography of the Holarctic: area relationships, ancestral areas and dispersal of non-marine animals. – Cladistics **11**: 223–263.

- FIBIGER, M. 1990: Noctuidae Europaeae. Vol. 1, Noctuinae I. – Entomological Press, Sorø: pp. 207.
- FIBIGER, M. 1993: Noctuidae Europaeae. Vol. 2, Noctuinae II. – Entomological Press, Sorø: pp. 230.
- FJELDSÅ, J. 2013: The global diversification of songbirds (Oscines) and the build-up of the Sino-Himalayan diversity hotspot. – Chinese Birds 4 (2):132–143. – DOI10.5122/cbirds.2013.0014.
- HAROLD, A. S. & MOOI, R. D. 1994: Areas of endemism: definition and recognition criteria. – Systematic Biology 43 (2): 261–266.
- HUEMER, P. 1998: Endemische Schmetterlinge der Alpen – ein Überblick (Lepidoptera). – Stafnia 55: 229–256.
- HUEMER, P. & KARSHOLT, O. 2020: Commented checklist of European Gelechiidae (Lepidoptera). – ZooKeys 921: 65–140. – <https://doi.org/10.3897/zookeys.921.49197>.
- HUEMER, P.; KARSHOLT, O. & MUTANEN, M. 2014: DNA barcoding as a screening tool for potential unrecognized cryptic diversity exemplified by the genus *Caryocolum*, with description of a new species (Lepidoptera, Gelechiidae). – ZooKeys 404: 91–111. – <https://doi.org/10.3897/zookeys.404.7234>.
- HUEMER, P. & MUTANEN, M. 2015: Alpha taxonomy of the genus *Kessleria* NOWICKI, 1864, revisited in light of DNA-barcoding (Lepidoptera, Yponomeutidae). – ZooKeys 503: 89–133. – <https://doi.org/10.3897/zookeys.503.9590>.
- HUEMER, P. & TIMOSSI, G. 2014: *Sattleria* revisited: unexpected cryptic diversity on the Balkan Peninsula and in the south-eastern Alps (Lepidoptera: Gelechiidae). – Zootaxa 3780 (2): 282–296. – <https://doi.org/10.11646/zootaxa.3780.2.4>.
- KANDUL, N. P.; LUKHTANOV V. A.; DANTCHENKO, A. V.; COLEMAN, J. W. S.; SEKERCIOGLU, C. H.; HAIG, D. & PIERCE, N. E. 2004: Phylogeny of *Agrodiaetus* HÜBNER, 1822 (Lepidoptera: Lycaenidae) inferred from mtDNA sequences of *COI* and *COII* and nuclear sequences of *EF1-α*: karyotype diversification and species radiation. – Systematic Biology 53: 278–298. – doi:10.1080/10635150490423692.
- KANDUL, N. P.; LUKHTANOV, V. A. & PIERCE, N. E. 2007: Karyotypic diversity and speciation in *Agrodiaetus* butterflies. – Evolution 61: 546–559. – doi:10.1111/j.1558-5646.2007.00046.x.
- KRYZHANOVSKY, O. A. 1965: Composition and Origin of the Terrestrial Fauna of Central Asia. (in Russian) – Nauka, Moscow-Leningrad: pp. 419.
- KRYZHANOVSKY, O. A. 1969: A contribution to the composition and origin of alpine faunas of Coleoptera (in Russian). – Zoologicheskiy Zhurnal 48: 1156–1165.
- LAFONTAINE, J. D. 1981: Classification and phylogeny of the *Euxoa detersa*-group (Lep.: Noctuidae). – Quaestiones entomologicae 17: 1–120.
- LAFONTAINE, J. D. 1987: Noctuoidea, Noctuidae (part), Noctuinae (part *Euxoa*). – In: The Moths of America North of Mexico including Greenland. – The Wedge Foundation, Washington, fasc. 27: pp. 1–237.
- LAFONTAINE, J. D. & KONONENKO, V. S. 1986: A revision of the genus *Trichosilia* HAMPSON (Lepidoptera: Noctuidae) with descriptions of four new species. – Canadian Entomologist 118 (11): 1079–1113.
- LAFONTAINE, J. D. & KONONENKO, V. S. 1988: A revision of the *Lasionycta skraelingia* (HERRICH-SCHAEFFER) species complex (Lepidoptera: Noctuidae). – Canadian Entomologist 120: 903–916.
- LAFONTAINE, J. D. & MIKKOLA, K. 1987: Lock-and-key systems in the inner genitalia of Noctuidae (Lepidoptera) as a taxonomic character (in Swedish with English summary) – Entomologisk Meddelser 55: 161–167.
- LAFONTAINE, J. D.; MIKKOLA, K. & KONONENKO, V. S. 1983: A revision of the genus *Xestia* subg. *Schoeyenia* Aurivillius (Lep.: Noctuidae) with descriptions of four new species and a new subspecies. – Entomologica scandinavica 14: 337–369.
- LAFONTAINE, J. D.; MIKKOLA, K. & KONONENKO, V. S. 1987: A revision of the genus *Xestia* subg. *Pachnobia* (Lep.: Noctuidae) with descriptions of two new subspecies. – Entomologica scandinavica 18: 305–331.
- LAFONTAINE, J. D. & WOOD, D. M. 1988: A zoogeographic analysis of the Noctuidae (Lepidoptera) of Beringia and some inferences about past Beringian habitats. – Memoires of the entomological Society of Canada 144: 109–123.
- LI, B. P.; LIU, Z. W. & ZHENG Z-M. 2011: Phylogeny and classification of the Catantopidae at the tribal level (Orthoptera, Acridoidea). – ZooKeys 148: 209–255. doi:10.3897/zookeys.148. 2081.
- LUKHTANOV, V. A.; KANDUL, V. A.; PLOTKIN, J. B.; DANTCHENKO, A. V.; HAIG, D. & PIERCE, N. E. 2005: Reinforcement of pre-zygotic isolation and karyotype evolution in *Agrodiaetus* butterflies. – Nature 436: 385–389. – doi: 10.1038/nature03704.
- MIKKOLA, K. 1992: Evidence for lock-and-key mechanisms in the internal genitalia of the Apamea moths (Lep.: Noctuidae). – Systematic Entomology 17: 145–153.
- MIKKOLA, K.; LAFONTAINE, J. D. & GROTFELT, P. 1987: A revision of the Holarctic *Chersotis andereggi* complex (Lep.: Noctuidae). – Nota lepidopterologica 10: 140–157.
- MIKKOLA, K.; LAFONTAINE, D. J. & KONONENKO, V. S. 1991: Zoogeography of the Holarctic species of the Noctuidae (Lepidoptera): importance of the Beringian refuge. – Entomologica fennica 2: 158–173.
- MÜLLER, B.; ERLACHER, S.; HAUSMANN, A.; RAJAEI, H.; SIHVONEN, P. & SKOU, P. 2019: The Geometrid Moths of Europe. Volume 6. Subfamily Ennominae II (Boarmiini, Gnophini, additions to previous volumes). – Part 1: 1–562, part 2: 563–906. – Brill, Leiden & Boston.
- PAPPAS, C.; FATICHI, S. & BURLANDO, P. 2016: Modeling terrestrial carbon and water dynamics across climatic gradients: does plant trait diversity matter? – New Phytologist 209: 137–151. – doi: 10.1111/nph.13590.

- PRAVDIN, F. N. & MISHTSHENKO, L. L. 1980: Formation and evolution of the ecological faunal types of insects in Central Asia. – Nauka, Moscow: pp. 156. [In Russian].
- PRINCE, S. D. & GOWARD, S. N. 1995: Global Primary Production: A Remote Sensing Approach. – Journal of Biogeography **22** (4/5): 815–835.
- RYABOV, M. A. 1956: Types of annual life-cycles in cut-worms (Lepidoptera, Agrotidae) (in Russian). – Revue d'Entomologie de l'URSS (Entomologicheskoe Obozrenie) **35**: 70–79.
- RODIN, L. E. & BAZILEVIC, N. I. 1966: The biological productivity of the main vegetation types in the Northern Hemisphere of the Old World. – Forestry Abstracts **27**: 365–375.
- RONKAY, L. 2000: Revision of the genus *Lophoterges* HAMPSON, 1906 (s. l.) (Lepidoptera, Noctuidae, Cuculliinae). Part I. – Annales historico-naturales Musei nationalis hungarici **92**: 193–213.
- RONKAY, L. & VARGA, Z. 1997: New taxa of the genera *Auchmis* HÜBNER [1821] and *Nekrasovia* RONKAY & VARGA, 1993. (Lepidoptera, Noctuidae). – Acta Zoolo Academiae Scientiarum Hungaricae **43**: 149–161.
- RONKAY, L. & VARGA, Z. 1999: Revision of the genus *Eugnorisma* BOURSIN, 1946, Part V. New genera and species of *Eugnorisma* genus group from Pakistan and China (Lepidoptera, Noctuidae). – Acta Zoologica Academiae Scientiarum Hungaricae **45** (4): 345–373.
- RUNNING, S. W.; NEMANI, R. R.; HEINSCH, F. A.; ZHAO, M.; REEVES, M. & HASHIMOTO, H. 2004: A continuous satellite-derived measure of global terrestrial primary production. – BioScience **54** (6): 547–560.
- SAULICH, A. KH.; SOKOLOVA, I. V. & MUSOLIN, D. C. 2017: Seasonal Cycles of Noctuid Moths of the Subfamily Plusiinae (Lepidoptera, Noctuidae) of the Palaearctic: Diversity and Environmental Control. – Entomological Review **97** (2): 143–157.
- SHUMAKOV, E. M. 1963: Locusts of Afghanistan and Iran. (In Russian). – Trudy Vsesoyuznogo Entomologicheskogo Obstshchestva **49**: 1–248.
- SPECHT, A.; ANGULO, A. O.; OLIVARES, T. S.; FRONZA, E.; ROQUE-SPECHT, V. F.; VALDUGA, E.; ALBRECHT, F.; POLETTI, G. & BARROS, N. M. 2013: Life cycle of *Agrotis malefida* (Lepidoptera: Noctuidae): a diapausing cutworm. – Zoologia **30** (4): 371–378.
- STEGMANN, B. 1938: Grundzüge der ornithogeographischen Gliederung des paläarktischen Gebietes. – Fauna SSSR, nov. ser. **19**: pp. 158.
- STEGMANN, B. 1958: Herkunft der europäischen Steppevögel. – Bonner zoologische Beiträge **9**: 208–230.
- STEIN, A. M. Sir 1928: Innermost Asia: Detailed Report of Explorations in Central Asia, Kan-su and Eastern Iran, Vol. 1–5. – Oxford, Clarendon Press. (Reprint: New Delhi, Cosmo Publications, 1981).
- SUKHAREVA, I. L. 1999: Family Noctuidae. – In: KUZNETSOV, V. I. (ed.): Insects and Acarines as Agricultural Pests, Vol. 3: Lepidoptera, Part 2. (In Russian). – Nauka, St. Petersburg: pp. 332–376.
- SUN, Y.; YIN, Q.; CRUCIFIX, M. et al. & AN, ZH. 2019: Diverse manifestations of the mid-Pleistocene climate transition. – Nature Communications **10**: 352. – <https://doi.org/10.1038/s41467-018-08257-9>.
- ÜNAL, M. 2016: Pamphagidae (Orthoptera: Acridoidea) from the Palaearctic Region: taxonomy, classification, keys to genera and a review of the tribe Nocarodeini I. Bolívar. – Zootaxa **4206** (1): 1–223. – ISBN 978-1-77670-047-9.
- VARGA, Z. 1977: Das Prinzip der areal-analytischen Methode in der Zoogeographie und die Faunenelemente-Einteilung der europäischen Tagschmetterlinge (Lep.: Diurna). – Acta Biologica Debrecina **14**: 223–285.
- VARGA, Z. 1989a: Zweiter Beitrag zur Kenntnis der Gattung *Xenophysa* BOURSIN, 1969 (Lep.: Noctuidae) mit Beschreibung fünf neuer Arten. – Zeitschrift der Arbeitsgemeinschaft österreichischer Entomologen **41** (1–2): 48–77.
- VARGA, Z. 1989b: The origin and division of the northern-montane disjunct areas in Palaearctic Lepidoptera – Their importance in solving zoogeographical and evolutionary problems. – Acta Biologica Debrecina **21**: 91–116.
- VARGA, Z. 1990: New species of Noctuidae from Afghanistan and adjacent territories I. The Genera *Euxoa* HÜBNER (1821) 1816, *Dichagyris* LEDERER, 1857, *Rhyacia* HÜBNER (1821) 1816. (Lep.: Noctuidae, Noctuinae). – Esperiana (Schwanfeld) **1**: 167–198.
- VARGA, Z. 1992: Taxonomic notes on the Genus *Haderonia* STAUDINGER, 1896 with the description of a new Genus *Ctenoceratoda* and four new species (Lep.: Noctuidae). – Acta Zoologica Academiae Scientiarum Hungaricae **38** (1–2): 113–124.
- VARGA, Z. 1993: Beiträge zur Kenntnis der Gattung *Dichagyris* LEDERER, 1857 (Lep.: Noctuidae) I. – Acta Zoologica Academiae Scientiarum Hungaricae **39** (1–4): 289–300.
- VARGA, Z. 1995a: Isolates of Arctic-alpine and Alpine Lepidoptera in SE Europe. – Proceedings of the EIS Colloquium, Helsinki: pp. 140–151.
- VARGA, Z. 1995b: Geographical Patterns of Biological Diversity in the Palearctic Region and the Carpathian Basin. – Acta Zoologica Academiae Scientiarum Hungaricae **41** (2): 71–92.
- VARGA, Z. 1996a: Entomologische Aspekte der räumlichen und biotischen Diversität in einer mitteleuropäischen Mosaiklandschaft. – Verh. 14. int. Symposium für Entomofaunistik in Mitteleuropa, SIEEC, München (1994): p. 33–67.
- VARGA, Z. 1996b: New species and subspecies of *Dichagyris* LEDERER 1857, *Chersotis* BOISDUVAL 1840 and *Rhyacia* HÜBNER [1821] 1816 (Noctuidae, subfam. Noctuinae) from Central Asia. – Acta Zoologica Academiae Scientiarum Hungaricae **42** (3): 195–230.

- VARGA, Z. 1997: Biogeography and Evolution of the boreal Lepidoptera in the Palearctic. – *Acta Zoologica Hungarica Academiae Scientiarum Hungaricae* **42** (4): 289–330.
- VARGA, Z. 1998: Sibling species and species groups in the genus *Chersotis* BOISDUVAL, 1840 (Lepidoptera, Noctuidae: Noctuinae) with description of two new species. – *Acta Zoologica Academiae Scientiarum Hungaricae* **44** (4): 341–372.
- VARGA, Z. 2003: The geographical distribution of high mountain Macrolepidoptera in Europe. – In: GRABHERR, G. & NAGY, L. (eds): Alpine biodiversity in Europe. – Springer-Verlag: pp. 239–257. – https://doi.org/10.1007/978-3-642-18967-8_13.
- VARGA, Z. 2010a: Extra-Mediterranean refugia, post-glacial vegetation history and area dynamics in Eastern Central Europe. – In: HABEL, J. & ASSMANN, TH. (eds.): Relict species: Phylogeography and Conservation Biology. – Springer-Verlag: pp. 57–87.
- VARGA, Z. 2010b: Biogeography of West Palaearctic Noctuidae. – In: FIBIGER, M.; RONKAY, L.; YELA, J. L. & ZILLI, A.: Noctuidae Europaea **12** (incl. Suppl. 1–12): pp. 265–274. – Entomological Press, Sorø.
- VARGA, Z. 2011: Revision of the genus *Xenophysa* BOURSIN, 1969 (Lepidoptera, Noctuidae). – *Zootaxa* **3094**: 1–29. – www.mapress.com/zootaxa/.
- VARGA, Z. 2014: New and revised *Euxoa* HÜBNER, 1821 species from Western and Central Asia (Lepidoptera, Noctuidae, Noctuinae). – *Fibigeriana Suppl.* **2**: pp. 9–30. Heterocera Press, Budapest.
- VARGA, Z. & GYULAI, P. 1999: Taxonomy of the genus *Ctenoceratoda* VARGA, 1992 (Lepidoptera, Noctuidae) with the description of seven new species. – *Acta Zoologica Academiae Scientiarum Hungaricae* **45** (2): 169–197.
- VARGA, Z.; GYULAI, P.; RONKAY, G. & RONKAY, L. 2013: Noctuinae I. A Taxonomic Atlas of the Eurasian and North African Noctuoidea. – The Witt Catalogue, Vol. **6**. Heterocera Press, Budapest: pp. 394.
- VARGA, Z.; GYULAI, P.; RONKAY, G. & RONKAY, L. 2015: Noctuinae II. A Taxonomic Atlas of the Eurasian and North African Noctuoidea. – The Witt Catalogue, Vol. **8**. Heterocera Press, Budapest: pp. 394.
- VARGA, Z. & RONKAY, L. 1987: Revision of the Genus *Eugnorisma* BOURSIN, 1946 (Lep.: Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **33**: 187–262.
- VARGA, Z. & RONKAY, L. 1996: New and Revised Taxa from Genera *Chersotis* BOISDUVAL, 1840 and *Dichagyris* LEDERER, 1857 from Central Asia (Lep.: Noctuidae). – *Esperiana* (Schwanfeld) **5**: 103–132.
- VARGA, Z. & RONKAY, L. 2002: A revision of the Palaearctic species of the *Eugrapha* HÜBNER, [1821] 1816 generic complex. Part I. The genera *Eugrapha* and *Goniographa* (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **49**: 333–374.
- VARGA, Z.; RONKAY, L. & HACKER, H. 1990: Revision of the Genus *Bryopolia* BOURSIN, 1954 (Lep.: Noctuidae). – *Esperiana* (Schwanfeld) **1**: 427–469.
- VARGA, Z.; RONKAY, L. & YELA, J. L. 1990: Revision of the Genus *Eugnorisma* BOURSIN, 1946. Part II. Taxonomic news, biogeographic and phylogenetic considerations and descriptions of two new genera: *Ledereragrotis* and *Pseudohermenassa* (Lep.: Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **36** (3–4): 331–360.
- VOOUS, K. H. 1960: Atlas of European birds. – Nelson, London: pp. 284.
- VOOUS, K. H. 1963: The concept of faunal elements or faunal types. – Proceedings of the 13 international Congress of Ornithology: pp. 1104–1108.
- WILLEIT, M.; GANOPOLSKI, A.; CALOV, R. & BROVKIN, V. 2019: Mid-Pleistocene transition in glacial cycles explained by declining CO₂ and regolith removal. – *Science Advances* 2019, **5**: eaav7337.


Tab. 1: Allopatric pairs and groups of *Dichagyris* (subg. *Dichagyris*) species (*with peripatric overlap, *polytypic species).

Western species	Eastern species
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>vallesiaca</i> (Boisduval, 1837) ⁴ : SW and Central Alps, Crimea, South Russia, South Urals, Asia Minor, Zagros, Elburs, Binaloud Mts, Kopet-Dagh, Tien Shan, Pamirs, Altai, Khangai, Sayan Mts.	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>fuscashmiriana</i> Varga, Ronkay & Ronkay, 2020: Hindukush East, Karakoram, W Himalayas
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>striata</i> Kozhantschikov*, 1930: Zagros, Elburs, Binaloud Mts, Kopet-Dagh	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>tyrannus</i> (A. Bang-Haas, 1912)*: Tien Shan, Hissaro-Darwaz, Pamirs, Hindukush, Karakoram, W Himalayas
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>squalidior</i> (Staudinger, 1901) ⁵ : Asia Minor, Zagros, Elburs, Binaloud Mts, Kopet-Dagh	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>payotiorum</i> Varga, Ronkay & Ronkay: Hissaro-Darwaz, Hindukush, Paghman Mts. Karakoram, W Himalayas
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>clara</i> (Staudinger, 1888)*: Tien Shan, Mongol Altai, Gobi Altai	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>leucographa</i> Varga, 1990: Pamirs <i>Dichagyris</i> (<i>Dichagyris</i>) <i>scotographa</i> Varga, 1990: Pamirs, Karakoram, W Himalayas (Kashmir) <i>Dichagyris</i> (<i>Dichagyris</i>) <i>kautti</i> Varga, 1996: W Himalayas (Ladakh)
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>jacobsoni</i> Kozhanchikov, 1930: Kopet-Dagh	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>umbrifera</i> (Alphéraky, 1882): Tien Shan
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>naumannii</i> Varga, 1996: Kugitang-Tau, Hindukush	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>kaszabi</i> Varga, 1973: Mongol & Gobi Altai
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>cataleipa</i> Varga, 1993: Eastern Asia Minor	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>afghana</i> Boursin, 1963: Central Hindukush
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>psammochroa</i> (Boursin, 1940)*: Zagros, Elburs, Kopet-Dagh	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>kurbatskyi</i> Varga, Ronkay & Ronkay, 2021: W Tien Shan <i>Dichagyris</i> (<i>Dichagyris</i>) <i>apochora</i> Gyulai & Varga, 2001: Pamirs
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>taftana</i> Brandt, 1941*: Elburs, Zagros	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>guenterbereti</i> Varga, Ronkay & Ronkay, 2021: Central Hindukush
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>humilis</i> (Boursin, 1940): Zagros, Elburs, Kopet-Dagh	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>hypotacta</i> Varga, Ronkay & Ronkay 2021: East Tien Shan, Pamirs, Hindukush
<i>Dichagyris</i> (<i>Dichagyris</i>) <i>verecunda</i> (Püngeler, 1898)*: Tien Shan system, Pamirs, Mongol Altai, Gobi Altai	<i>Dichagyris</i> (<i>Dichagyris</i>) <i>karakorealis</i> Varga, Ronkay & Ronkay, 2020: West Himalayas

⁴ Sympatric endemic species: *Dichagyris* (*Dichagyris*) *griseotincta* (Wagner, 1931) – Central Anatolia.⁵ Northern sister species (?subspecies): *Dichagyris* *lux* Fibiger & Nupponen, 2002 – South Urals.

Tab. 2: Allopatric pairs of *Euxoa* species (exceptionally with peripatric overlap).

Western species	Eastern species
<i>Euxoa homicida</i> (Staudinger, 1900)*: Asia Minor, Transcaucasia, Zagros, Elburs	<i>Euxoa transcaspica</i> Kozhantshikov, 1928*: Kopet-Dagh, Binaloud Mts (ssp.); Hindu Kush, Karakoram, W. Himalayas (ssp.)
<i>Euxoa decora</i> ([Denis & Schiffermüller], 1775): Pyrenees, Alps, Carpathians, Europa Centr., Apennines, Balkans	<i>Euxoa kurushensis</i> Boursin, 1940: Daghestan (NE Caucasus)
<i>Euxoa heringi</i> (Staudinger, 1877): Asia Minor, Elburs, Kopet-Dagh	<i>Euxoa cretaporos</i> Varga, G. Ronkay & L. Ronkay, 2020: Daghestan (NE Caucasus)
<i>Euxoa aneucta</i> Brandt, 1938*: Zagros Mts, Binaloud Mts (ssp.)	<i>Euxoa subeucta</i> Varga, 2014: Hissar Mts., Transalai Mts, Hindu Kush Central & East
<i>Euxoa perierga</i> Brandt, 1938*: Zagros Mts, Elburs (ssp.)	<i>Euxoa dichagyroides</i> Varga, 1979: Darwaz Mts., Pamirs, Transalai Mts, Hindu Kush Central & East
<i>Euxoa sigma</i> Kozhantshikov, 1928: Kopet-Dagh, Binaloud Mts	<i>Euxoa metasigmata</i> Varga, 2014: Hissar Mts, Hindu Kush Central & East
<i>Euxoa conifera</i> Christoph, 1885* ⁶ : Daghestan, Transcaucasia, Asia Minor (East), Elburs, Kopet-Dagh	<i>Euxoa bactriana</i> Varga, 2014: Tien Shan (Seravshan), Transalai, Hissar Mts, Pamirs, Hindu Kush Central & East, Karakoram, West Himalayas
<i>Euxoa bogdanovi</i> (Erschoff, 1873): Tien Shan	<i>Euxoa xanthophylla</i> Varga, 1990: Darwaz Mts, Pamirs

Fig. 18: Early autumnal steppic *Euxoa* species: *E. distinguenda* (1), *E. christophi* (2).⁶ Senior synonym of *E. difficillima* Draudt, 1937.

Tab. 3: Allopatric pairs of *Chersotis* species (exceptionally with peripatric overlap).

Western species	Eastern species
<i>Ch. firdusii</i> Schwingenschuss, 1937: East Anatolia, Zagros, Elburs, Binaloud Mts, Kopet-Dagh	<i>Ch. fidahusseini</i> Varga, Gyulai, Ronkay & Ronkay, 2013: E Tien Shan, Hissar and Darwaz range, Central and East Hindukush, Karakoram, West Himalayas
<i>Ch. sordescens</i> (Staudinger, 1900): Tien-Shan range, Hindukush, Hissar and Darwaz range	<i>Ch. herczigi</i> Varga, 1996: West Himalayas
<i>Ch. ronkayorum</i> Fibiger, Hacker & Varga, 1993: Central and East Anatolia	<i>Chersotis sterilis</i> Brandt, 1938: Zagros Mts
<i>Ch. nitens</i> Brandt, 1941: Binaloud Mts, Kopet-Dagh	<i>Ch. metrapha</i> Varga, 1975: Afghanistan, Hindukush Mts, Badakhshan
<i>Ch. capnistis</i> (Lederer, [1870]): South Russia, Caucasus, Transcaucasia, Asia Minor, South Balkans	<i>Ch. leucostola</i> Varga & Ronkay, 1996: Tien-Shan Mts, Dzhungarian Ala-too
<i>Ch. semna</i> (Püngeler, 1906): Asia Minor, Trans-caucasia, Elburs, Binaloud Mts, Kopet-Dagh	<i>Ch. pachnosa</i> Varga, 1975: Afghanistan, Badakhshan
<i>Ch. juvenis</i> (Staudinger, 1901): Asia Minor, Transcaucasia, Elburs Mts, Zagros Mts.	<i>Ch. kouros</i> Varga & Ronkay, 1996: Kopet-Dagh, Hindukush
<i>Ch. calorica</i> (Corti, 1930): North & East Tien-Shan Mts	<i>Ch. shandur</i> Varga, 1998: Pakistan, Karakoram Mts, Afghanistan, Paghman Mts
<i>Ch. petermarci</i> Varga, 1998: Seravshan Mts, W Tien-Shan Mts	<i>Ch. vicina</i> (Corti, 1930): N and E Tien-Shan Mts
<i>Chersotis delear</i> Boursin, 1970 ⁷ : Central Afghanistan, Nimla	<i>Ch. electrographa</i> Varga, 1990: Afghanistan, Darwaz Mts. <i>Ch. vargai</i> Hacker, 1992 Hindukush Mts, Karakoram
<i>Ch. binaloudi</i> Brandt, 1941 ⁸ : Khorassan, Binaloud Mts	<i>Ch. antigrapha</i> Boursin, 1961: Hindukush Mts <i>Ch. argyllographa</i> Varga & Gyulai, 2001: W Pamirs
<i>Ch. elegans</i> (Eversmann, 1837): Iberian Mts, Pyrenees, Alps, Balkans, Asia Minor, Transcaucasia, S Russia, W Tien-Shan and W Altai Mts	<i>Ch. eberti</i> Dufay & Varga, 1995: Elburs Mts, Zagros Mts
<i>Ch. sarhada</i> Brandt, 1941 ⁹ : Asia Minor, Transcaucasia, Elburs and Kopet-Dagh Mts	<i>Ch. lehmanni</i> Varga, Gyulai, Ronkay, Ronkay, 2010: Hindukush, Hissar and Darwaz Mts <i>Ch. lukhtanovi</i> Varga & Gyulai, 2001: S Tadjikistan
<i>Ch. rungsi</i> Boursin, 1944: Atlas Mts	<i>Ch. stenographa</i> Varga, 1979: Asia Minor East, Transcaucasia
<i>Ch. gratissima</i> (Corti, 1932)	<i>Ch. zagroica</i> Gyulai & Varga, 2006: Zagros Mts
<i>Ch. nupponenorum</i> Varga, Gyulai, Ronkay & Ronkay, 2013: W. Transcaspia	<i>Ch. nekrasovi</i> Varga, 1996: Pamir Mts

⁷ A „triplet” of species.⁸ A „triplet” of species.⁹ A „triplet” of species.

Tab. 4: Partly sympatric pairs and groups of *Chersotis* species.

<i>Ch. rectangula</i> ([Denis & Schiffermüller], 1775): Central and Southern Europe, Caucasus, Asia Minor	<i>Ch. andereggii</i> (Boisduval, [1837]): N Europe, Alps, Balkans, Asia Minor, Caucasus, Transcaucasia, Iran, Mts of Southern Siberia <i>Ch. acutangula</i> (Staudinger, 1892) ¹⁰ : Transalai and Hindukush Mts
<i>Ch. hahni</i> (Christoph, 1885): Transcaucasia, Elburz, Kopet-Dagh Mts	<i>Ch. curvispina</i> Boursin, 1961: Kopet-Dagh Mts, Hindukush Mts
<i>Ch. elegans</i> (Eversmann, 1837): Iberian Mts, Pyrenees Alps, Balkans, Asia Minor, Caucasus, Transcaucasia, S Russia, W Tien-Shan and W Altai Mts <i>Ch. eberti</i> Dufay & Varga, 1995: Elburz Mts, Zagros Mts	<i>Ch. anatolica</i> (Draudt, 1936) ¹¹ : Iberian Mts, Pyrenees, SW Alps, Central Apennines, Balkans, Asia Minor, Caucasus, Transcaucasia, S Russia (Europa + Siberia), Kopet-Dagh, W Tien-Shan Mts
<i>Ch. fimbriola</i> (Esper, [1803]): Atlas, Iberian Mts, Pyrenees, Sicily, W Alps, Pannonian Basin, Balkans Asia Minor, Crimea, Transcaucasia, Zagros, Elburz Mts, Kopet-Dagh Mts	<i>Ch. laeta</i> (Rebel, 1904): Balkans, W coastal area of Black Sea, Asia Minor, Caucasus, Transcaucasia, Elburz Mts




Fig. 19: Adaptation to the arid environment in Noctuinae genera: two types of univoltine life-cycles with autumnal period of oviposition and dispersal. Type I. (e.g. *Euxoa*, *Eugnorisma* – aestivating pupae, late summer, early autumn imagoes), type II. (e.g. *Chersotis*, *Spaelotis*, *Standfussiana*, aestivating imagoes).

¹⁰ Latter species is strictly localised in the Tien-Shan and Hindukush mountain systems.

¹¹ These species are not sibling species, because they are strongly differentiated from each other (see: Dufay & Varga 1995).





Fig. 20: General distribution of Poliina (Noctuidae) genera with primary centre of diversity (light blue) in the monsoonic mountainous areas of South-Eastern Asia (Himalayan – Sino-Tibetan faunal type, *Tricheurois*, *Haderonia*, *Metallopolia*) and secondary core area (light green) in the South Siberian mountains (*Polia* spp.). The main faunal corridor is extended from the Himalaya to the South Siberian mountain systems (arrows).



Appendix I

Biogeographical types of xeromontane Noctuidae (with special regard to *Noctuinae* genera: *Actebia* s.l., *Dichagyris* s.l., *Euxoa* s.l., *Rhyacia* s.l., *Chersotis*, *Eugnorisma* s.l., and *Xenophysa*) with taxonomic notes

1. Mediterranean- (Western Palearctic-) xeromontane species

(= „Mediterranean-Asiatic“ p.p. in most publications, e.g. Boursin 1964, Dufay 1981, Fibiger 1990, etc.). Species marked with* are polytypic species.

1.1. Holo-Mediterranean-xeromontane: *Dichagyris (Dichagyris) candelisequa* ([Denis & Schiffermüller], 1775)*, *D. (D.) renigera* (Hübner, [1808])*, *D. (Yigoga) forcipula* ([Denis & Schiffermüller], 1775)*, *Euxoa (Euxoa) decora* ([Denis & Schiffermüller], 1775)*, *E. (E.) birivia* ([Denis & Schiffermüller], 1775)*, *E. (E.) hastifera* (Donzel, 1847)*, *E. (E.) cos* (Hübner, 1827)*, *Chersotis alpestris* (Boisduval, 1834)*, *Ch. elegans* (Eversmann, 1837)*, *Ch. anatolica* (Draudt, 1936)*, *Ch. larixia* (Guenée, 1852)*, *Ch. fimbriola* (Esper, [1803])*, *Rhyacia (Lafontainea) helvetica* (Boisduval, 1833)*, *Hadena clara* (Staudinger, 1901)*, *H. tephroleuca* (Boisduval, 1833)*, *Apamea platinea* (Treitschke, 1825)* (maps: Hacker & Varga 1990, Varga 1997).

1.2. Ponto-Mediterranean-(Ponto-Mediterranean-Iranian)-xeromontane: *Dichagyris (D.) melanura* (Kollar, 1846)*, *D. (Y.) nigrescens* (Höfner, 1888)*, *D. (Y.) celsicola* (Bellier, 1859)*, *Euxoa (E.) glabella* F. Wagner, 1930*, *Euxoa (Pleonectopoda) derrae* Hacker, 1985, *Chersotis laeta* (Rebel, 1904)*, *Eugnorisma (Metagnorisma) pontica* (Staudinger, 1901)*, *Hadena vulcanica* (Turati, 1907) (= *H. urumovi* Drenovsky, 1931)*, *H. drenowskii* (Rebel, 1930)*, *Calophasia barthae* (F. Wagner, 1929), *Behounekia freyeri* (Frivaldszky, 1835), *Heterophysa dumetorum* (Geyer, 1828)*.

1.3. Cretan-endemic-xeromontane: *Dichagyris rhadamanthys* Reisser, 1958, *Euxoa malickyi* Varga, 1990.

1.4. Cyprus-endemic-xeromontane: *Dichagyris (D.) endemica* Fibiger, Nielsen & Svensen, 1999, *D. (D.) adelphi* Fibiger, Nielsen & Svensen, 1999, *Ammoconia aholai* Fibiger, 1996, *Polymixis (Parabrachionycha) aphrodite* Fibiger, 1997.

1.5. Atlanto-Mediterranean-xeromontane: *Dichagyris (D.) constanti* (Millière, 1860), *D. (D.) serraticornis* (Staudinger, 1897), *Euxoa (E.) mendelis* Fernandez, 1918*, *E. (Pleonectopoda) nevadensis* Corti, 1928, *E. (P.) haverkampfi* (Standfuss, 1893)*, *Cladocerotis optabilis* (Boisduval, 1834), *Eugnorisma (Metagnorisma) arenoflava* (Schawerda, 1934).

1.6. Atlanto-Ponto-Mediterranean-xeromontane: *Euxoa (E.) vitta* (Esper, 1786)*, *Epipsilia cervantes* Reisser, 1935*, *Conisania renati* (Oberthür, 1890)*.

1.7. Mauretanian-xeromontane: *Actebia (Parexarnis) photophila* (Guenée, 1852), *Euxoa (E.) rugifrons* (Mabille, 1888), *E. (E.) oranaria* (Bang-Haas, 1906) (= *E. psimmythiosa* Boursin, 1958), *Chersotis rungsi* Boursin, 1944, *Epipsilia boursini* Rungs, 1972, *Eremohadena roseonitens* (Oberthür, 1887).

1.8. Atlanto-Mediterranean–Mauretanian-xeromontane: *Dichagyris (Stenosomides) mansoura* (Chrétien, 1911)*, *Euxoa (E.) abdallah* (Oberthür, 1918)* (= *E. suffusa* Fernandez, 1918, *E. (E.) ambrosiana* Boursin, 1927), *E. (Pleonectopoda) powelli* (Oberthür, 1912)*, *Calophasia almoravida* (Graslin, 1863), *C. hamifera* Staudinger, 1863.

1.9. Lebanese-xeromontane: *Dichagyris (Yigoga) libanicola* (Corti, 1933), *Pachyagrotis tischendorfi* (Püngeler, 1925).

Legend of Figure 21, page 24. – Fig. 21: Dispersal was canalised by the constraints of passing two main filter-corridors. 1 (dark red) Xeromontane way of dispersal; 2. (dark green) Boreo-Arboreal way of dispersal; 3. Core area of diversity of xeromontane species (*Ctenoceratoda*); 4. Core area of boreo-montane *Polia* (s. str.) species; Rh: “Rhododendron” filter of dispersal; TB: Trans-Beringian way of dispersal; X: Xeromontane filter of dispersal; G: Arrow of dispersal from the Gondwana to the Sino-Himalayan core area.

2. West-and West-Central Asiatic xeromontane species

(„Mediterranean-Asiatic” species *p. p.* in Boursin 1964, Dufay 1981, Fibiger 1990, 1993)

2.1. Anatolian-xeromontane: *Actebia (Protexarnis) squalidiformis* (Corti & Draudt, 1933), *Dichagyris (D.) griseotincta* (F. Wagner, 1931), *D. (D.) catalaipa* Varga, 1993, *Euxoa (E.) luteomixta* F. Wagner, 1930, *E. (E.) anatolica* Draudt, 1936, *E. (E.) robiginosa* (Staudinger, 1894), *Agrotis scruposa* (Draudt, 1934), *Pachyagrotis ankarensis* (Rebel, 1930), *Chersotis ronkayorum* Fibiger, Hacker & Varga, 1992, *Eugnorisma (Phacognorisma) enargiaris* (Draudt, 1934), *Hadena cappadocia* Hacker, 1987, *H. cavalla* Pinker, 1980, *Ostheldera gracilis* (Osthelder, 1933), *Victrix gracilis* (F. Wagner, 1931), *V. hackeri* Varga & Ronkay, 1991 (maps: Varga 1997).

2.2. Anatolian–Transcaucasian-xeromontane: *Dichagyris (D.) armeniaca* Kozhantshikov, 1930*, *D. (D.) romanovi* (Christoph, 1885), *Stenosomides carthalina* (Christoph, 1983), *Euxoa (E.) homicida* (Staudinger, 1900)*, *E. (Pleonectopoda) rjabovi* Kozhantshikov, 1929, *Chersotis stenographa* Varga, 1979, *Ch. gratissima* Corti, 1932, *Ch. friedeli* Pinker, 1974, *Polymixis (Myxinia) csorbagabori* Ronkay, Varga & Hreblay, 1998, *Dasypolia transcaucasica* Ronkay & Varga, 1985, *Luperina diversa* (Staudinger, 1892), *Victrix karsiana* Staudinger, 1879*, *V. artaxias* Varga & Ronkay, 1989 (maps: Varga 1997).

2.3. Transcaucasian–Kurdestanian-xeromontane: *Dichagyris (Yigoga) wiltshirei* (Boursin, 1936), *Xenophysa junctimacula huberi* Varga, 1989, *Eicomorpha kurdestanica* de Freina & Hacker, 1985, *Ammoconia anonyma* Ronkay & Varga, 1984, *Ostheldera arne* Ronkay & Varga, 1994, *Resapamea vaskeni* Varga, 1979*.

2.4. Anatolian–Transcaucasian–N-Iranian-xeromontane: *Actebia (Protexarnis) opisoleuca* (Staudinger, 1881), *Dichagyris (D.) pfeifferi* Corti & Draudt, 1933 (= *D. fredi* Brandt, 1938), *D. (D.) forficula* (Eversmann, 1851)*, *Dichagyris (D.) erubescens* (Staudinger, 1892), *Dichagyris (D.) anastasia* (Draudt, 1936), *D. (Yigoga) weigerti* Hacker & Varga, 1992, *Euxoa (E.) heringi* Christoph, 1877)*, *E. (E.) adjemi* Brandt, 1941, *E. (E.) dsheiron* Brandt, 1938, *E. (E.) conifera* Christoph, 1877* (= *E. difficillima* Draudt), *Euxoa (E.) sulcifera* (Christoph, 1893), *E. (E.) scurrilis* Draudt, 1937, *Chersotis semna* (Püngeler, 1906), *Ch. sarhada* Brandt, 1941, *Chersotis obnubila* (Corti, 1926)*, *Ch. illauta* (Draudt, 1936)*, *Eugnorisma (Metagnorisma) heuristica* Varga & Ronkay, 1987, *Polymixis (P.) rosinae* (Bohatsch, 1908) (= *P. paradiisiaca* Boursin, 1944), *Eremohadena (Megahadena) rjabovi* (Boursin, 1970), *Ostheldera persa* Ronkay & Varga, 1994 (maps: Varga 1997).

2.5. Daghestanian–Transcaucasian-xeromontane: *Dichagyris (D.) achtalensis* (Kozhantshikov, 1929)*, *Euxoa (E.) kuruschensis* Boursin, 1940, *E. (E.) cretaporos* Varga, G. Ronkay & L. Ronkay, 2020, *E. (Pleonectopoda) uncarpa* Kozhantshikov, 1929, *Agrotis fratercula* Pekarsky, Varga, G. Ronkay & L. Ronkay, 2020.

2.6. Iranian–Farsistanian-xeromontane: *Dichagyris (D.) taftana* Brandt, 1941*, *D. (D.) zagoica* Hacker & Ebert, 2002, *D. (D.) gyulaiivani* Gyulai & Varga, *D. (D.) strenua* (Corti, 1926), *Euxoa (E.) charlesboursini* Varga, 2014, *Chersotis firdusii* Schwingenschuss, 1937, *Ch. sterilis* Brandt, 1938, *Ch. eberti* Dufay & Varga, 1995, *Ch. zagoica* Gyulai & Varga, 2006, *Xenophysa cacumena* (Brandt, 1938), *Polymixis (Brandticola) dubiosa* (Brandt, 1938).

2.7. Iranian–Turcomanian-xeromontane: *Dichagyris (D.) psammochroa* Boursin, 1940*, *D. (D.) darius* Boursin, 1940, *D. (D.) humilis* Boursin, 1940*, *Euxoa (E.) subdecora* Hampson, 1903 (= *E. clauda* Püngeler, 1906), *E. (E.) aneucta* Brandt, 1938*, *Rhyacia (Anchorhyacia) gabori* Varga, 1996, *Xenophysa junctimacula* (Christoph, 1887)*, *Ctenoceratoda lupa* (Christoph, 1893), *Polymixis (Boursinixis) pericaspicus* Ronkay, Varga & Hreblay, 1998, *Anchoscelis oropotamica* (Wiltshire, 1941)*.

2.8. Turcomanian-xeromontane: *Actebia (Hemixarnis) iuguma* (Brandt, 1938), *Dichagyris (D.) korshunovi* Varga, 1996, *D. (D.) devota* (Christoph, 1884), *D. (D.) herzi* Kozhantshikov, 1930, *D. (D.) spintheropis* Varga & Ronkay, 1996, *D. (D.) thylacina* Varga, 1996, *D. (D.) exornata* Varga, 1990, *D. (Yigoga) glaucescens* (Christoph, 1887), *Euxoa (E.) sigmata* Kozhantshikov, 1928, *Pachyagrotis benigna* (Corti, 1926), *Chersotis nitens* Brandt, 1941, *Ch. kouros* Varga & Ronkay, 1996, *Ch. binaloudi* Brandt, 1941, *Eugnorisma (Holognorisma) cuneiferum* Varga & Ronkay, 1994, *Ostheldera minna* Ronkay & Varga, 1994, *Polymixis (Myxinia) schistochlora* Ronkay, Varga & Hreblay, 1998.

2.9. Transcaucasian–Turcomanian–Iranian-xeromontane: *Chersotis hahni* (Christoph, 1885), *Ch. sjuntensis* Kusnetzov, 1958, *Polymixis (M.) philippi* (Püngeler, 1911), *P. (M.) achrysa* Ronkay, Varga & Hreblay, 1998, *P. (E.) colluta* (Staudinger, 1888) (= *E. apoteinea* (Brandt, 1938), *P. (Lophotyna) crinomima* (Wiltshire, 1946)*, *Rhiza (Graphantha) gnorima* (Püngeler, 1907)*).

2.10. N-Iranian–Turcomanian–Turkestanian-xeromontane: *Dichagyris (D.) grisescens* Staudinger, 1878*, *D. (D.) leucomelas* Brandt, 1841*, *D. (D.) terminicincta* Corti, 1933*, *D. (D.) squalidior* (Staudinger, 1901), *D. (D.) celebra* (Alphéraky, 1897)*, *D. (D.) elbursica* Draudt, 1937, *D. (D.) singularis* (Staudinger, 1877)*, *D. (D.) eureteocles* (Boursin, 1940)*, *D. (D.) amoena* (Staudinger, 1891)*, *D. (Yigoga) disturbans* (Püngeler, 1914), *D. (Y.) improba* (Staudinger, 1888)*, *Euxoa (E.) transcaspica* Kozhantshikov, 1928 (= *E. catervaria* Corti, 1929, *E. lugubris* Brandt, 1941)*, *E. (E.) perierga* Brandt, 1938*, *E. (E.) mustelina* (Christoph, 1877)*, *E. (E.) cespitis* (Swinhoe, 1885)* (= *E. praestigiosa* Brandt, 1941), *Agrotis bifurca* Staudinger, 1881, *A. psammocharis* Boursin, 1950, *Chersotis curvispina* Boursin, 1961, *Rhyacia (Anchorhyacia) psammia* (Püngeler, 1899)* (= *R. nyctymerides* (Bang-Haas, 1922), *Eugnorisma (Holognorisma) spodia* (Püngeler, 1899)*, *E. (H.) kristensi* Varga, G. Ronkay & L. Ronkay, 2015, *E. (H.) atrabaelbops* Varga, 1975*, *Eugrapha marcida* (Christoph, 1887)*, *Rhiza (G.) laciniosa* (Christoph, 1887)*, *Heptapotamia* (= *Guselderia eustratii* Alphéraky, 1882 (= *G. lutea* Hacker, 1987)*.

2.11. Anatolian–N-Iranian–Tien-Shan–Altai-xeromontane: *Dichagyris (D.) truculenta* (Lederer, 1853)*, *Eugnorisma (H.) eminens* (Lederer, 1855)*, *E. insignata* (Lederer, 1853)*.

3. Central- and Inner-Asiatic xeromontane species

3.1. Turkestanian-(Tienshan)-Mongolian-xeromontane: *Actebia (Parexarnis) poecila* (Staudinger, 1895), *A. (P.) laetifica* (Staudinger, 1889), *A. (P.) candida* (Staudinger, 1888), *A. (Protexarnis) confusa* (Alphéraky, 1882)*, *Dichagyris (D.) verecunda* (Staudinger, 1898)*, *D. clara* (Staudinger, 1888)*, *D. (D.) pudica* (Staudinger, 1895)*, *D. (subg. ?) plumbea* (Alphéraky, 1887), *Xestia (Pachnobia) senescens* (Staudinger, 1881)*, *Ammogrotis suavis* (Staudinger, 1888)*, *Ctenoceratoda zetina* (Staudinger, 1900)*, *Polia lama* (Staudinger, 1886) (= *P. enodata* Bang-Haas, 1912)*, *Auchmis detersina* (Staudinger, 1889), *A. curva* (Staudinger, 1889)*, *Aedophron eos* Varga & Ronkay, 1991.

3.2. W-Mongolian-xeromontane species: *Xenophysa sharhu* Varga, 1989, *Ctenoceratoda sukharevae* (Varga, 1973)*, *C. oxyptera* Varga, 1992, *C. julianae* Varga, 1992, *C. argyrea* Varga, 1992, *C. persephone* Varga, Ronkay & Ronkay, 2017, *C. scotosparsa* Varga, Ronkay & Ronkay, 2017, *C. cyanochrea* Varga, Ronkay & Ronkay, 2017, *Rhiza (Graphantha) calligrapha* (Ronkay & Varga, 1989), *Orohadena clementissima* (Ronkay & Varga, 1993), *Auchmis mongolica* (Staudinger, 1896)*.

3.3. Mongolian–S-Siberian-xeromontane species: *Actebia (Ledereragrotis) difficilis* (Ershov, 1877), *Dichagyris (D.) kaszabi* Varga, 1973, *D. (D.) ignara* (Staudinger, 1896), *D. (Stenosomides) spissilinea* (Staudinger, 1896), *Trichosilia nigrita* (Graeser, 1892) (= *T. maerens* Staudinger, 1896), *T. honesta* (Staudinger, 1892) (= *T. pulchrella* Bang-Haas, 1912), *Euxoa (E.) decorans* (Staudinger, 1896), *E. (E.) fissa* (Staudinger, 1895), *E. (Orosagrotis) tristis* (Staudinger, 1897), *Chersotis transiens* (Staudinger, 1896), *Prognorisma albifurca* (Ershov, 1877), *Pseudohermenassa melancholica* (Lederer, 1857), *P. ononensis* (Bremer, 1861), *Versutographa versuta* (Püngeler, 1901), *Estimata herrichschaefferi* (Alphéraky, 1898), *Ctenoceratoda peregovitsi* Varga & Gyulai, 1999, *Oncocnemis senica* (Eversmann, 1857), *Palaeagrotis inops* (Lederer, 1857).

3.4. W-Tienshan–Alai-xeromontane: *Dichagyris (Yigoga) subturbans* (Boursin, 1948), *Euxoa (E.) enixa* Püngeler, 1906, *E. (E.) bogdanovi* (Ershov, 1873), *Chersotis petermarci* Varga, 1988, *Rhyacia (Dichorhyacia) ignobilis* (Staudinger, 1888), *Eugnorisma (E.) jubilans* Varga, Ronkay & Gyulai, 1995, *Ctenoceratoda tancrei* (Graeser, 1892), *Bryomixis lichenosa* Ronkay & Varga, 1990.

3.5. Turkestanian–E. Tienshan-xeromontane: *Actebia (Parexarnis) violetta* (Staudinger, 1895), *Dichagyris (D.) stellans* Corti & Draudt, 1933, *D. (D.) umbrifera* (Alphéraky, 1882)*, *D. (Albocosta) juldussi* (Alphéraky, 1882), *Euxoa* (subg. ?) *cuprina* (Staudinger, 1899), *Chersotis vicina* (Corti, 1930), *Ch. calorica* (Corti, 1930), *Rhyacia (Anchorhyacia) nyctymerina* (Staudinger, 1888), *R. (A.) diplogramma* (Hampson, 1903), *R. (Ororhyacia) hampsoni* (Bang-Haas, 1910), *Rh. (Standfussrhyacia) junonia* (Staudinger, 1881)*, *Xenophysa agnoscita* Varga, 1989, *Ctenoceratoda khorgossi* (Alphéraky, 1882), *Bryopolia chrysospora* Boursin, 1944.

3.6. Tienshan-(Alai)-Hissaro-Darwaz-xeromontane: *Dichagyris (Albocosta) lasciva* (Staudinger, 1888)*, *Chersotis leucostola* Varga & Ronkay, 1996, *Ch. sordescens* (Staudinger, 1899), *Eugnorisma (Eugnorisma) gaurax* (Püngeler, 1899), *E. (E.) variago* (Staudinger, 1882), *Xestia ornata* (Staudinger, 1891), *Bryoxena tenuicornis* (Alphéraky, 1887).

3.7. Hissaro-Darwaz-(Shugnan)-xeromontane: *Euxoa (E.) hissarica* Varga, 1990, *E. (E.) xanthophylla* Varga, 1990, *Chersotis electrographa* Varga, 1990, *Eugnorisma (E.) deleasma* Boursin, 1970, *Xenophysa paradoxa* Varga, 1989, *Ostheldera kondara* Ronkay & Varga, 1994.

3.8. Tienshan-W-Pamir-xeromontane: *Euxoa (E.) flavisignata* Corti, 1931, *E. (E.) plumbina* (F. Wagner, 1913), *Rhyacia (Stenorhyacia) electra* (Staudinger, 1888)* (= *R. griseoalba* Kozhantshikov, 1937), *Rh. (Anchorhyacia) subdecora* Staudinger, 1887*, *Rh. (A.) similis* Staudinger, 1881, *Xenophysa agnoscita* Varga, 1990, *Ctenoceratoda turpis* (Staudinger, 1900), *Bryopolia chamaeleon* (Alphéraky, 1887).

3.9. W-Pamir-xeromontane: *Dichagyris (D.) apochora* Varga & Gyulai, 2001, *Dichagyris (Yigoga) nekrasovi* Varga & Gyulai, 2001, *Euxoa (E.) melanochroa* Varga, 1990, *E. (E.) psammospora* Varga & Gyulai, 2001, *Chersotis argyllographa* Varga & Gyulai, 2001, *Ch. lukhtanovi* Varga & Gyulai, 2001, *Ch. nekrasovi* Varga, 1998, *Dasypolia shugnana* Varga, 1982, *Polymixis (Boursinixis) pamiridia* Boursin, 1970.

3.10. Transalai-E. Pamir-xeromontane: *Euxoa (Pleonectopoda) murzini* Varga, 1990, *E. (P.) puengeleri* (F. Wagner, 1913), *Ctenoceratoda longicornis* (Graeser, 1892), *C. lukhtanovi* Varga & Gyulai, 1999, *C. psychrogena* Varga & Gyulai, 1999, *C. aksakal* Varga & Gyulai, 1999, *Apamea nekrasovi* Mikkola, Varga & Gyulai, 1995.

3.11. Tienshan-Hissar-Hindukush-xeromontane: *Eugnorisma (E.) trigonica* Alphéraky, 1872*, *Goniographa funkei* Püngeler, 1901, *G. decussa* Staudinger, 1896.

3.12 W-Pamir-Hindukush-xeromontane: *Actebia (Hemiexarnis) berezkii* (Kozhantshikov, 1937), *Dichagyris (D.) chrysopyga* Boursin, 1963, *D. (Yigoga) unifica* (Kozhantshikov, 1937), *Euxoa (E.) submelanochroa* Gyulai & Varga, 2006, *E. (Pleonectopoda) kotzschii* Draudt, 1937 (= *E. plumbescens* Kozhantshikov, 1937).

3.13 (Hissaro-Darwaz)-Pamirs-West-Himalayan-xeromontane: *Actebia (Protexarnis) monogramma* (Hampson, 1903), *Actebia (Hemiexarnis) peperida* (Hampson, 1903), *Dichagyris (D.) leucographa* Varga, 1990, *D. (D.) scotographa* Varga, 1990, *D. (D.) hypotacta* Varga, G. Ronkay & L. Ronkay, 2021, *Euxoa (E.) naumannii* Varga, 1990, *E. (E.) hypochlora* Boursin, 1964*, *Agrotis semivirens* Kozhantshikov, 1937, *Rhyacia (A.) oxytheca* Boursin, 1957, *Rh. (A.) oromys* Varga, 1990, *Xenophysa poecilogramma* Varga, 1985, *Bryopolia monotona* Varga & Ronkay, 1990, *B. tsvetaevi* Varga & Ronkay, 1990*, *Bryoxena plantei* Varga & Ronkay, 1990.

3.14. Hindukush-xeromontane: *Dichagyris (D.) euteles* (Boursin, 1940), *D. (D.) stenoptera* Boursin, 1961, *D. (D.) ammoxanthoides* Varga, 1975, *Euxoa (E.) subeucta* Varga, 2014, *E. (E.) metasigmata* Varga, 2014, *E. (E.) eremorealis* Varga, 1975, *E. (Orosargotis) triumregium* Varga, 1979, *Chersotis pachnosa* Varga, 1975, *Ch. antigrapha* Boursin, 1961, *Ch. metagrapha* Varga, 1975, *Ch. delear* Boursin, 1970, *Rhyacia (A.) evartianae* Varga, 1990, *Rh. (A.) scythropa* Boursin, 1961, *Standfussiana socors* Corti, 1925, *Xenophysa boursini* Varga, 1985, *X. argyrogramma* Varga, 1985, *X. xenogramma* Boursin, 1969, *X. monastica* Boursin, 1969, *Eicomorpha epipsilioides* Boursin, 1969, *Ctenoceratoda septemlacustris* Gaal-Haszler, Lödl, G. Ronkay, L. Ronkay & Varga, *Polymixis (Boursinixis) zophodes* (Boursin, 1960), *P. (B.) polymorpha* (Boursin, 1960), *P. (B.) stictineura* (Boursin, 1960), *P. (Eremophysa) roehrei* Boursin, 1961, *Bryopolia holosericea* Boursin, 1960, *Bryoxena boursini* (Plante, 1983), *B. tribulis* (Plante, 1983), *Victrix chloroxantha* Boursin, 1957, *V. lichenodes* Boursin, 1969, *V. illustris* Varga & Ronkay, 1991.

3.15. Hindukush-West-Himalayan-xeromontane: *Actebia (Hemiexarnis) peperida* (Hampson, 1903), *Dichagyris (D.) payotiorum* Varga, G. Ronkay & L. Ronkay, 2020, *D. (D.) vietteana* Plante, 1979, *D. (Yigoga) acutijuxta* Boursin, 1957, *D. (subg. ?) poliogramma* (Hampson, 1903), *D. (subg. ?) chersotoides* Hacker, 1990, *Euxoa (E.) vartianica* Boursin, 1963, *E. (E.) bactriana* Varga, 2014, *Chersotis fidahusseini* Varga, Gyulai, G. Ronkay & L. Ronkay, 2014, *Eugnorisma (E.) asad* Boursin, 1963*, *E. (E.) conformis* (Swinhoe, 1885), *Anagnorisma eucratides* (Boursin, 1957), *A. glareomima* (Varga & Ronkay, 1991), *Xenophysa poecilogramma* Varga, 1985, *X. naumannii* Varga, 1990, *Polymixis (subg. ?) vartianorum* (Varga, 1979), *P. (Boursinixis) fabiani* Ronkay, Varga & Hreblay, 1998, *P. (Eremophysa) acharis* (Püngeler, 1903), *P. (E.) argyllosa* Boursin, 1970, *Bryopolia virescens* (Hampson, 1894)*.

3.16. West-Himalayan–(Karakoram)-xeromontane: *Dichagyris* (subg. ?) *nivisparsa* (Butler, 1889), *D.* (subg. ?) *draesekei* (Corti, 1928), *Euxoa* (E.) *hyperythra* Boursin, 1964, *E. dimorpha* (Hampson, 1919), *E.* (E.) *amorpha* Boursin, 1964*, *E.* (E.) *vargai* Hacker, 1996, *Standfussiana herbuloti* Plante, 1987*, *Rhyacia* (*Standfussrhyacia*) *chimaera* Hacker & Varga, 1993, *R.* (S.) *peksi* Hacker, 1990, *R.* (S.) *karakoreas* Hacker & Varga, 1990, *R.* (S.) *horroreas* Varga, 2011, *Anagnorisma goniophora* (Varga, Ronkay & Hacker, 1990), *Ctenoceratoda gandhara* Hacker & Varga, 1990, *C. weigerti* Hacker & Varga, 1990, *C. mallopyga* Varga, Gyulai, Ronkay & Ronkay, 2018, *Bryopolia thomasi* Hacker, 1990, *B. ronkayorum* Hacker, 1996, *Polymixis* (E.) *calamistis* (Hampson, 1894).

3.17. Trans-Himalayan–Tibetan-xeromontane species: *Actebia* (*Hemiexarnis*) *moechilla* (Püngeler, 1906), *A.* (H.) *epiphana* (Boursin, 1940), *Dichagyris* (*Albocosta*) *dulcis* (Alphéraky, 1892), *D.* (A.) *musivula* (Staudinger, 1895), *D.* (subg. ?) *subplumbea* (Staudinger, 1895), *D.* (subg. ?) *geochroides* Boursin, *D.* (subg. ?) *astigmata* (Hampson, 1906), *D.* (subg. ?) *vargazoli* Gyulai & Ronkay, 2001, *D.* (subg. ?) *kormos* Gyulai & Ronkay, 2001, *D.* (subg. ?) *minuta* Gyulai & Ronkay, 2001, *Euxoa* (E.) *polytela* Boursin, 1940, *E.* (E.) *hypobscura* Hreblay & Ronkay, 1998, *E.* (E.) *ratna*, Hreblay & Ronkay, 1998, *E.* (E.) *trivibia* Hreblay & Ronkay, 1998, *Euxoa* (E.) *zugmayeri* Boursin, 1948, *Agrotis lamprosericea* Hreblay & Ronkay, 1998, *A. macrobscura* Hreblay & Ronkay, 1998, *Rhyacia* (*Standfussrhyacia*) *oreas* (Püngeler, 1904); *R.* (S.) *mirabilis* Boursin, 1954, *R.* (S.) *admiranda* Gyulai & Ronkay, 2001.

4. Xeromontane (mostly polycentric) species with expansion into the zonal steppe areas (expansive species of different faunal types!)

Dichagyris (*D.*) *nigrolineata* Kozhantshikov, 1930* (syn: *D. squalorum* Auct. nec Eversmann, 1856)*, *D.* (*D.*) *celebrata* (Alphéraky, 1897)*, *Dichagyris* (*D.*) *truculenta* (Lederer, 1853)*, *D.* (*D.*) *lutescens* (Eversmann, 1844), *D.* (*Yigoga*) *flavina* (Herrich-Schäffer, 1852), *D.* (*Yigoga*) *orientis* (Alphéraky, 1882)*, *Euxoa* (E.) *basigramma* (Staudinger, 1870)*, *E.* (E.) *christophi* (Staudinger, 1870), *E.* (E.) *distinguenda* (Lederer, 1857)*, *E.* (E.) *phantoma* Kozhantshikov, 1928, *E.* (E.) *mustelina* (Christoph, 1876)*, *E.* (E.) *deserta* (Staudinger, 1870)*, *E.* (E.) *foeda* (Lederer, 1885), *E.* (E.) *fallax* (Eversmann, 1854), *E.* (E.) *triaena* Kozhantshikov, 1929 (= *E. sagitta* Hübner, 1827), *E.* (E.) *acuminifera* (Eversmann, 1854), *E.* (*Orosagrotis*) *tristis* (Staudinger, 1897), *Chersotis capnistis* (Lederer, 1871)*, *Ch. andereggii* (Boisduval, 1837), *Ch. alpestris* (Boisduval, 1834)*, *Ch. transiens* (Staudinger, 1896), *Eugnorisma* (*Holognorisma*) *puengeleri* Varga & Ronkay, 1987, *E.* (H.) *ignoratum* Varga & Ronkay, 1994, *E.* (H.) *chaldaica* (Boisduval, 1840)*, *Xestia sareptana* (Herrich-Schäffer, 1851) (= *Amathes iobaphes* Boursin, 1940), *Polia serratilinea* (Treitschke, 1825)*, *Apamea leucodon* (Eversmann, 1837)*, *Rhiza* (*Rh.*) *commoda* (Staudinger, 1889), *G. indigna* (Christoph, 1887), *Rhiza* (*Rh.*) *minuta* (Püngeler, 1900), *Eremohadena siri* (Ershov, 1874), *E. adscripta* (Püngeler, 1914), *Oncocnemis confusa* (Freyer, 1842), *O. strioligera* (Lederer, 1853), *O. nigricula* (Eversmann, 1847), etc.

Appendix II

Allopatric and parapatric sister species in Palaearctic Noctuinae

2.1. Species groups in the genus *Dichagyris*

Lederer, 1857 (subg. *Dichagyris*)

Subg. *Dichagyris* (type species *Agrotis melanura* Kollar, 1846)

Dichagyris melanura group

Dichagyris melanura (Kollar, 1846)

Dichagyris melanura melanura (Kollar, 1846)

Dichagyris melanura albida (Caradja, 1931)

Dichagyris melanura dufayi Moberg & Fibiger, 1990

Dichagyris melanura hyrcanica Boursin, 1963

Dichagyris rhadamanthus Reisser, 1958

Dichagyris korshunovi Varga, 1996

Dichagyris leucomelas group

Dichagyris leucomelas Brandt, 1941

Dichagyris leucomelas leucomelas Brandt, 1941

Dichagyris leucomelas endemica Fibiger, Svendson & Nilsson, 1999

Dichagyris leucomelas ladakhensis Hacker & Peks, 1990

Dichagyris grisescens (Staudinger, 1878)

Dichagyris melanuroides Kozhantshikov, 1930 (sensu Boursin et Auctorum!)

Dichagyris stellans (Corti & Draudt, 1933)

Dichagyris duskei Moberg & Fibiger, 1990

Dichagyris imperator group

Dichagyris imperator (Bang-Haas, 1912)

Dichagyris pfeifferi group

Dichagyris pfeifferi (Corti & Draudt, 1933) (= *fredi* Brandt, 1941)

Dichagyris vallesiaca group

Dichagyris vallesiaca Boisduval, 1932

Dichagyris vallesiaca vallesiaca Boisduval, 1932

Dichagyris vallesiaca squalorum (Eversmann, 1856)

Dichagyris vallesiaca subsqualorum Kozhantshikov, 1930

Dichagyris vallesiaca crymaea Kozhantshikov, 1930

Dichagyris vallesiaca inexpectata Kozhantshikov, 1925

Dichagyris vallesiaca venosa Kozhantshikov, 1930

Dichagyris vallesiaca opulenta Brandt, 1941 (species?!)

Dichagyris fuscashmiriana Varga, G. Ronkay & L. Ronkay, 2020 (subspecies?!)

Dichagyris griseotincta Wagner, 1931

Dichagyris nigrolineata group

Dichagyris nigrolineata Kozhantshikov, 1930

Dichagyris nigrolineata nigrolineata Kozhantshikov, 1930 (= *squalorum* sensu Boursin et Auctorum, nec Eversmann, 1856)

Dichagyris nigrolineata rubidior Corti, 1933

Dichagyris tyrannus (Bang-Haas, 1912)

Dichagyris striata Kozhantshikov, 1930

Dichagyris striata striata Kozhantshikov, 1930

Dichagyris striata beluchus Brandt, 1941

Dichagyris striata kalastrata Varga, G. Ronkay & L. Ronkay, 2020

Dichagyris eremicola group

Dichagyris eremicola (Standfuss, 1888)

Dichagyris wilsoni Fibiger, 2002 (subspecies ?!)

Dichagyris squalidior group

Dichagyris squalidior (Staudinger, 1901)

Dichagyris lux Fibiger & Nupponen, 2002 (subspecies of *D. squalidior*?!)

Dichagyris payotiorum Varga, G. Ronkay & L. Ronkay, 2020

Dichagyris taftana group

Dichagyris taftana Brandt, 1941

Dichagyris taftana taftana Brandt, 1941

Dichagyris taftana elborsasta Varga, G. Ronkay & L. Ronkay, 2021

Dichagyris taftana safavida Varga, G. Ronkay & L. Ronkay, 2021

Dichagyris guenterberti Varga, G. Ronkay & L. Ronkay, 2021

Dichagyris kirghisa group

Dichagyris kirghisa (Eversmann, 1856)

Dichagyris terminicincta group

Dichagyris terminicincta (Corti & Draudt, 1933)

Dichagyris terminicincta terminicincta (Corti & Draudt, 1933)

Dichagyris terminicincta maraschi Corti & Draudt, 1933 (= *phaeotaenia* (Boursin, 1940))

Dichagyris terminicincta ssp. from the Kopet-Dagh Mts

Dichagyris terminicincta capnista (Boursin, 1963)

Dichagyris ilseae Stangelmaier, Wieser & Fibiger, 2003

Dichagyris euteles (Boursin, 1963)

Dichagyris psammochroa group

Dichagyris psammochroa Boursin, 1940

Dichagyris psammochroa dichroa Boursin, 1940

Dichagyris psammochroa kopetdaghimena Varga, G. Ronkay & L. Ronkay, 2021

Dichagyris catalaipa Varga, 1993

Dichagyris afghana Boursin, 1963

Dichagyris kurbatskyi Varga, G. Ronkay & L. Ronkay, 2021

Dichagyris apochora Varga & Gyulai 2001

- Dichagyris humilis* group**
- Dichagyris humilis* Boursin, 1940
Dichagyris hypotacta Varga, G. Ronkay & L. Ronkay, 2021
- Dichagyris clara* group**
- Dichagyris clara* Staudinger, 1888
Dichagyris clara clara Staudinger, 1888
Dichagyris clara gobialtaica Varga, 1996
Dichagyris leucographa Varga, 1990
Dichagyris scotographa Varga, 1990
Dichagyris kautti Varga & Ronkay, 1991
- Dichagyris hymalayensis* group**
- Dichagyris hymalayensis* Turati, 1933 (= *despecta* Corti & Draudt, 1933, subsp.?)
Dichagyris chrysopyga (Boursin, 1963)
- Dichagyris celebrata* group**
- Dichagyris armeniaca* Kozhantshikov, 1930
Dichagyris armeniaca armeniaca Kozhantshikov, 1930
Dichagyris armeniaca centranatolica Varga, G. Ronkay & L. Ronkay, 2020
Dichagyris celebrata (Alphéraky, 1897)
Dichagyris celebrata celebrata (Alphéraky, 1897)
Dichagyris celebrata pashtu Varga, 1996
Dichagyris kongur Varga, 1996
- Dichagyris umbrifera* group**
- Dichagyris umbrifera* (Alphéraky, 1882)
Dichagyris umbrifera umbrifera (Alphéraky, 1882)
Dichagyris umbrifera inumbrata Varga, Gyulai & Miatleuski, 2002
Dichagyris kaszabi Varga, 1973
Dichagyris naumannii Varga, 1996
Dichagyris herzi Kozhantshikov, 1930
Dichagyris jacobsoni Kozhantshikov, 1930
Dichagyris boursini Brandt, 1941
Dichagyris achtalensis Kozhantshikov, 1929
Dichagyris korsak Varga, Gyulai & Miatleuski, 2002
- Dichagyris pudica* group**
- Dichagyris pudica* (Staudinger, 1895)
Dichagyris pudica pudica (Staudinger, 1895)
Dichagyris pudica griseola (Staudinger, 1895)
- Dichagyris singularis* group**
- Dichagyris singularis* (Staudinger, 1877)
Dichagyris singularis singularis (Staudinger, 1877)
Dichagyris singularis mesopotamica Hacker & Weigert, 1986
Dichagyris melanofusca Varga, G. Ronkay & L. Ronkay, 2020
- Dichagyris amoena* group**
- Dichagyris amoena* (Staudinger, [1892])
Dichagyris amoena amoena (Staudinger, [1892])
Dichagyris amoena diffluentior Varga, G. Ronkay & L. Ronkay, 2020
- Dichagyris amoena fuscarenosa* Varga, G. Ronkay & L. Ronkay, 2020**
- Dichagyris anastasia* (Draudt, 1936)**
- Dichagyris constanti* group**
- Dichagyris constanti* (Millière, 1860)
Dichagyris constanti constanti (Millière, 1860)
Dichagyris constanti eos (Oberthür, 1913)
- Dichagyris candelisequa* group**
- Dichagyris candelisequa* ([Denis & Schiffermüller], 1775)
Dichagyris candelisequa candelisequa ([Denis & Schiffermüller], 1775)
Dichagyris candelisequa zernyi (Rungs, 1952)
Dichagyris candelisequa cyrnos (Schawerda, 1928)
Dichagyris candelisequa achaemenidica Hacker, 1990
Dichagyris candelisequa rana (Kindermann, 1853)
Dichagyris candelisequa serpens Varga, G. Ronkay & L. Ronkay, 2020
Dichagyris elbursica Draudt, 1937
- Dichagyris verecunda* group**
- Dichagyris verecunda* (Püngeler, 1898)
Dichagyris verecunda verecunda (Püngeler, 1898)
Dichagyris verecunda psammodis Varga, 1993
Dichagyris karakorealis Varga, G. Ronkay & L. Ronkay, 2020
- Dichagyris renigera* group**
- Dichagyris renigera* (Hübner, [1808])
Dichagyris renigera renigera (Hübner, [1808])
Dichagyris renigera funestissima (Bubacek, 1926)
Dichagyris renigera argentina (Caradja, 1930)
Dichagyris renigera funebris (Staudinger, 1892)
Dichagyris renigera subsp. (from Lebanon)
Dichagyris poecilopetala Varga, 1979
- Dichagyris forficula* group**
- Dichagyris forficula* (Eversmann, 1851)
Dichagyris forficula forficula (Eversmann, 1851)
Dichagyris forficula hadjina (Staudinger, 1892)
Dichagyris forficula adefti Fibiger, Svendson & Nilsson, 1999
Dichagyris forficula devota (Christoph, 1884)
Dichagyris turana (Staudinger, 1892)
Dichagyris turana turana (Staudinger, 1892)
Dichagyris turana furiosa (Bang-Haas, 1912)
Dichagyris contermina (Corti, 1930) (= *D. wolfi* Hacker, 1985)
Dichagyris erubescens (Staudinger, 1892)
- Dichagyris argentea* group**
- Dichagyris argentea* Kozhantshikov 1929
Dichagyris darius Boursin, 1940
Dichagyris darius darius Boursin, 1940
Dichagyris darius acroptera Varga, 1996
Dichagyris gyulaiivani Gyulai & Varga, 2002

Dichagyris eureteocles* groupDichagyris eureteocles* Boursin, 1940*Dichagyris eureteocles eureteocles* Boursin, 1940*Dichagyris eureteocles kappadotis* Varga, G. Ronkay & L. Ronkay, 2020*Dichagyris eureteocles karayalchis* Varga, G. Ronkay & L. Ronkay, 2020*Dichagyris danilevskyi* (Shtshetkin, 1965)*Dichagyris zagorica* Hacker & Ebert, 2002*Dichagyris nekrasovi* Varga & Gyulai, 2001***Dichagyris multicuspis* group***Dichagyris multicuspis* (Eversmann, 1852)*Dichagyris aequicuspis* (Staudinger, 1900)***Dichagyris spintheropis* group***Dichagyris spintheropis* Varga & Ronkay, 1996*Chersotis transiens* (Staudinger, 1896)*Chersotis stridula* (Hampson, 1903)*Chersotis stridula stridula* (Hampson, 1903)*Chersotis stridula chimgana* Varga, Gyulai, L. Ronkay & G. Ronkay, 2013*Chersotis cortifera* Rézbányai-Reser, 1997 (species status questioned)***Chersotis capnistis* group***Chersotis capnistis* (Lederer, 1871)*Chersotis capnistis capnistis* (Lederer, 1871)*Chersotis capnistis glabripennis* (Corti, 1926)*Chersotis capnistis schnacki* Fibiger & Moberg, 1993*Chersotis leucostola* Varga & Ronkay, 1996*Chersotis nitens* Brandt, 1941*Chersotis metrapha* Varga, 1975*Chersotis sterilis* (Brandt, 1938)*Chersotis ronkayorum* Fibiger, Hacker & Varga, 1993***Chersotis multangula* group***Chersotis multangula* (Hübner, [1803])*Chersotis andreae* Dufay, 1973*Chersotis semna* (Püngeler, 1906)*Chersotis pachnosa* Varga, 1975***Chersotis griseivena* group***Chersotis griseivena* (Hampson, 1894)*Chersotis harutai* Ronkay & Varga, 1998*Chersotis delear* Boursin, 1970*Chersotis vargai* Hacker, 1992*Chersotis electrographa* Varga, 1990***Chersotis juvenis* group***Chersotis juvenis* (Staudinger, 1901)*Chersotis kouros* Varga & Ronkay, 1996*Chersotis kouros kouros* Varga & Ronkay, 1996*Chersotis kouros zahirshahi* Varga, Gyulai, L. Ronkay & G. Ronkay, 2013*Chersotis shandur* Varga, 1998*Chersotis calorica* (Corti, 1930)***Chersotis vicina* group***Chersotis vicina* (Corti, 1930)*Chersotis petermarci* Varga, 1998***Chersotis binaloudi* group***Chersotis binaloudi* Brandt, 1941*Chersotis antigrapha* Boursin, 1961*Chersotis argyllographa* Varga & Gyulai, 2001***Chersotis deplanata* group***Chersotis deplanata* (Eversmann, 1843)***Chersotis hahni* group***Chersotis hahni* (Christoph, 1885)*Chersotis curvispina* Boursin, 1961*Chersotis cryptographa* Varga & Gyulai, 2002*Chersotis zukowskyi* (Draudt, 1936)***Chersotis acutangula* (Staudinger, 1892)***Chersotis acutangula acutangula* (Staudinger, 1892)*Chersotis acutangula klapperichi* Varga, Gyulai,

L. Ronkay & G. Ronkay, 2013

Chersotis acutangula subtilis Hacker & Peks, 1990***Chersotis juncta* (Grote, 1878)*****Chersotis sordescens* group***Chersotis sordescens* (Staudinger, 1900)*Chersotis herczi* Varga, 1996***Chersotis firdusii* Schwingenschuss, 1937***Chersotis fidahusseini* Varga, Gyulai, L. Ronkay & G. Ronkay, 2013*Chersotis fidahusseini fidahusseini* Varga, Gyulai, L. Ronkay & G. Ronkay, 2013*Chersotis fidahusseini heptalimna* Varga, Gyulai, L. Ronkay & G. Ronkay, 2013***Chersotis ocellina* group***Chersotis ocellina* ([Denis & Schiffermüller], 1775)***Chersotis alpestris* (Boisduval, 1837)***Chersotis alpestris alpestris* (Boisduval, 1837)*Chersotis alpestris ponticola* (Draudt, 1936)*Chersotis alpestris caucasica* Varga, Gyulai, L. Ronkay & G. Ronkay, 2013*Chersotis oreina* Dufay, 1984¹ Coloured letters: allo- (para-)patric sister species

Chersotis obnubila group
Chersotis obnubila (Corti, 1926)

Chersotis sjuntensis group
Chersotis sjuntensis (Kuznetsov, 1958)

Chersotis elegans group
Chersotis elegans (Eversmann, 1837)
Chersotis eberti Dufay & Varga, 1995
Chersotis kacem (Le Cerf, 1933)
Chersotis anatolica (Draudt, 1936)
Chersotis larixia (Guenée, 1852)

Chersotis fimbriola group
Chersotis fimbriola (Esper, [1803]) (several subspecies, see Fig. 16)
Chersotis laeta (Rebel, 1904) (several subspecies, see Fig. 16)

Chersotis cuprea ([Denis & Schiffermüller], 1775)
Chersotis cryptocuprea Varga, Gyulai, L. Ronkay & G. Ronkay, 2013
Chersotis rungsi Boursin, 1944
Chersotis stenographa Varga, 1979
Chersotis gratissima (Corti, 1932)
Chersotis zaghros Gyulai & Varga, 2006
Chersotis nekrasovi Varga, 1996
Chersotis nupponenorum Varga, Gyulai, L. Ronkay & G. Ronkay, 2013
Chersotis friedeli Pinker, 1974

Chersotis illauta group
Chersotis illauta (Draudt, 1936)

Chersotis glebosa group
Chersotis glebosa (Staudinger, 1900)

Chersotis sarhana group
Chersotis sarhana Brandt, 1941
Chersotis hoppei Varga, Gyulai, L. Ronkay & G. Ronkay, 2013
Chersotis lehmanni Varga, Gyulai, L. Ronkay & G. Ronkay, 2013
Chersotis lukhtanovi Varga & Gyulai, 2001

Chersotis margaritacea group
Chersotis margaritacea (de Villers, 1789)
Chersotis orophila Rungs, 1967

Chersotis anachoreta group (subg. *Cyrebia*)
Chersotis anachoreta (Herrich-Schäffer, 1851)
Chersotis soganli Matov, Gyulai, Varga, Ronkay & Ronkay, 2013
Chersotis adili (Koçak, 1987)
Chersotis tshetverikovi Matov, Gyulai, Varga, Ronkay & Ronkay, 2013
Chersotis shaposhnikovi Matov, Gyulai, Varga, Ronkay & Ronkay, 2013

Chersotis luperinoides (Guenée, 1852)
Chersotis romanovi Matov, Gyulai, Varga, Ronkay & Ronkay, 2013

2.3. Species groups in the genus *Rhyacia* Hübner [1821] 1816

Rhyacia subg. *Rhyacia* Hübner [1821] 1816

Rhyacia (subg. *Rhyacia*) *lucipeta* group:

Rhyacia (*Rhyacia*) *lucipeta* ([Denis & Schiffermüller], 1775) – type species of the genus.

Rhyacia (subg. *Rhyacia*) *ledereri* group

Rhyacia (*Rhyacia*) *ledereri* (Erschoff, 1870)
Rhyacia (*Rhyacia*) *quadrangula* (Zetterstedt, 1839)
Rhyacia (*Rhyacia*) *clemens* (Smith, 1890)

Rhyacia (subg. *Rhyacia*) *simulans* group

Rhyacia (*Rhyacia*) *simulans* (Hufnagel, 1766)
Rhyacia (*Rhyacia*) *arenacea* (Hampson, 1907) – probably a complex of closely related species.
Rhyacia (*Rhyacia*) *afghanidia* Boursin, 1968

Rhyacia subg. *Stenorhyacia* Varga 2011

Rhyacia (subg. *Stenorhyacia*) *electra* group

Rhyacia (*Stenorhyacia*) *electra* (Staudinger, 1888)
 (= *R. griseoalba* Kozhantshikov, 1937)
Rhyacia (*Stenorhyacia*) *caradrinoides* (Staudinger, 1896).

Rhyacia subg. *Dichorhyacia* Varga 2011

Rhyacia (*Dichorhyacia*) *ignobilis* Staudinger, 1888
Rhyacia (*Dichorhyacia*) *fabiani* Varga, 1996

Rhyacia subg. *Ororhyacia* Varga 2011

Rhyacia (*Ororhyacia*) *hampsoni* (Bang-Haas, 1910)

Rhyacia subg. *Anchorhyacia* Varga 2011

Rhyacia (*Anchorhyacia*) *psammia* group
Rhyacia (*Anchorhyacia*) *psammia* (Püngeler, 1906)
 = *nyctymerides* (Bang-Haas, 1922)
Rhyacia (*Anchorhyacia*) *psammia psammia* (Püngeler, 1906)
Rhyacia (*Anchorhyacia*) *psammia stavroitiacus* Touleshkoff, 1951
Rhyacia (*Anchorhyacia*) *psammia alagesica* Boursin, 1962
Rhyacia (*Anchorhyacia*) *psammia rehnensis* (F. Wagner, 1937)

- Rhyacia (Anchorhyacia) psammia roseoflava* (Corti, 1933)
- Rhyacia (Anchorhyacia) psammia nyctymerides* (Bang-Haas, 1922)
- Rhyacia (Anchorhyacia) latebrosa* Gyulai, 2021 (Pakistan, Hindukush) (syn. or ssp.??)
- Rhyacia (Anchorhyacia) nyctymerina* (Staudinger, 1888)
- Rhyacia (Anchorhyacia) gabori* Varga, 1996
- Rhyacia (Anchorhyacia) evartiana* Varga, 1990
- Rhyacia (Anchorhyacia) oxytheca* Boursin, 1957
- Rhyacia (Anchorhyacia)* sp. from Iran, Khorasan
- Rhyacia (Anchorhyacia) subdecora* (Staudinger, 1887)
- Rhyacia (Anchorhyacia) scythropa* Boursin, 1961
- Rhyacia (Anchorhyacia) diplogramma* group**
- Rhyacia (Anchorhyacia) diplogramma* (Hampson, 1903)
- Rhyacia (Anchorhyacia) oromys* Varga, 1990
- Rhyacia (Anchorhyacia) similis* group:**
- Rhyacia (Anchorhyacia) similis* (Staudinger, 1888)
(= *decorata* Staudinger, 1881, praecocc.)
- Rhyacia* subg. *Standfussrhyacia* Hacker & Varga, 1990**
- Rhyacia (Standfussrhyacia) chimaera* group:**
- Rhyacia (Standfussrhyacia) chimaera* (Hacker & Varga, 1990)
- Rhyacia (Standfussrhyacia) mirabilis* group:**
- Rhyacia (Standfussrhyacia) mirabilis* Boursin, 1954
- Rhyacia (Standfussrhyacia) mirabilis mirabilis* Boursin, 1954
- Rhyacia (Standfussrhyacia) mirabilis nepalensis* Boursin, 1954
- Rhyacia (Standfussrhyacia) admiranda* Gyulai & Ronkay, 2001
- Rhyacia (Standfussrhyacia) miranda* Gyulai, 2021
- Rhyacia (Standfussrhyacia) junonia* group:**
- Rhyacia (Standfussrhyacia) junonia* (Staudinger, 1881)
- Rhyacia (Standfussrhyacia) junonia junonia* (Staudinger, 1881)
- Rhyacia (Standfussrhyacia) junonia alaina* (Staudinger, 1888)
- Rhyacia (Standfussrhyacia) junonia alexandrina* Corti & Draudt, 1933
- Rhyacia (Standfussrhyacia) junonia calamochroa* Varga, 1973
- Rhyacia (Standfussrhyacia) schistochroa* Varga, 1973
- Rhyacia (Standfussrhyacia) kulinuna* Gyulai, 2021
- Rhyacia (Standfussrhyacia) illustris* Hacker & Kautt, 1996
- Rhyacia (Standfussrhyacia) horroreas* Varga, 2011
- Rhyacia (Standfussrhyacia) oreas* (Püngeler, 1904)
- Rhyacia (Standfussrhyacia) unicornis* Varga 2011
- Rhyacia (Standfussrhyacia) karakoreas* Hacker & Varga, 1990
- Rhyacia (Standfussrhyacia) peksi* Hacker, 1990

2.4. Species groups in the genus *Eugnorisma*

Boursin, 1846

Eugnorisma, subg. *Holognorisma* Varga, Gyulai, Ronkay & Ronkay, 2014

- Eugnorisma (Holognorisma) chaldaica* group**
- Eugnorisma (Holognorisma) chaldaica* (Boisduval, 1840)
- Eugnorisma (Holognorisma) chaldaica chaldaica* (Boisduval, 1840)
- Eugnorisma (Holognorisma) chaldaica kurdistana* Hacker, 1986
- Eugnorisma (Holognorisma) chaldaica rubicunda* Varga & Ronkay, 1990
- Eugnorisma (Holognorisma) kristensi* Varga, Gyulai, Ronkay & Ronkay, 2014
- Eugnorisma (Holognorisma) spodia* (Püngeler, 1900)
- Eugnorisma (Holognorisma) spodia spodia* (Püngeler, 1900)
- Eugnorisma (Holognorisma) spodia psammochrea* Varga & Ronkay, 1987

- Eugnorisma (Holognorisma) ignoratum* group**
- Eugnorisma (Holognorisma) ignoratum* Varga & Ronkay, 1994
- Eugnorisma (Holognorisma) cuneiferum* Varga & Ronkay, 1994
- Eugnorisma (Holognorisma) puengeleri* Varga & Ronkay, 1987
- Eugnorisma (Holognorisma) mikkolai* Varga, Gyulai, Ronkay & Ronkay, 2014
- Eugnorisma (Holognorisma) tamerlana* (Hampson, 1903)
- Eugnorisma (Holognorisma) eminens* group**
- Eugnorisma (Holognorisma) eminens* (Lederer, 1855)
- Eugnorisma (Holognorisma) eminens eminens* (Lederer, 1855)
- Eugnorisma (Holognorisma) eminens clarior* Varga, 1975
- Eugnorisma (Holognorisma) atrabaelbops* Varga, 1975
- Eugnorisma (Holognorisma) atrabaelbops atrabaelbops* Varga, 1975
- Eugnorisma (Holognorisma) atrabaelbops firyuza* Varga, Gyulai, Ronkay & Ronkay, 2014
- Eugnorisma (Holognorisma) atrabaelbops scotophalia* Varga, Gyulai, Ronkay & Ronkay, 2014

Eugnorisma, subg. *Eugnorisma* Boursin, 1946

- Eugnorisma (Eugnorisma) trigonica* group**
- Eugnorisma (Eugnorisma) trigonica* (Alphéraky, 1882)
- Eugnorisma (Eugnorisma) trigonica trigonica* (Alphéraky, 1882)
- Eugnorisma (Eugnorisma) trigonica gauracoides* Hacker & Peks, 1990

- Eugnorisma (Eugnorisma) gaurax* (Püngeler, 1900)
 (= *funebris* Varga & Ronkay, 1990)
- Eugnorisma (Eugnorisma) deleasma* Boursin, 1967
 Eugnorisma (Eugnorisma) deleasma deleasma Boursin,
 1967
- Eugnorisma (Eugnorisma) deleasma reducta* Boursin,
 1968
- Eugnorisma (Eugnorisma) deleasma hissarica* Varga &
 Ronkay, 1987
- Eugnorisma (Eugnorisma) jubilans* Varga, Ronkay &
 Gyulai, 1995
- Eugnorisma (Eugnorisma) insignata* group**
- Eugnorisma (Eugnorisma) insignata* (Lederer, 1853) –
 Polytypic species, subspp. not clarified
- Eugnorisma (Eugnorisma) conformis* (Swinhoe, 1885)
- Eugnorisma (Eugnorisma) rafidain* (Boursin, 1936)
 (= *semiramis* (Boursin, 1940))
- Eugnorisma (Eugnorisma) asad* Boursin, 1963
 Eugnorisma (Eugnorisma) asad asad Boursin, 1963
- Eugnorisma (Eugnorisma) asad plantei* Varga, Ronkay
 & Hacker, 1990
- Eugnorisma (Eugnorisma) asad eva* Varga, Gyulai,
 Ronkay & Ronkay, 2014
- Eugnorisma (Eugnorisma) variago* (Staudinger, 1882)
 Eugnorisma (Eugnorisma) variago variago (Staudinger,
 1882)
- Eugnorisma (Eugnorisma) variago xanthiago* Varga &
 Ronkay, 1987
- Subgenus *Metagnorisma* Varga & Ronkay, 1987**
- Eugnorisma (Metagnorisma) depuncta* group**
- Eugnorisma (Metagnorisma) depuncta* (Linnaeus, 1761)
 Eugnorisma (Metagnorisma) depuncta depuncta
 (Linnaeus, 1761)
- Eugnorisma (Metagnorisma) depuncta transcaucasica*
 Varga, Gyulai, Ronkay & Ronkay, 2014
- Eugnorisma (Metagnorisma) arenoflavida* (Schawerda,
 1934)
- Eugnorisma (Metagnorisma) janhabeli* Varga, Gyulai,
 Ronkay & Ronkay, 2014
- Eugnorisma (Metagnorisma) pontica* group**
- Eugnorisma (Metagnorisma) pontica* (Staudinger, 1891)
 Eugnorisma (Metagnorisma) pontica pontica (Stau-
 dinger, 1891)
- Eugnorisma (Metagnorisma) pontica anis* Varga &
 Ronkay, 1987
- Eugnorisma (Metagnorisma) pontica zaghros* Varga &
 Ronkay, 1987
- Eugnorisma (Metagnorisma) deserta* Varga & Ronkay,
 1987
- Eugnorisma (Metagnorisma) heuristica* Varga & Ronkay,
 1987
- Eugnorisma (Metagnorisma) hermannhackeri* Varga,
 Gyulai, Ronkay & Ronkay, 2014
- 2.5. Species groups in the genus *Goniographa***
 Ronkay & Varga, 2002
- Goniographa decussa* group**
- Goniographa decussa* (Staudinger, 1897)
- Goniographa discussa* Varga & Ronkay, 2002
- Goniographa shchetkini* Varga & Ronkay, 2002
- Goniographa funkei* group**
- Goniographa funkei* (Püngeler, 1901)
- Goniographa metafunkei* Varga & Ronkay, 2002
- Goniographa naumanni* Varga & Ronkay, 2002
- Goniographa marcida* group**
- Goniographa marcida* (Christoph, 1893)
- Goniographa gyulaipeteri* Varga & Ronkay, 2002