



IMPACT OF FUNGAL METABOLITES ON SEED MYCOFLORA AND SEED GERMINATION OF HERBAL PLANTS OF *THESPIESIA POPULNEA*

Harane Smita S¹ Harpale Dattatraya V²

¹SPH Arts, Science and Commerce Mahila Mahavidyalaya, Nashik-423105

²HPT Arts & RYK Science College, Nashik- 05

Corresponding Author- Harane Smita S

Email- smitaharane@gmail.com

DOI-

Abstract

Effects of fungal metabolites on seed mycoflora, seed germination of medicinal plants of Thespesia populnea seeds were studied. The metabolites affect the seed germination as well as the length of radicle. Production of the metabolites depends on the species, isolates or strain of the fungus, and on the ecological conditions like temperature humidity etc. and the nature of the substrate. Therefore, experiments were performed to study the effect of extracellular metabolites, produced by eight the dominant storage fungi, which are represented in the research paper. Seeds of herbal plant, like those of agricultural and horticultural crops, carry a wide variety of micro-organisms like fungi, bacteria and even some viruses. The metabolites affect the seed germination as well as the length of radicle. Production of the metabolites depends on the species, isolates or strain of the fungus, and on the ecological conditions like temperature humidity etc. and the nature of the substrate.

Key: Metabolites, seed mycoflora, medicinal seeds.

Introduction

Fungi are known to produce various extracellular metabolites; they may be enzymes, toxins, organic acids etc. All these play a role in disease development. Fungi growing on stored seeds produce highly toxic metabolites (Mycotoxins), that are poisonous, sometimes fatal to man and animals. These fungi are also known to produce cell wall degrading enzymes like pectolytic and cellulolytic enzymes.

Some of them are also known to produce the enzyme amylase which attack the starch content of the seed and deteriorate them and the enzyme lipase that attack on the lipids present in the cell walls. Seeds are used for the medicine. These seeds are found to be frequently contaminated by fungi (Roy *et al.* 1988, Mamatha *et al.*, 2009).⁽¹⁾ Chaurasia (1990) investigated that almost all medicinal seed samples were associated with a large number of fungi. Effects of fungal metabolites on seed mycoflora, seed

germination of herbal plants of *Thespesia populnea* seeds were studied.⁽²⁾

Material and Method

The toxicity of the culture filtrates of fungi on seed germination and seedlings was assessed by the method adopted by Papdiwal and Deshpande (1978).⁽³⁾ The toxicity was tested by using as follows: The toxic effect of culture filtrates on seed germination was studied by keeping 12 seeds on filter paper in a petridish. The filter paper was soaked in 8 ml of the above solution. Filter paper soaked in 8 ml tapwater served as control. The filter paper was kept moist by adding the cultures filtrate or water in the respective petridishes. The plates were incubated for 80 hours and percentage germination was recorded.

Observations

Seeds of herbal plant *viz.* *Thespesia populnea*, was surface sterilized with 0.1 % HgCl₂ solution for four minutes and washed repeatedly with sterile distilled water. For the study of fungal metabolites,

8 dominant fungi, occurring on the herbal seeds under investigation, was selected. They were *Penicillium corylophilum*, *Pythium indigoferae*, *Aspergillus carbonarius*, *A. flavus*, *A. niger*, *Cladosporium cladosporioides*, *Fusarium oxysporum*, and *Rhizopus oryzae*. The extracellular metabolites of these fungi were collected, as per the method described earlier.

Effect on Seed Germination

Suitable controls were maintained with sterile distilled water. The seed germination of *Thespesia populnea* herbal

plant under investigation was noted, and expressed as percent germination. The length of radicle of every germinated seed was recorded, and its mean was calculated. The results were compared with control and percent inhibition of seed germination, and average radicle length were calculated. The results are presented in table 1, fig. 1 and 2. The effect of these metabolites of respective fungi on seed germination was studied as per the method described earlier (Papdiwal and Deshpande, 1978).⁽⁴⁾

Table 1: Effect of culture filtrates on seed germination and length of radicle of *Thespesia populnea*

SN	Metabolites from the fungus	Seed germination		Average Radicle length	
		% germination	% inhibition	Length in cm	% inhibition
1	<i>Aspergillus carbonarius</i>	50	45.45	0.6	40.00
2	<i>A. flavus</i>	40	55.56	0.4	60.00
3	<i>A. niger</i>	50	45.45	0.6	40.00
4	<i>Cladosporium cladosporioides</i>	60	33.34	0.6	40.00
5	<i>Fusarium oxysporum</i>	55	38.89	0.7	30.00
6	<i>Penicillium corylophilum</i>	40	55.56	0.4	60.00
7	<i>Pythium indigoferae</i>	65	27.78	0.7	30.00
8	<i>Rhizopus oryzae</i>	60	33.34	0.6	40.00
9	Control	90	---	1.0	--
Mean			56.67		0.62
S. D.			15.21		0.18
C. V.			26.84		28.72

Source: Fieldwork, 2022




Fig. 1




Fig. 2

The extracellular metabolites secreted by the 8 storage fungi, under investigation, were inhibitory for the seed germination and radicle length, of *Thespesia populnea*. It is observed from table 1, fig. 1 and 2 that in case of *Thespesia populnea*, the highest inhibition (55.56%) of seed germination was with culture filtrates of *A. flavus* and *P. corylophilum*. The culture filtrates of *A. flavus* and *P. corylophilum* were found to cause maximum inhibition (60%) of the length of the radicle. The culture filtrate of *P. indigoferae* caused minimum inhibition of seed germination (27.78%) while the culture filtrate of *F. oxysporum* and *P. indigoferae* were minimum inhibitory for the radicle growth (30%). It is further noted that the metabolites from *P. indigoferae* were comparatively less inhibitory for seed

germination and radicle length of *T. populnea*; compared to other fungi under investigation (Plate 1).

Result

The result reveals that the culture filtrate of *Aspergillus flavus* and *P. corylophilum* cause maximum inhibition, of seed germination, among the 8 storage fungi studied. The culture filtrate of the fungus was found to cause maximum inhibition of seed germination of the medicinal plants investigated. The maximum inhibition of seed germination was observed in the culture filtrates of *A. flavus* and *P. corylophilum* (inhibition 60%). Minimum inhibition was observed in culture filtrates of *Pythium indigoferae* (27.78%). The adverse effect of culture filtrates was also observed on radicle length.



PLATE :- 4.9 Fig. 1) *Thespesia populnea* tree 2) and 3) Inflorescence 4) Fruits 5) Seeds

Plate.1: *Thespesia Populnea* Tree

Harane Smita S Harpale Dattatraya V

Reference

1. Afzal. R., Mughal, S.M. Munir, M., Sultana K, Qureshi,R., Arshad M. and M. K. Laghari., (2010) Mycoflora associated with seeds of different sunflower cultivars and it's management. Pak. J. Bot. 42(1) 435-445.
2. Agrawal, R. H. (1995) Seed Technology. Oxford and IBH Publishing Co., New Delhi.
3. Bhanumathi A and V. Ravishankarra(2008) Seed mycoflora of some important forest tree species. Seed Res. 36(1):95-98.
4. Bharjan S. K., V. (1995) Studies on antifungal activity of some medicinal plant product on Jowar, grains during storage. Proc. 82nd Science Congress, Part III.
5. Chourasia, H.K. and A. K. Roy (1991) Effect of Temperature, relative humidity and light on aflatoxin-B production in Neem and Datura seeds, Inst. J. Pharmacognosy 29 (3):197-202.
6. Doyer, I. C. (1938) Manual for the Determination of Seed borne diseases. ISTA, Washington, pp. 59.
7. Papdiwal, P. B. and K. B. Deshpande (1978) Studies on *in vitro* production of toxic metabolites by *Xanthonas malvacearum* and their phytotoxic effects, *Ind. J. Bot.* 1:1-4.