

OPEN ACCESS

***Correspondence:** Duwa, R S.
Department of Biological
Sciences, Nigeria Police
Academy, Wudil, Kano,
Nigeria.

Email: rsnb8100@gmail.com

Specialty Section:

This article was submitted to
Biology, a section of NAPAS.

Accepted: 1 May, 2022

Published: 1 May 2022

Citation:

Duwa, R S and Ononamadu, C
J. (2022). *Furcocercus*
Cercariae and Other Parasitic
Infections in Freshwater Snails
of Jakara Dam, Kano State,
Nigeria. Nig Annals of Pure &
Appl Sci. 5(1):171-177.
DOI:10.5281/zenodo.6969552

Publisher:

cPrint, Nig. Ltd

E-mail:

cprintpublisher@gmail.com

Access Code

<http://napas.org.ng>

Furcocercus Cercariae and other Parasitic Infections in Freshwater Snails of Jakara Dam, Kano State, Nigeria

Duwa, RS¹, Ononamadu, CJ².

¹ Department of Biological Sciences and ² Department of Biochemistry, Nigeria Police Academy, Wudil, Kano, Nigeria.

ABSTRACT

Freshwater Snails from different parts of Jakara dam, Kano state, Nigeria were examined for the presence of cercariae. The locations were Wasai, Bangare and Yadakunya in three local government areas. A total of 613 snails comprising of the genus *Bulinus*, *Lymnea*, *Bithynia* were collected out of which 104 (17%) were infected with different types of cercariae, nematode larva and annelids (*Oligochaetes*). There was no significant association between the locations sampled and the prevalence of infected snails. A prevalence of 17.24% of infected snails was observed in Bangare; this was not significantly different from the 19.20% observed in Wasai (PR: 0.89, χ^2 :0.11, P = 0.7370) and 16.14% observed in Yadakunya (PR:1.06, χ^2 :0.04, P = 0.8322). The prevalence observed in Wasai was also not significantly different from that of Yadakunya (PR:1.19, χ^2 :0.73, P = 0.3915). *Lymnea* and *Bulinus* species were the only snails infected with cercariae while *Bithynia* species were infected with nematodes larva. Five types of *Furcocercus* cercariae were recovered from the *Bulinus* snails only during the cold harmattan season while *Fasciola* cercariae and other cercariae were recovered at the end of the harmattan season.

Keywords: *Cercaria*, *Furcocercus*, *Fasciola*, *Jakara dam*, *Snail intermediate hosts*

INTRODUCTION

Freshwater snails play a very important role in the transmission of trematode infections. The larval forms of these trematodes known as cercariae gets into the body of the host either by ingestion with vegetables or penetration of the skin when the host comes in contact with the water body. Some *Furcocercus* cercariae are the infective forms of trematodes of the genus *Schistosoma* which causes Schistosomiasis. Schistosomiasis is a neglected tropical disease. In Africa alone, it is estimated that some 200 million people are infected with this blood fluke. Nigeria is one of most affected country with an estimated 101.28 million people at risk of infection

MATERIALS AND METHODS

This study was carried out in Wasai, Yadakunya and Bangare parts of Jakara dam between February and December 2018. Jakara dam was constructed in 1976 and this part of it is situated in Minjibir Local Government Area (LGA) in the North Eastern part of Kano metropolis about 41.5Km from the city centre. Bangare part of Jakara dam is the most polluted.



Snail collection

Three parts of Jakara dam were investigated for the presence of freshwater snails. These parts are located in three local government areas of the state (Gezawa, Minjibir and Ungoggo). Snails were collected by hand picking only (due to accessibility) into a wide mouthed plastic container and brought to the laboratory. It was then sorted out into different genera and species for further investigation. The snails were fed with lettuce during the study.

Cercarial shedding

Snails were separately placed in compartmented petridishes and exposed to artificial light (rechargeable lamp) for 2-3 hours to stimulate cercarial shedding, they were also exposed to sunlight on a sunny day. Few drops of water were then taken from the petridish containing the snail and placed on a glass slide using a pipette and narcotized with neutral red to demobilize them before examination with a compound microscope for more detailed identification. Those snails that do not shed cercaria on the first day were re-exposed on the second day and those that shed cercariae were continuously exposed and observed under the microscope until the water in which they were placed was exhausted. Photomicrograph of the cercariae were taken using Toupview 3.1 digital camera fitted to the microscope through a USB and cercariae were identified using keys provided by Frandsen and Christensen (1984) and Key to cercarial identification.

Statistical analysis

The statistical analysis was performed using Stata version 12.1 for Windows (Stata Corp LP, United States). Chi-square test and classical test of hypothesis by *csi* (Tables for Epidemiologist: cohort study risk-ratio calculator) command in Stata were used to test for association and differences between prevalence of snail infection and sampled location. This program implements Pearson Chi square as it determines prevalence,

prevalence (Risk) difference (PD), prevalence (Risk) ratio (PR), Odd Ratio and confidence intervals.

RESULTS

The prevalence of infection in *Lymnea* and *Bulinus* species in the different parts of Jakara dam was shown in Table 1 (16.96%). A total of 613 snails were collected out of which 104 were infected with different types of cercariae. The highest prevalence was found in Wasai (19.2%) (Minjibir LGA) part of the dam followed by Bangare (17.24%, Gezawa LGA) and the least in Yadakunya (Ungoggo LGA) 16.1% although sample sizes were not equal. The infection with furcocercus cercariae was only observed in *Bulinus* species in the month of December in Yadakunya. The *Fasciola* cercaria was observed in *Lymnea* species. The month with the highest infection rate is December. There was no snail in the rainy season during the months of July and September. One of the furcocercus cercaria named *Cercaria bulini yadakunya* I was the most abundant, the tail furca curved backwards at intervals during the swimming period until the body separates from the tail while still under the pressure of a cover slip. They have no eyespots.




Fig.2a: Furcocercuscercariae (a-e) from Yadakunya part of Jakara dam they were named *Cercariae bulini yadakunya I V*




Fig.2b: Furcocercus cercariae (a-e) from Yadakunya part of Jakara dam they were named *Cercariae bulini yadakunya I V*.




Fig.3: a. *Echinostome* cercaria b. *Fasciola* cercaria. c. *Gymnocephalus* cercariae



Fig.4: Other infections:a,b. larva of a Nematode from *Bithynia tentaculata* c. Annelid (Oligochaeta) from a *Lymnea* snail






Fig.5: *Bulinus* species

**Fig. 6: Lymnea species****Fig.7: Bithynia specie****Statistical analysis**

There was no significant association between the locations sampled and the prevalence of infected snails. A prevalence of 17.24% of infected snails was observed in Bangare; this was not significantly different from the 19.20% observed in Wasai (PR: 0.89, χ^2 :0.11, $P = 0.7370$) and 16.14% observed in Yadakunya (PR:1.06, χ^2 :0.04, $P = 0.8322$). The prevalence observed in Wasai was also not significantly different from that of Yadakunya (PR:1.19, χ^2 :0.73, $P = 0.3915$).

DISCUSSION

The cercarial infections observed in snails in the different parts of this water body shows that there is likely to be different types of trematode infections taking place. Considering the fact that the water body is accessible to different animals, other animal schistosome cercaria can be harboured by the different snails. Okeke and Ubachukwu (2016) reported the infection of Biomphalaria snails with different morphotypes of cercaria in south-east Nigeria. Although many

Table 1: Prevalence of infection in the three locations

Location	Snails examined	Snails infected	Prevalence (%)
Bangare	58	10	17.24
Wasai	140	27	19.2
Yadakunya	415	67	16.1
Total	613	104	16.96

cercariae have been discovered in the different parts of Jakara dam, this is the first time different types of *Furcercus* cercariae are encountered despite continuous reports of Schistosomiasis in the area (Duwa *et al*, 2009; Abubakar, *et al*, 2015; Dawaki, *et al*, 2016). The *Bulinus* species were shedding only *furcercus* cercaria. There was no *Biomphalaria* snail during this study but the predominant cercariae in this study named *Cercaria bulini yadakunya* I (after the location where the snail intermediate host was collected) has the characteristics of a *Schistosoma mansoni* cercaria and has the tail furcae curved backwardly in the resting position when observed under pressure of the coverslip on the slide. According to Frandsen and Christensen (1984) the backward curvature of the tail furca is rarely seen in preserved cercariae or in living cercariae under pressure of the coverslip on the slide. Multiple infections were only observed in *Lymnea* snails. According to Webster *et al* (2013) there is the likelihood of novel zoonotic hybrid schistosomes evolving with subsequent changes in the parasites' life history traits, transmission potential and virulence when humans and their livestock start to frequent the same water bodies. In view of this there is the need for the proper identification of snails and cercariae obtained to ascertain the particular infection being transmitted as there may be animal forms of schistosomiasis considering the morphology of the *furcercus* cercariae emitted and for proper control strategy of snails, fascioliasis and schistosomiasis in the area.

REFERENCES

- Abubakar S., Oyeyi T. O., Yunusa I. and Ahmed M. K (2015). Co-infection of urinary schistosomiasis and haematuria *Annals of Experimental Biology*, 3 (1):29-32.
- Aisha Salihu Jibril (2016). Studies on Rice Fields Snails and Urinary Schistosomiasis Among Rice Farmers in Kura Local Government Area, Kano State, Nigeria. MSc Thesis.
- Dawaki, S., H M. Al-Mekhlafi, Init, I., J, Ibrahim, Awatif., *Et al* (2016). The Menace of Schistosomiasis in Nigeria: Knowledge, Attitude, and Practices Regarding Schistosomiasis among Rural Communities in Kano State *PLOS*.
- Duwa, M.R., Oyeyi, T.I and Bassey, S.E (2009). Prevalence and Intensity of Urinary Schistosomiasis Among Primary School Pupils in Minjibir L G A of Kano state *Bayero Journal of Pure and Applied Sciences*, 2(1):75-78.
- Frandsen, F. and Christensen, N.Ø (1984). An introductory guide to the identification of cercariae from African freshwater snails with special reference to cercariae of trematode species of medical and veterinary importance: *Acta Tropica* 4L 181-202.
- Okeke, O.C. and Ubachukwu, P.O. (2017). Trematode infections of the freshwater snail *Biomphalaria pfeifferi* from a south-east Nigerian community with emphasis on cercariae of *Schistosoma*. *Journal of Helminthology* 91, 295301.
- Yousif, F., Ayoub, M., Tadros, M., El Bardicy, S. and Abolarinwa, S. (2014) Description of two new cercariae (an echinostome cercaria and xiphidiocercaria) procured from *Biomphalaria pfeifferi* (Krauss) from Nigeria. *Journal of the Egyptian Society of Parasitology* 44, 373380.
- Webster BL, Emery AM, Webster JP, Gouvras A, Garba A, Diaw O, *et al* (2012). Genetic diversity within *Schistosoma haematobium*: DNA barcoding reveals two distinct groups. *PLoS Neglected Tropical Diseases*.;6:e1882.