



Article

Heterostigmatic mites (Acari: Trombidiformes: Prostigmata) associated with Coleoptera and Hymenoptera in Mazandaran province, northern Iran

Hamidreza Hajiqanbar* and Elham Arjomandi

Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, 14115-336, Tehran, Iran; E-mails: hajiqanbar@modares.ac.ir, elham.arjomandi@modares.ac.ir

* Corresponding author

ABSTRACT

In a study on insect-associated heterostigmatic mites (Acari: Trombidiformes: Prostigmata) carried out in Mazandaran province, northern Iran, during spring and summer 2016–2017, a total of 13 species from 11 genera of six families were identified. Among them, three species *Imparipes sebastianovi* Khaustov, 2008, *Scutacarus hystrichocentrus* Sevastianov, 1983 (Scutacaridae), and *Petalonium kurosai* Khaustov, 2014 (Neopygmephoridae) are new records for mite fauna of Iran. The *S. hystrichocentrus* is also new for Asian mite fauna. Eight new host records are documented and the distributions of the mites are revised. Beetles of the family Cerambycidae are recorded as host of the mite families Microdispidae and Scutacaridae for the first time.

KEY WORDS: Ant; bee; beetle; new record; new host record.

PAPER INFO.: Received: 24 August 2019, Accepted: 26 September 2019, Published: 15 October 2019

INTRODUCTION

Association with arthropods, mostly insects, have been evolved in many Heterostigmata (Acari: Trombidiformes: Prostigmata) (Kaliszewski *et al.* 1995; Mortazavi *et al.* 2018). For some fungivorous families, this association is restricted to phoresy including Dolichocybidae, most Microdispidae, Neopygmephoridae, Pygmephoridae and Scutacaridae (Kaliszewski *et al.* 1995; Walter *et al.* 2009). Females of the family Trochometridiidae appear to be kleptoparasites of their bee associates (Cross and Bohart 1969, 1978) however, in recent years, they have been found on some other insects as well (Hajiqanbar *et al.* 2009; Mortazavi *et al.* 2011; Loghmani *et al.* 2014a; Jacinavicius *et al.* 2018; Khaustov and Frolov 2018).

During a survey on heterostigmatic mites associated with insects in Mazandaran province, northern Iran, in 2016 and 2017, a total of 13 species from 11 genera of six families were identified. Among them, three species *Imparipes sebastianovi* Khaustov, 2008, *Scutacarus hystrichocentrus* Sevastianov, 1983 (Scutacaridae), and *Petalonium kurosai* Khaustov, 2014 (Neopygmephoridae) are new records for mite fauna of Iran. The *S. hystrichocentrus* is also new for Asia. Eight new host records are documented and the distributions of the mites are revised.

How to cite: Hajiqanbar, H. & Arjomandi, E. (2019) Heterostigmatic mites (Acari: Trombidiformes: Prostigmata) associated with Coleoptera and Hymenoptera in Mazandaran province, northern Iran. *Persian Journal of Acarology*, 8(4): 343–352.

MATERIALS AND METHODS

The study was carried out from 2016 to 2017 in Mazandaran province, northern Iran. The host insects were collected directly in their habitats, by sweeping or light traps. Mite specimens were removed using an Olympus stereomicroscope, cleared in a mixture of lactophenol and Nesbitt's solution and mounted in Hoyer's medium. The morphology of mites was studied using a light microscope (model BX51, Olympus, Tokyo, Japan) equipped with phase contrast illumination. The systematics of Heterostigmata follows that of Khaustov (2008). All materials are deposited in the Acarological Collection, Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.

Superfamily Dolichocyboidea Mahunka, 1970
Family Dolichocybidae Mahunka, 1970
Genus *Pavania* Lombardini, 1949

***Pavania kamalii* Hajiqanbar & Khaustov, 2010**

Material examined – 7 ♀, Mazandaran province, Noor, forest park, from under elytra of the beetle *Scarabaeus* sp. (Col.: Scarabaeidae), 4 August 2016, collected from livestock dung.

World distribution – Iran, North Khorasan province, phoretic on *Scarabaeus* spp. (Hajiqanbar and Khaustov 2010); Mazandaran province (current study).

Superfamily Trochometridoidea Mahunka, 1970
Family Trochometridiidae Mahunka, 1970
Genus *Trochometridium* Cross, 1965

***Trochometridium kazachstanicum* Khaustov & Eidelberg, 2002**

Material examined – 5 ♀, Mazandaran province, Miankaleh, 15 August 2016, from under elytra of *Copris lunaris* (L.) (Col.: Scarabaeidae) attracted to light trap.

World distribution – Kazakhstan, associated with a carabid beetle *Machozetus* sp. (Khaustov and Eidelberg 2002); Iran, Razavi Khorasan province, phoretic on ventral body surfaces of *Gymnopleurus* (s. str.) *mopsus persianus* Reitter (Coleoptera: Scarabaeidae) (Hajiqanbar *et al.* 2009); Mazandaran province (current study).

Remarks – The *Copris lunaris* is a new host record for this mite.

***Trochometridium kermanicum* Mortazavi & Hajiqanbar, 2011**

Material examined – 3 ♀, Mazandaran province, Miankaleh peninsula, 15 August 2016, from under elytra of *Coccobius schreberi* (L.) (Col.: Scarabaeidae), collected from cows dung.

World distribution – Iran, Kerman province, associated with *Paulusiella* sp. (Coleoptera: Elateridae) (Mortazavi *et al.* 2011); Razavi Khorasan province, associated with *Komarowia tartara* (Saussure) (Hymenoptera: Tiphidae) (Loghmani *et al.* 2014a); Golestan province, associated with *Labidura riparia* (Pallas) (Dermaptera: Labiduridae) (Rahiminejad *et al.* 2014); Mazandaran province (current study).

Remarks – The *Coccobius schreberi* is a new host record for this mite.

Superfamily Pygmephoroidae Cross, 1965
Family Microdispidae Cross, 1965
Genus *Paramicrodispus* Khaustov, 2009

***Paramicrodispus crenulatus* (Savulkina, 1978)**

Material examined – 5♀, Mazandaran province, Noor Forest, 17 May 2016, under elytra of *Gnorimus subcostatus* (Mentries) (Col.: Scarabaeidae), collected from rotten logs; 4♀, Mazandaran Province, vicinity of Behshahr, 17 July 2016, under elytra of *Parandra caspia* Ménétriès, 1832 (Col.: Cerambycidae).

World distribution – Bulgaria, in the nest of a small mammal (Savulkina 1978); Crimea, beneath elytra of a carabid beetle *Pterostichus niger* (Schaller) and in a rotten log of *Fagus orientalis* (Khaustov 2009). Iran, Gorgan province, phoretic on *Oryctes nasicornis* L. (Col.: Scarabaeidae) (Hajiqanbar *et al.* 2012); Mazandaran province (current study).

Remarks – This is the first report of the relationship between mites of family Microdispidae and beetles of the family Cerambycidae. Also, *Gnorimus subcostatus* is new host record for the mite *Paramicrodispus crenulatus*.

Family Neopygmephoridae Cross, 1965
Genus *Petalomium* Cross, 1965

***Petalomium kurosai* Khaustov, 2014**

Material examined – 3♀, Mazandaran province, Neka Forest, 14 August 2016, associated with *Lasius turcicus* Santschi (Hymenoptera: Formicidae).

World distribution – Russia (Western Siberia), associated with *Lasius niger* (L.) and *Tetramorium caespitum* (L.) (Khaustov 2014); Iran (current study).

Remarks – This species is a new record for mite fauna of Iran. *Lasius turcicus* is also a new host record for this mite species.

Family Pygmephoridae Cross, 1965
Genus *Elattoma* Mahunka, 1969

***Elattoma cerambycidum* Rahiminejad & Hajiqanbar, 2011**

Material examined – 7♀, Mazandaran province, Neka forest, 4 July 2016, from under elytra of *Morimus verecundus* (Falderman) (Coleoptera: Cerambycidae).

World distribution – Iran, Golestan province, phoretic on *Morimus verecundus* (Coleoptera: Cerambycidae) (Rahiminejad *et al.* 2011); Mazandaran province (current study).

Genus *Metapygmeophorellus* Rahiminejad, Hajiqanbar & Khaustov, 2015

***Metapygmeophorellus colydius* Rahiminejad & Hajiqanbar, 2015**

Material examined – 1♀, Mazandaran province, Noor forest, 12 June 2016, from under elytra of

Parandra caspia Ménétriès (Coleoptera: Cerambycidae).

World distribution – Iran, Golestan province, phoretic on *Colydium elongatum* (F.) (Coleoptera: Zopheridae) and *Dorcus parallelipipedus* (L.) (Coleoptera: Lucanidae) (Rahiminejad *et al.* 2015); Mazandaran province (current study).

Remarks – Cerambycid beetle *Parandra caspia* is a new phoretic host for mites of the genus *Metapygmephorellus*.

Genus *Propygmeophorus* Cross, 1974

***Propygmeophorus crossi* Katlav & Hajiqanbar, 2016**

Material examined – 2♀, Mazandaran province, Sisangan forest, 19 May 2016, associated with *Lucanus ibericus* Motschulsky (Coleoptera: Lucanidae).

World distribution – Iran, Mazandaran province, phoretic on carabid beetles *Pseudoophonus rufipes* (De Geer) and *Pterostichus caspius* (Menetries) (Coleoptera: Carabidae) (Katlav *et al.* 2016); and current study.

Remarks – Lucanid beetle *Lucanus ibericus* is a new phoretic host for mites of the genus *Propygmeophorus*.

Genus *Pseudopygmephorellus* Khaustov, 2008

***Pseudopygmephorellus mazandaranicus* Katlav & Hajiqanbar, 2015**

Material examined – 7♀, Mazandaran province, Mazichal elevations, 14 June 2016, from under elytra of *Onthophagus* sp. (Coleoptera: Scarabaeidae), collected from cow's dung.

World distribution – Iran, Mazandaran province, phoretic on scarabaeid beetles *Onthophagus* sp., *Aphodius depressus* (Kugelann) and *A. varians* Dufschmid (Coleoptera: Scarabaeidae) (Katlav *et al.* 2015); and current study.

Family Scutacaridae Oudmans, 1916

Genus *Archidispus* Karafiat, 1959

***Archidispus minor* Karafiat, 1959**

Material examined – 2♀, Mazandaran province, Miankaleh, 6 August 2016, from under elytra of *Harpalus* sp. (Coleoptera: Carabidae), attracted to light trap.

World distribution – Austria, Germany, Greece, Hungary, India, Poland, Ukraine and Iran (Razavi Khorasan province) associated with several beetles of the family Carabidae including *Acupalpus* spp., *Agonum* spp., *Amara* spp., *Anisodactylus binotatus*, *Calathus* spp., *Harpalus* spp., *Poecilus* spp., *Pterosticus* spp., *Pseudoophonus calceatus* Duft, *Stenolophus* spp. (Ebermann 1991; Khaustov 2008; Loghmani *et al.* 2014b); Mazandaran province (current study).

Genus *Imparipes* Berlese, 1903

Imparipes sebastianovi Khaustov, 2008

Material examined – 2♀, Mazandaran province, Behshahr, Abbas-Abad forest, 16 August 2016, associated with *Lasius flavoniger* Seifert (Hymenoptera: Formicidae).

World distribution – Ukraine, associated with *Lasius fuliginosus* Latreille (Khaustov 2008); Russia (Western Siberia) associated with *L. fuliginosus* (Khaustov 2016); Iran (current study).

Remarks – This is the first record of *I. sebastianovi* from Iran. Moreover, *Lasius flavoniger* is a new host record for *I. sebastianovi*.

Genus Scutacarus Gros, 1845

Scutacarus acarorum (Goeze, 1780)

Material examined – 2♀, Mazandaran province, Behshahr, 19 July 2017, phoretic on *Bombus* sp. (Hymenoptera: Apidae).

World distribution – Holarctic, associated with various bumblebees or their nests (Khaustov 2008) and sometimes as hyperphoresy on deutonymphs of *Parasitellus* mites (Mesostigmata: Parasitidae) (Schousboe 1986). Iran, Ardabil province, associated with *Bombus argillaceus* (Scopoli) (Kazemi and Kamali 2006); Mazandaran province (current study).

Scutacarus hystrichocentrus Sevastianov, 1983

Material examined – 1♀, Mazandaran province, Behshahr, Abbas-Abad forest, 16 August 2016, associated with *Parandra caspia* (Coleoptera: Cerambycidae).

World distribution – Ukraine, associated with *Lasius niger* (L.) and *L. alienus* (Foerster) (Khaustov 2008); Iran (current study).

Remarks – This is a new record for mite fauna of Asia. Moreover, *Parandra caspia* is a new host record for mites of the family Scutacaridae.

DISCUSSION

This study revealed one of representatives of the beetle family Cerambycidae, *Parandra caspia*, hosting three species of pygmephoroids belonging to three different families. They are *Paramicrodispus crenulatus* (Microdispidae), *Metapygmephorellus colydius* (Pygmephoridae) and *Scutacarus hystrichocentrus* (Scutacaridae). *Parandra caspia* is distributed in Azerbaijan (Europe) and Iran (Asia) with a three-year life cycle and is polyphagous in deciduous trees. This beetle usually attacks two-year old dead host plants which are first attacked by other wood boring insects. Therefore, such habitat can probably provide a suitable niche for their progeny and also some fungivorous mites phoretic on adult beetles.

Previously, mites of the families Microdispidae and Scutacaridae were never found to be associated with beetles of the family Cerambycidae [see Hajiqanbar and Hosseininaveh (2014) for microdispids; and Khaustov (2008) and Baumann (2018) for scutacarids]. In Pygmephoridae, only some members of the genus *Elattoma* have been recorded from cerambycids (see Rahiminejad *et al.* 2011, 2016). *Paramicrodispus crenulatus* and *Metapygmephorellus* spp. have been earlier recovered from xylophagous or subcortical beetles (Hajiqanbar *et al.* 2012; Rahiminejad *et al.* 2016), although

S. hystrichocentrus has been collected from some ants (Khaustov 2008). Rotten woods and coarse woody debris mostly contain a large diversity of saprophytic fungi in forest ecosystems (like northern forests of Iran) and discovery of different fungivorous mites on any xylophagous, subcortical and saprophytic insects is quite possible.

Eleven genera of different families of Heterostigmata have been found in this study. In Table 1, numbers of included species of these genera in the world and Iran have been compared. It can be concluded that scutacarid species-rich genera (*Archidispus*, *Imparipes* and *Scutacarus*) need more attention and are to be further collected in various parts of Iran.

Table 1. Comparison of species number of some heterostigmatic genera in the world and Iran.

Genus name	No. of world species	Reference	No. of Iranian species	Reference
<i>Pavania</i>	27	Hajiqanbar <i>et al.</i> (2019)	15	Sobhi <i>et al.</i> (2017b); Hajiqanbar <i>et al.</i> (2019)
<i>Trochometridium</i>	6	Loghmani <i>et al.</i> (2014a)	5	Loghmani <i>et al.</i> (2014a)
<i>Paramicrodispus</i>	3	Hajiqanbar <i>et al.</i> (2012)	2	Hajiqanbar and Sobhi (2018)
<i>Petalomium</i>	50	Silva <i>et al.</i> (2019)	10	Abbasi-Moqadam <i>et al.</i> (2018); Navabi <i>et al.</i> (2018); this study
<i>Elattoma</i>	10	Rahiminejad <i>et al.</i> (2011)	3	Rahiminejad <i>et al.</i> (2011, 2014)
<i>Metapygmephorellus</i>	3	Rahiminejad <i>et al.</i> (2015)	1	Rahiminejad <i>et al.</i> (2015)
<i>Propygmechorus</i>	2	Katlav <i>et al.</i> (2016)	1	Katlav <i>et al.</i> (2016)
<i>Pseudopygmephorellus</i>	8	Khaustov and Frolov (2019)	2	Katlav <i>et al.</i> (2015); Sobhi <i>et al.</i> (2017b)
<i>Archidispus</i>	>70	Khaustov (2008); Kurosa (2009); Katlav <i>et al.</i> (2016); Baumann (2018)	7	Sobhi <i>et al.</i> (2017a)
<i>Imparipes</i>	>160	Ebermann (1998); Khaustov (2008); Baumann and Ferrugut (2019)	17	Sobhi <i>et al.</i> (2017a); this study
<i>Scutacarus</i>	>400	Khaustov (2008); Khaustov and Minor (2018)	18	Sobhi <i>et al.</i> (2017a); this study

ACKNOWLEDGEMENTS

We are grateful to the following entomologists for help to identify different host insects: A. Anichtchenko (Latvia), A. Frolov (Russia), A. Radchenko (Ukraine), and B. Seifert (Germany).

REFERENCES

- Abbasi-Moqadam, F., Hajiqanbar, H. & Mehrabadi, M. (2018) Contribution to the knowledge of the genus *Petalomium* (Acari: Neopygmephoridae) associated with ants from Iran. *Systematic and Applied Acarology*, 23(6): 1180–1189.
<http://doi.org/10.11158/saa.23.6.13>
- Baumann, J. (2018) Tiny mites on a great journey – a review on scutacarid mites as phoronts and inquilines (Heterostigmatina, Pygmephoridae, Scutacaridae). *Acarologia*, 58(1): 192–251.

- <http://doi.org/10.24349/acarologia/20184238>
- Baumann, J. & Ferragut, F. (2019) Description and observations on morphology and biology of *Imparipes clementis* sp. nov., a new termite associated scutacarid mite species (Acari, Heterostigmata: Scutacaridae; Insecta, Isoptera: Rhinotermitidae). *Systematic and Applied Acarology*, 24(2): 303–323.
<https://doi.org/10.11158/saa.24.2.12>
- Cross, E.A. & Bohart, G.E. (1969) Phoretic behavior of four species of alkali bee mites as influenced by season and host sex. *Journal of the Kansas Entomological Society*, 42: 195–219.
- Cross, E.A. & Bohart, G.E. (1978) Some observations of the habits and distribution of *Trochometridium* Cross, 1965 (Acarina: Pyemotidae). *Acarologia*, 20: 286–293.
- Ebermann, E. (1991) Das Phänomen Polymorphismus in der Milbenfamilie Scutacaridae (Acari, Heterostigmata, Tarsonemina, Scutacaridae). *Zoologica*, 141: 1–76.
- Ebermann, E. (1998) *Imparipes (Sporichneuthes)* nov. subgen., a remarkable new taxon in the family Scutacaridae (Acari, Heterostigmata). In: Ebermann E. (Ed.), *Arthropod biology: Contributions to morphology, ecology and systematics*. Biosystematics and Ecology Series, Österreichische Akademie der Wissenschaften, Vienna, 14, pp. 179–214.
- Hajiqanbar, H. & Hosseiniaveh, F. (2014) A new genus and species of the family Microdispidae (Acari: Prostigmata) associated with *Oryctes nasicornis* (Coleoptera: Scarabaeidae) and redescription of the monotypic genus *Vietodispus* Mahunka, 1975. *Zoological Studies*, 53, 58.
<https://doi.org/10.1186/s40555-014-0058-7>
- Hajiqanbar, H. & Khaustov, A. (2010) A new species group and five new species of the genus *Pavania* (Acari: Dolichocybidae) associated with insects, with notes on leg chaetotaxy and the distribution of genera. *European Journal of Entomology*, 107: 441–453.
<https://doi.org/10.14411/eje.2010.051>
- Hajiqanbar, H., Khaustov, A., Kamali, K., Saboori, A. & Kamali, H. (2009) New taxa of the family Trochometridiidae (Acari: Heterostigmata) associated with insects from Iran. *Journal of Natural History*, 43(43–44): 2701–2722.
<http://doi.org/10.1080/00222930903243962>
- Hajiqanbar, H., Khaustov, A. & Mortazavi, A. (2019) New species of the genera *Dolichocybe* and *Pavania* (Acari: Dolichocybidae), phoretic on scarab beetles (Coleoptera: Scarabaeidae) from Iran. *Systematic and Applied Acarology*, 24(8): 1363–1382.
<http://doi.org/10.11158/saa.24.8.3>
- Hajiqanbar, H., Rahiminejad, V. & Fathipour, Y. (2012) New insect host records for mites of the family Microdispidae (Acari: Heterostigmata), with description of a new species of the genus *Paramicrodispus*. *Entomological Science*, 15: 309–313.
<http://doi.org/10.1111/j.1479-8298.2012.00515.x>
- Hajiqanbar, H. & Sobhi, M. (2018) New records of the microdispid mites (Acari: Heterostigmata: Microdispidae) associated with ants with a review of the family in Iran. *Persian Journal of Acarology*, 7(2): 105–113.
<https://doi.org/10.22073/pja.v7i2.35131>
- Jacinavicius, F.C., Silva, R.A., Bassini-Silva, R., Lopez, V.M., Silvestre, R. & Khaustov, A.A. (2018) A new species of *Petalonium* Cross, 1965 (Heterostigmata: Neopygmephoridae) and records of *Trochometridium tribulatum* Cross, 1965 (Heterostigmata: Trochometridiidae) from Brazil. *Acarina*, 26(2): 167–174.
<http://doi.org/10.21684/0132-8077-2018-26-2-167-174>
- Kaliszewski, M., Athias-Binche, F. & Lindquist E.E. (1995) Parasitism and parasitoidism in Tarsonemina (Acari: Heterostigmata) and evolutionary considerations. *Advances in Parasitology*, 35: 335–367.
[https://doi.org/10.1016/S0065-308X\(08\)60074-3](https://doi.org/10.1016/S0065-308X(08)60074-3)

- Katlav, A., Hajiqanbar, H. & Talebi, A.A. (2015) *Pseudopygmephorellus mazandaranicus* sp. nov. (Acari: Heterostigmata: Pygmephoridae), phoretic on scarabaeid dung beetles (Coleoptera: Scarabaeidae) from Iran. *Zootaxa*, 3919(1): 100–110.
<https://doi.org/10.11646/zootaxa.3919.1.5>
- Katlav, A., Hajiqanbar, H. & Talebi, A.A. (2016) Two remarkable new species of the superfamily Pygmephorooidea (Acari: Heterostigmata) associated with beetles (Coleoptera: Carabidae, Staphylinidae). *Annals of the Entomological Society of America*, 109: 136–144.
<http://doi.org/10.1093/aesa/sav099>
- Kazemi, Sh. & Kamali, K. (2006) New records of mites (Astigmata, Mesostigmata, Prostigmata) associated with *Bombus argillaceus* (Hymenoptera, Apoidea) from Iran. In: *Abstract Book of the 12th International Congress of Acarology, 21–26 August 2006, Amsterdam, The Netherlands*, p. 90.
- Khaustov, A.A. (2008) *Mites of the family Scutacaridae of Eastern Palaearctic*. Akademperiodyka, Kiev, 291 pp.
- Khaustov, A.A. (2009) A new genus and three new species of the mite family Microdispidae (Acari: Heterostigmata) from Crimea. *Acarina*, 17: 65–73.
- Khaustov, A.A. (2014) A new species of the genus *Petalonium* (Acari: Heterostigmatina: Neopygmephoridae) from Western Siberia with redescription of *Petalonium nataliae* (Sebastianov, 1967). *Acarina*, 22: 100–108.
- Khaustov, A.A. (2016) Myrmecophilous pygmephoroid mites (Acari: Pygmephorooidea) associated with *Lasius fuliginosus* (Hymenoptera: Formicidae) in Western Siberia, Russia. *International Journal of Acarology*, 42: 92–105.
- Khaustov, A.A. & Eidelberg, M.M. (2002) A new species mites of the genus *Trochometridium* (Acarina: Heterostigmata: Trochometridiidae) from Kazakhstan. *Acarina*, 10: 43–45.
- Khaustov, A.A. & Frolov, A.V. (2018) A new species, new genus and new records of heterostigmatic mites (Acari: Heterostigmata) phoretic on scarab beetles of the subfamily Orphninae (Coleoptera: Scarabaeidae). *Zootaxa*, 4514(2): 181–201.
<http://doi.org/10.11646/zootaxa.4514.2.3>
- Khaustov, A.A. & Frolov, A.V. (2019) Two new species of *Pseudopygmephorellus* (Acari: Heterostigmata: Pygmephoridae) phoretic on beetles of the genus *Rhyparus* (Coleoptera: Scarabaeidae: Aphodiinae) from Papua New Guinea and Indonesia. *Systematic and Applied Acarology*, 24(8): 1541–1554.
<http://doi.org/10.11158/saa.24.8.14>
- Khaustov, A.A. & Minor, M.A. (2018) New species and a new record of the mite family Scutacaridae (Acari: Heterostigmata) from alpine New Zealand. *Zootaxa*, 4508(4): 507–550.
<http://doi.org/10.11646/zootaxa.4508.4.2>
- Kurosa, K. (2009) The scutacarid mites of Japan. XIV. A new species of *Archidispus* phoretic on stenolophine carabid beetles. *Journal of Acarological Society of Japan*, 18 (2): 85–90.
<http://doi.org/10.2300/acari.18.85>
- Loghmani, A., Hajiqanbar, H. & Talebi, A.A. (2014a) An illustrated key to world species of the mite family Trochometridiidae (Acari: Prostigmata), with description of a new species and new insect host records. *Canadian Entomologist*, 146(5): 471–480.
<http://doi.org/10.4039/tce.2014.3>
- Loghmani, A., Hajiqanbar, H. & Talebi, A.A. (2014b) New records of mites of the superfamily Pygmephorooidea (Acari: Heterostigmatina) associated with insects from northeastern Iran and new host records. *Systematic and Applied Acarology*, 19(2): 154–159.
<https://doi.org/10.11158/saa.19.2.5>
- Mortazavi, A., Hajiqanbar, H. & Lindquist, E.E. (2018) A new family of mites (Acari: Prostigmata: Raphignathina), highly specialized subelytral parasites of dytiscid water beetles (Coleoptera: Dytiscidae: Dytiscinae). *Zoological Journal of the Linnean Society*, 184(3): 695–749.

- <http://doi.org/10.1093/zoolinnean/zlx113>
- Mortazavi, A., Hajiqanbar, H. & Saboori, A. (2011) A new species of the family Trochometridiidae (Acari: Heterostigmatina) associated with *Paulusiella* sp. (Coleoptera: Elateridae) from Iran. *Zootaxa*, 2746: 57–68.
- Navabi, A., Hajiqanbar, H. & Mortazavi, A. (2018) New records of the genus *Allopygmephorus* (Acari: Neopygmephoridae) with a list of the species of the family Neopygmephoridae in Iran. *Persian Journal of Acarology*, 7(4): 315–322.
<https://doi.org/10.22073/pja.v7i4.40314>
- Rahiminejad, V., Hajiqanbar, H. & Fathipour, Y. (2011) Two new species of the genus *Elattoma* (Acari: Heterostigmatina: Pygmephoridae) phoretic on *Morimus verecundus* (Coleoptera: Cerambycidae) from Iran. *Zootaxa*, 2903: 48–56.
- Rahiminejad, V., Hajiqanbar, H. & Talebi, A.A. (2014) First record of the phoretic relationship between the mite *Trochometridium kermanicum* (Acari: Heterostigmata) and the earwig *Labidura riparia* (Insecta: Dermaptera). In: Aminian, H. (Ed.), *Proceedings of the 21th Iranian Plant Protection Congress, 23–26 August 2014, Urmia University, Urmia, Iran*, p. 955.
- Rahiminejad, V., Hajiqanbar, H., Khaustov, A.A. & Talebi, A.A. (2015) A new genus and two new species of the family Pygmephoridae (Acari: Heterostigmata) associated with beetles (Insecta: Coleoptera). *Annals of the Entomological Society of America*, 108(5): 893–901.
<http://dx.doi.org/10.1093/ae/aes073>
- Rahiminejad, V., Hajiqanbar, H. & Talebi, A.A. (2016) A new genus and species of the family Pygmephoridae (Acari: Heterostigmata) associated with *Carpelimus rivularis* (Coleoptera: Staphylinidae). *Systematic and Applied Acarology*, 21(4): 461–470.
<http://doi.org/10.11158/saa.21.4.7>
- Savulkina, M.M. (1978) Neue Pygmephiden-Arten (Trombidiformes, Pygmephoridae) aus Nagernestern von Bulgarien und der Sowjetunion. *Parasitologica Hungarica*, 11: 127–140.
- Schousboe, C. (1986) On the biology of *Scutacarus acarorum* Goeze (Acari: Trombidiformes). *Acarologia*, 27(2): 151–158.
- Silva, R.A.D., Khaustov, A.A., Delabie, J.H.C. & Oliveira, A.R. (2019) A new species of *Scutacarus* and description of larva and male of *Petalonium megasolenidiatum* Silva, Khaustov & Oliveira (Acari: Heterostigmatina: Neopygmephoridae, Scutacaridae) from Brazil. *Systematic and Applied Acarology*, 24(8): 1343–1362.
<http://doi.org/10.11158/saa.24.8.2>
- Sobhi, M., Hajiqanbar, H. & Mortazavi, A. (2017a) A contribution to the knowledge of scutacarid mites (Acari: Pygmeporoidea: Scutacaridae) associated with Coleoptera and Hymenoptera (Arthropoda: Insecta) from northwestern Iran. *Acarologia*, 57(4): 1103–1111.
<https://doi.org/10.24349/acarologia.20174222>
- Sobhi, M., Hajiqanbar, H. & Mortazavi, A. (2017b) New species and records of heterostigmatic mites (Acari: Prostigmata: Heterostigmata) phoretic on scarabaeid dung beetles (Coleoptera: Scarabaeidae) from northwestern Iran. *Zootaxa*, 4276: 427–434.
<https://doi.org/10.11646/zootaxa.4276.3.7>
- Walter, D.E., Lindquist, E.E., Smith, I.M., Cook, D.R., Krantz, G.W. (2009) Order Trombidiformes. In: Krantz, G.W. & Walter, D.E. (Eds.), *A manual of acarology*. 3rd ed., Texas, USA: Texas Tech Universiy Perss, pp. 233–420.

COPYRIGHT

Hajiqanbar and Arjomandi. Persian Journal of Acarology is under a free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

کنه‌های هترواستیگما (Acari: Trombidiformes: Prostigmata) مرتبط با سخت‌بالپوشان و بال‌غشائیان در استان مازندران، شمال ایران

حمیدرضا حاجی‌قبر* و الهام ارجمندی

گروه حشر‌شناسی، دانشکده کشاورزی، دانشگاه تربیت مدرس، ۱۴۱۱۵-۳۳۶، تهران، ایران؛ رایانامه‌ها:
hajiqanbar@modares.ac.ir
elham.arjomandi@modares.ac.ir

*نویسنده مسئول

چکیده

در مطالعه‌ای که در مورد کنه‌های هترواستیگما مرتبط با حشرات در استان مازندران، شمال ایران، طی سال‌های ۱۳۹۵ و ۱۳۹۶ انجام گرفت در مجموع ۱۳ گونه متعلق به ۱۱ جنس از شش خانواده شناسایی شد. در بین آنها سه گونه *Imparipes sebastianovi* Khaustov, 2008 *Petalomium kurosai* Khaustov, 2014 (Scutacaridae) و *Scutacarus hystrichocentrus* Sevastianov, 1983 (Neopygmephoridae) گزارش‌های جدیدی برای فون کنه‌های ایران هستند. گونه *S. hystrichocentrus* نیز برای فون کنه‌های آسیا جدید است. هشت گزارش میزانی جدید مورد استناد قرار گرفته و مناطق انتشار کنه‌ها بازبینی شده است. سوسک‌های خانواده Cerambycidae برای نخستین بار به عنوان میزان کنه‌های خانواده Microdispidae و Scutacaridae در جهان گزارش می‌شوند.

واژگان کلیدی: مورچه؛ زنبور؛ سوسک؛ گزارش جدید؛ گزارش میزانی جدید.

اطلاعات مقاله: تاریخ دریافت: ۱۳۹۸/۶/۲، تاریخ پذیرش: ۱۳۹۸/۷/۴، تاریخ چاپ: ۱۳۹۸/۷/۲۳