

**Physiological characteristics of putative Enterobacteriaceae associated with meat and fish available in Southern Brazilian retail markets: antimicrobial susceptibility, toxic metal tolerance and expression of efflux pumps**

Renata da Costa Barros Silva, Jéssica Andrade, Vanessa Cordeiro Dias, Jéssica Sequeto, Nayara Felga Santos, Vânia Lúcia da Silva, Cláudio Galuppo Diniz

Laboratory of Microbial Physiology and Molecular Genetics, Center for Studies in Microbiology, Department of Parasitology, Microbiology and Immunology, Federal University of Juiz de Fora, 36036-330 Juiz de Fora, Brazil

Corresponding author:

Claudio Galuppo Diniz, PhD

Email: [claudio.diniz@ufjf.br](mailto:claudio.diniz@ufjf.br)

Laboratory of Microbial Physiology and Molecular Genetics, Center for Studies in Microbiology, Department of Parasitology, Microbiology and Immunology, Federal University of Juiz de Fora, 36036-330 Juiz de Fora, Brazil

## Supplementary Information

Table S1 – Antimicrobial resistance phenotypes observed among mesophilic facultatively anaerobic Gram-negative rods presumptively characterized as enterobacteria isolated from meat and fish samples in Southern Brazilian retail markets.

AMR phenotype	Number of bacteria isolates (n) and frequency (%)				
	Chicken (n= 91)	Beef (n= 91)	Pork (n= 112)	Fish (n= 136)	Total (n= 436)
CFZ	12 (13.2)	25 (27.5)	19 (17.0)	30 (22.1)	86 (20.0)
AMP	11 (12.1)	8 (8.8)	9 (8.0)	7 (5.1)	34 (7.9)
CFO	-	4 (4.4)	4 (3.6)	1 (0.7)	9 (2.0)
TET	-	1 (1.1)	3 (2.7)	1 (0.7)	5 (1.1)
SUT	-	-	2 (1.8)	1 (0.7)	3 (0.7)
ATM	1 (1.1)	-	-	1 (0.7)	2 (0.4)
AMP-CFZ	4 (4.4)	1 (1.1)	1 (0.9)	5 (3.7)	11 (2.5)
CFZ-CFO	3 (3.3)	2 (2.2)	2 (1.8)	1 (0.7)	8 (1.8)
AMP-TET	-	1 (1.1)	2 (1.8)	1 (0.7)	4 (0.9)
AMC-AMP	1 (1.1)	1 (1.1)	2 (1.8)	-	3 (0.7)
AMP-SUT	1 (1.1)	-	1 (0.9)	-	2 (0.4)
SUT-TET	-	1 (1.1)	1 (0.9)	-	2 (0.4)
CFO-SUT	1 (1.1)	-	-	-	1 (0.2)
AMI-CLO	-	1 (1.1)	-	-	1 (0.2)
AMP-CFO	-	1 (1.1)	-	-	1 (0.2)
CFO-TET	-	-	1 (0.9)	-	1 (0.2)
ATM-CAZ	1 (1.1)	-	-	-	1 (0.2)

AMC-CFZ-AMP	4 (4.4)	4 (4.4)	9 (8.1)	4 (2.9)	21 (4.8)
AMC-CFO-AMP	-	-	-	-	5 (1.1)
AMP-AMC-CFZ	-	-	-	-	7 (1.6)
AMP-CLO-SUT	1 (1.1)	-	1 (0.9)	-	2 (0.4)
AMP-AMC-CFO	1 (1.1)	-	-	-	1 (0.2)
AMP-CFZ-CFO	1 (1.1)	-	-	-	1 (0.2)
CFZ-SUT-TET	-	1 (1.1)	-	-	1 (0.2)
AMP-CFZ-TET	-	-	1 (0.9)	-	1 (0.2)
AMP-CIP-CLO	-	-	1 (0.9)	-	1 (0.2)
AMP-SUT-TET	-	-	-	1 (0.7)	1 (0.2)
CLO-SUT-TET	-	-	-	1 (0.7)	1 (0.2)
AMP-AMC-CFZ-CFO	14 (15.4)	12 (13.2)	10 (8.9)	6 (4.4)	42 (9.7)
AMP-CFZ-CFO-CLO	2 (2.2)	-	1 (0.9)	-	3 (0.7)
AMP-AMC-CFZ-TET	1 (1.1)	-	-	-	1 (0.2)
AMP-CIP-LEV-TET	1 (1.1)	-	-	-	1 (0.2)
AMP-AMC-CFZ-CAZ	1 (1.1)	-	-	-	1 (0.2)
AMC-CFZ-CFO-TET	-	-	-	1 (0.7)	1 (0.2)
AMP-AMC-CFZ-CFO-TET	1 (1.1)	1 (1.1)	1 (0.9)	1 (0.7)	4 (0.9)
AMP-AMC-CFZ-CLO-SUT	-	-	2 (1.8)	-	2 (0.4)
ATM-CPM-CFO-CAZ-CRO	-	-	-	2 (1.5)	2 (0.4)
AMP-AMC-ATM-CFZ-CAZ	1 (1.1)	-	-	-	1 (0.2)
AMP-AMC-CFZ-CFO-CLO	1 (1.1)	-	-	-	1 (0.2)
AMP-AMC-CIP-CLO-LEV-TET	1 (1.1)	1 (1.1)	2 (1.8)	-	4 (0.9)

AMP = ampicillin, 10 µg); ATM = aztreonam, 30 µg; AMC = amoxicillin/clavulanic acid, 30 µg, AMI = amikacin, 30 µg, GEN gentamicina, 10 µg; CIP = ciprofloxacin, 5 µg; LEV = levofloxacin, 5 µg; CFZ = cefazolin, 30 µg; CFO = ceftazidime, 30 µg; CAZ = ceftazidime, 30 µg; CRO = ceftriaxone, 30 µg; COM = cefepime, 30 µg; CLO = chloramphenicol, 30 µg; MPM = meropenem, 10 µg; SUT = sulfamethoxazole/trimethoprim, 25 µg; TET = tetracycline, 30 µg.