



Biodiversity of Sematophyllaceae s. str. in northeastern Brazil

Milena Evangelista-dos-Santos^{1*}, Emilia de B. Valente²

1 Programa de Pós-Graduação em Botânica, Herbário da UEFS, Universidade Estadual de Feira de Santana, Feira de Santana, BA, Brazil • milenaevangelista16@gmail.com  <https://orcid.org/0000-0003-0816-2832>

2 Departamento de Ciências Biológicas, Universidade Estadual de Feira de Santana, Feira de Santana, BA, Brazil • ebvalente@gmail.com  <https://orcid.org/0000-0002-9801-2370>

* Corresponding author

Abstract

In this study, we list 25 moss species belonging to Sematophyllaceae s. str. Six species are reported for the first time from some of the states in the Northeast Region of Brazil. *Sematophyllum succedaneum* (Hook.f. & Wilson) Mitt. stands out for being found for the first time in this region. The Atlantic Forest is the phytogeographic domain with the greatest number of Sematophyllaceae species, which are either exclusive to this domain or shared with the Caatinga and/or Cerrado. Our study expands on the previous knowledge of these species and provides a foundation for further studies.

Keywords

Atlantic Forest, Caatinga, Cerrado, Hypnales, pleurocarpous mosses

Academic editor: Anna Luiza Ilkiu-Borges | Received 1 July 2022 | Accepted 10 October 2022 | Published 26 October 2022

Citation: Evangelista-dos-Santos ME, Valente EB (2022) Biodiversity of Sematophyllaceae s. str. in northeastern Brazil. Check List 18 (5): 1181–1189. <https://doi.org/10.15560/18.5.1181>

Introduction

The Northeast Region of Brazil is the third largest region and has the largest number of bryophyte species, with an estimated 689 species (Costa and Peralta 2015). This region presents four phytogeographic domains: Caatinga, Cerrado, Amazon Forest (in Maranhão), and Atlantic Forest. According to Drumond et al. (2000), the Caatinga is the largest domain occupying the central, northeastern, and southern area of the Northeast Region of Brazil. The Cerrado occupies the western and southeastern portions, whereas the Atlantic Forest, with its area drastically reduced, lies in the easternmost portion of the Northeast region. Although the Caatinga covers the largest territory, this domain has the lowest species diversity

of bryophytes (Costa and Peralta 2015). The predominant ombrophilous forest in the Atlantic region is characterized by a high diversity (Costa and Peralta 2015).

Sematophyllaceae is a large moss family with approximately 49 species in Brazil (Câmara and Carvalho-Silva 2020). Descriptions of Sematophyllaceae species from the Northeast Region are found in the publications by Yano et al. (2009), Varão et al. (2011), Oliveira et al. (2018), Evangelista-dos-Santos et al. (2021), Evangelista-dos-Santos and Almeida (2020), and Souza and Oliveira (2019). Although there are existing published studies on Sematophyllaceae, additional investigations of herbarium collections are required to provide invaluable information.

The five herbaria with the most specimens collected in the Northeast Region are ALCB, HABIT, HUEFS, UB, and UFB (SpeciesLink 2021). Approximately 126 Sematophyllaceae specimens were found on the SpeciesLink website without identifications to species. Here, we provide a list of all Sematophyllaceae s. str. species occurring in the Northeast Region of Brazil, with taxonomic comments, photographs, ecological information, and geographic and altitudinal distribution.

Study Area

The Northeast Region of Brazil is located in the Southern Hemisphere tropics (Fig. 1). In the north and east borders the Atlantic Ocean. Rainfall is between 300 and 2220 mm per year, and the average temperature is between 20 and 40° C (Queiroz et al. 2018). The Northeast Region has a semi-arid climate (Araújo and Martins 1998). The region has the following physiognomies: Savana-estépica, Caatinga do Sertão, arid region with vegetational disjunctions, dense ombrophilous forest, ombrophilous open forest, seasonal semi-deciduous forest, seasonal deciduous forest, and savanna (IBGE 2012).

Methods

To obtain a preliminary list of the Sematophyllaceae s. str. species in the Northeast Region, we compiled data from published sources (i.e., Yano 1981, 1989, 1994, 2011; Costa et al. 2011; Varão et al. 2011; Oliveira et al. 2018; Evangelista-dos-Santos et al. 2019; Souza and Oliveira 2019). In addition, we consulted the online platforms SpeciesLink (<https://specieslink.net/>) and Flora e Funga do Brasil 2020 (<http://floradobrasil.jbrj.gov.br>).

For identification of Sematophyllaceae s. str. species, specialized literature was consulted (i.e., Florschutz 1964; Sharp et al. 1994; Buck 1994, 1998; Ramsay et al. 2004; Câmara et al. 2016), and type materials from NY, NHM, and PACA herbaria were requested on loan.

We undertook collection excursions in areas where fewer or dubious records existed (i.e., Serra da Jibóia, BA; Barra do Vento, BA; and Serra da Bela Vista, SE). Between 2019 and 2022, 212 samples were collected following the methodology proposed by Yano (1984). The collected material is deposited in HUEFS and ASE. Specimens were requested from the Alexandre Leal Costa (ALCB) and Anchieta (PACA) herbaria and analyzed in HUEFS. The specimens housed in the herbaria ASE, HABIT, HUESPI, HUNEB, UB, and UFP were observed during visits in 2019 and 2021. The taxonomic classification used here follows Carvalho-Silva et al. (2017) and Deshmukh et al. (2021).

The species are classified according to the substrate they were colonizing (Robins 1952; Bates 2009). The Brazilian phytogeographic domains follow the classification proposed by Fiaschi and Pirani (2009). Of the 25 species included in the list of species, we selected 13 to be described and illustrated, as these species are newly reported from Northeast Region or with little information in the literature on them.

Results

Twenty-five species belonging to the family Sematophyllaceae s. str. (Table 1) are known from the Northeast Region of Brazil. Twenty-three species occur in the Atlantic Forest, of which 12 are exclusive to that domain, and 11 are shared with the Cerrado and Caatinga




Figure 1. Map showing the study area (northeastern Brazil) and its phytogeographic domains.

Table 1. Check list of Sematophyllaceae s. str. species in the Northeast Region of Brazil. Phytogeographic domain: CA = Caatinga, CE = Cerrado, AF = Atlantic Forest. Briocenose: Co = Corticolous, Ep = Epimicont, Ex = Epixilous, Ru = Rupicolous, Te = Terricolous, Tm = Termiticola. Distribution in northeastern Brazilian state: AL = Alagoas, BA = Bahia, CE = Ceará, MA = Maranhão, PE = Pernambuco, PI = Piauí and SE = Sergipe. * New records.

Species	Phytogeographic domain	Briocenose	Elevation	Distribution in Northeastern Brazil by state
<i>Acroporium pungens</i> (Hedw.) Broth.	AF, CA	Co	93–800	BA, CE
<i>Aptychopsis estrellae</i> (Müll.Hal) P.E.A.S.Câmara, Carv.-Silva & W.R.Buck	AF	Co, Ex	800–1200	AL, BA, PE
<i>Aptychopsis pungifolia</i> (Hampe.) Broth.	AF	Co, Ex	392–800	BA, PE
<i>Aptychopsis pyrrhophylla</i> (Müll.Hal) Wijk & Margad.	AF	Co, Ru	1500–1900	BA
<i>Aptychopsis subpungifolia</i> (Broth.) Broth.	AF	Co, Ex	800–1000	BA
<i>Aptychopsis tequendamensis</i> (Hampe) P.E.A.S.Câmara, Carv.-Silva & W.R.Buck	AF	Co, Ex	800	BA
<i>Brittonodoxa lithophila</i> (Hornsch.) W.R.Buck, P.E.A.S. Câmara & Carv.-Silva	AF	Co	500	BA
<i>Brittonodoxa subpinnata</i> (Brid.) W.R.Buck, P.E.A.S.Câmara & Carv.-Silva	AF, CA, CE	Co, Ex, Ru, Te	50–1200	AL, BA, CE, MA, PE, PI, SE
<i>Colobodontium vulpinum</i> (Mont.) S.P.Churchill & W.R. Buck	AF, CE	Ru	1289	BA
<i>Donnellia commutata</i> (Müll.Hal.) W.R.Buck	AF, CA, CE	Co, Ex, Ru	50–1200	AL*, BA, CE, MA*, PE, SE
<i>Donnellia lageniformis</i> (Müll.Hal.) W.R.Buck	CA	Ex		BA
<i>Jirivanaea caespitosa</i> (Hedw.) U.B.Deshmukh & Rathor	AF, CA	Co, Ru	1289–1734	BA
<i>Jirivanaea cuspidifera</i> (Mitt.) U.B.Deshmukh & Rathor	AF, CA	Co, Ex, Ru	500–924	BA, CE*, MA
<i>Jirivanaea galipensis</i> (Müll. Hal.) U.B.Deshmukh & Rathor	AF, CA, CE	Co, Ep, Ex, Ru, Te, Tm	92–1690	BA, CE, MA, PE, PI, SE*
<i>Meiothecium boryanum</i> (Müll.Hal) Mitt.	AF	Co	50–383	BA
<i>Sematophyllum beyrichii</i> (Hornsch.) Broth.	AF, CA, CE	Co, Ex, Ru, Te	70–1734	AL, BA, CE, MA, PE, SE*
<i>Sematophyllum succedaneum</i> (Hook.f. & Wilson) Mitt.	AF	Co	800	BA*
<i>Sematophyllum swartzii</i> (Schw gr.) W.H.Welch & H.A.Crum	AF, CA	Ex, Ru	500–1200	BA, PE*
<i>Trichosteleum brachydictyon</i> (Besch.) A.Jaeger	AF	Ex		BA
<i>Trichosteleum glaucinum</i> (Besch.) A.Jaeger	AF	Co	583	BA
<i>Trichosteleum glaziovii</i> W.R.Buck	AF, CA	Ru		BA
<i>Trichosteleum lonchophyllum</i> (Mont.) Carv.-Silva, P.E.A.S. Câmara & W.R.Buck	AF, CE	Co, Ru, Te		BA
<i>Trichosteleum papillosum</i> (Hornsch.) A.Jaeger	AF, CA	Ex	50–200	BA, PE, SE
<i>Trichosteleum sentosum</i> (Sull.) A.Jaeger	AF	Co, Te	92–383	BA
<i>Trichosteleum subdemissum</i> (Besch.) A.Jaeger	CA, CE	Co, Ex, Ru, Te		BA, MA, PI

phytogeographic domains. Only *Donnellia lageniformis* (Müll.Hal.) W.R. Buck was found exclusively in campo rupestre in the Caatinga domain, growing on decaying wood. *Sematophyllum succedaneum* (Hook.f. & Wilson) Mitt. is newly recorded from the Northeast Region, and the geographic distributions of five other species are expanded: *Donnellia commutata* (Müll.Hal.) W.R.Buck, *Jirivanaea cuspidifera* (Mitt.) U.B.Deshmukh & Rathor, *J. galipensis* (Müll. Hal.) U.B.Deshmukh & Rathor, *Sematophyllum beyrichii* (Hornsch.) Broth., and *S. swartzii* (Schwägr.) W.H.Welch & H.A.Crum. The Brazilian states with the most species are Bahia (25 spp.), Pernambuco (8 spp.), and Maranhão and Ceará (each with 6 spp.). Altitudinal variation ranged between 93 and 1900 m. The diversity of substrates was great, with species growing on tree trunks, decaying logs, rocks, soil, fungi, and termite mounds. The substrate most colonized by Sematophyllaceae s. str. species was logs and decaying wood.

Aptychopsis estrellae (Müll.Hal) P.E.A.S.Câmara, Carv.-Silva & W.R.Buck

Figure 2A–C

Material examined. BRAZIL – Alagoas • Murici, Estação Ecológica de Murici; 09°22'46"S, 035°50'28"W; 02.XII.2004; K.C. Pôrto leg.; UFP 48852 – Bahia • Miguel Calmon, Parque Estadual das Sete Passagens; 11°38'14"S, 040°52'25"W; 1200 m alt.; 11.X.2007; J.

Ballejos 1982; ALCB 79378 • Santa Teresinha, Serra da Jiboia; 12°44'55"S, 039°47'55"W; 800 m alt.; 17.IX.2015; S.B.V. Bôas-Bastos 2734; ALCB 119927 – Pernambuco • Caruaru, Mata dos Altinhos, Brejo dos Cavalos; 08°28'33"S, 035°17'11"W; 25.XIII.1987; K.C. Pôrto 2443; UFP 20827 • São Vicente Férrer, Serra das Marendas; 07°47'45"S, 035°35'14"W; 23.III.2010; M.P.P. Silva 279; UFP 72135.

Identification. Leaves lanceolate to lanceolate-triangular, concave, rarely plane, apex gradually acuminate, margin subentire; cells linear, flexuose, thick-walled, smooth, porose at base; alar cells curved or not to insertion, enlarged, inflated, oblong; supra-alar cells quadratic, 1–2 rows. Capsules cylindric; exostome teeth triangular, bordered or not, cross-striolate below, trabeculate at back; endostome with a medium-high basal membrane, segments papillose, perforate, cilia single sometimes rudimentary.

Aptychopsis pyrrhophylla (Müll.Hal) Wijk & Margad.

Figure 2D–F

Material examined. BRAZIL – Bahia • Abaíra, Catolés, Serra do Barbado, Mata da Forquilha; 13°19'47"S, 041°53'12"W; 1655 m alt.; 05. IX.2008; C. Bastos 5166; ALCB 84217 • Rio de Contas, Pico das Almas; 13°57'59"S, 041°51'14"W; 1600 m alt.; 27.XI.1998; R.M. Harley 26634; HUEFS 28246.




Figure 2. Sematophyllaceae s. str. species in the Northeast Region of Brazil. **A–C.** *Aptychopsis estrellae*. **D–F.** *Aptychopsis pyrrhophylla*. **G–I.** *Aptychopsis tequendamensis*. **J–L.** *Brittonodoxa lithophila*. **M–O.** *Colobodontium vulpinum*. **P–R.** *Donnellia commutata*. Scale bars: A, D, G, J, M, P = 200 µm; B, C, E, F, H, I, L, O, R = 50 µm; K, N, Q = 20 µm.

Identification. Leaves oblong-lanceolate, concave or plane, apex long-acuminate, margins entire, revolute or not; cells linear-flexuose, thick-walled, smooth, porose at base; alar cells curved to the insertion, enlarged, inflated, oblong; supra-alar cells quadrate to subrectangular, 3–4 rows. Capsule not observed.

***Aptychopsis tequendamensis* (Hampe)**

P.E.A.S.Câmara, Carv.-Silva & W.R.Buck

Figure 2G–I

Material examined. BRAZIL – Bahia • Eunápolis, Estação Ecológica Veracruz, divisa com Imbiricu; 16°36'06"S, 039°16'26"W; 09.IX.1999; C. Bastos & S.B.V. Bôas-Bastos 1727; ALCB 41679 • Santa Teresinha, Serra da Jiboia; 12°55'55"S, 039°47'55"W; 800 m alt.; 12.III.2002; C. Bastos & S.B.V. Bôas-Bastos 3155; ALCB 51322.

Identification. Leaves lanceolate to lanceolate-triangular, plane, apex long-acuminate, margins mostly entire; cells linear-flexuose, short linear at apices, thick-walled, smooth, porose at base; alar cells rarely incurved to insertion, enlarged, rectangular to short-rectangular; supra-alar cells quadrate to rectangular, 1 row. Capsules ovoid to cylindric; exostome teeth triangular, bordered, cross-striolate below, trabeculate at back; endostome with a high basal membrane, segments finely papillose, perforate, cilia single.

***Brittonodoxa lithophila* (Horns.) W.R.Buck,**

P.E.A.S.Câmara & Carv.-Silva

Figure 2J–L

Material examined. BRAZIL – Bahia • Nova Esperança, Estação Ecológica Wenceslau Guimarães, Trilha Água Vermelha; 13°35'43"S, 039°43'10"W; 550 m alt.; 04.XII.2017; C. Bastos 6147; ALCB 13044.

Identification. Leaves orbicular to ovate, concave, rarely plane, apex mucrunate or obtuse, margin subentire; cells rhomboidal, thick-walled, smooth, not porose at base; alar cells not curved to insertion, inflated or not, oblong to short-rectangular; supra-alar cells quadratic to subrectangular, 2–4 rows. Capsule cylindric; exostome teeth triangular, bordered, cross-striolate below, trabeculate at back; endostome with a high basal membrane, segments papillose, perforate, cilia single.

***Colobodontium vulpinum* (Mont.) S.P.Churchill & W.R.Buck**

Figure 2M–O

Material examined. BRAZIL – Bahia • Piatã, Cachoeira do Patrício; 13°08'35"S, 041°55'46"W; 1289 m alt.; 04.VI.2007; E.B. Valente 622; HUEFS 153867.

Identification. Leaves oval, ovate or semi-orbiculate, concave or plane, apex acute or obtuse, margin subentire; cells rhomboidal to rhombic, thick-walled, smooth, not porose at base; alar cells not curvate to insertion, inflated or not, oblong; supra-alar cells quadratic, 2–4 rows. Capsule cylindric; exostome teeth triangular, not bordered,

cross-striolate below, trabeculate at base papillose above; endostome with segments filiform or rudimentary, cilia absent.

***Donnellia commutata* (Müll.Hal.) W.R.Buck**

Figure 2P–R

Material examined. BRAZIL – Alagoas • Murici, Estação Ecológica de Murici; 09°22'46"S, 035°50'28"W; 02.XII.2004; K.C. Pôrto leg.; UFP 48981 – Bahia • Boa Nova, Parque Nacional de Boa Nova, Recanto dos Pásaros; 14°39'20"S, 040°14'46"W; 20.I.2010; A.M. Souza 590; HUEFS 204888 • Igrapiúna, Reserva Ecológica da Michelin; 13°47'19"S, 039°11'51"W; 04.IV.2010; H.C. Oliveira 1737; ALCB 112639 • Itaberaba, Serra do Orobó; 12°35'13"S, 040°46'27"W; 752 m alt.; 10.X.2019; C.J.P. Bastos 5580; ALCB 119569 • Miguel Calmon, Parque Estadual das Sete Passagens; 11°38'14"S, 040°52'25"W; 1200 m alt.; 19.II.2007; J. Ballejos 1214; ALCB 77318 • Morro do Chapéu, Mata do Capão do Pinho; 929 m alt.; 17.VI.2008; E.B. Valente 1196; HUEFS 154441 • Santa Teresinha, Serra da Jiboia; 12°55'55"S, 039°47'55"W; 800 m alt.; 24.IX.2004; E.B. Valente 402; HUEFS 97274 – Ceará • Crato, Chapada do Araripe; 07°38'72"S, 040°21'70"W; 920 m alt.; 19.III.1983; S. Prazeres & K.C. Pôrto leg.; UFP 6069 – Maranhão • Carolina, RPPN Mansinha; 07°10'49"S, 047°41'21"W; 13.VII.2016; R.R.O. 493; HABIT 3187 – Pernambuco • Timbaúba, Engenho, Água Azul; 17.VII.1992; S.R. Germano leg; UFP 8683.

Identification. Leaves lanceolate, concave or plane, apex acuminate, margin entire; cells rhomboidal to long-rhomboidal, thick-walled, smooth, not porose at base; alar cells not curved to insertion, oblong; supra-alar cells quadratic, 2–4 rows. Capsules cylindric; exostome teeth triangular, bone-white, not bordered, with prominent cross walls, slightly trabeculate at back; endostome reduced with a very low basal membrane, sometimes absent, cilia absent.

***Sematophyllum beyrichii* (Horns.) Broth.**

Figure 3A–C

Material examined. BRAZIL – Alagoas • Murici, Estação Ecológica de Murici; 09°22'46"S, 035°50'38"W; 21.XII.2004; K.C. Pôrto leg.; UFP 48977 – Bahia • Abaíra, Catolés, Serra do Barbado, Mata do Cigano; 13°19'47"S, 041°53'12"W; 1734 m alt.; 06.IX.2008; C. Bastos 5246; ALCB 84328 • Boa Nova, Parque Nacional de Boa Nova, Recanto dos Pásaros; 14°39'20"S, 040°14'56"W; 27.VI.2013; A.M. Souza 653; HUEFS 204959 • Eunápolis, Estação Ecológica Veracruz, trilha pau-brasil; 16°36'60"S, 039°16'60"W; 08.IX.1999; C. Bastos & S.B.V. Bôas-Bastos 1639; ALCB 45051 • Igrapiúna, Reserva Ecológica da Michelin; 13°47'19"S, 039°11'51"W; 383 m alt.; 03.IV.2010; H.C. Oliveira 1679; ALCB 112604 • Itaberaba, Serra do Orobó; 12°35'13"S, 040°46'67"W; 752 m alt.; 18.I.2015; R. Freitas 120; ALCB 119497 • Miguel Calmon, Parque Estadual das Sete Passagens; 11°38'14"S, 040°52'25"W; 1200 m alt.; 19.II.2007; J. Ballejos 1256; HUEFS 187538 • Santa




Figure 3. Sematophyllaceae s. str. species in the Northeast Region of Brazil. **A–C.** *Sematophyllum beyrichii*. **D–F.** *Sematophyllum succedaneum*. **G–I.** *Trichosteleum brachydictyon*. **J–L.** *Trichosteleum glaucinum*. **M–O.** *Trichosteleum glaziovii*. **P–R.** *Trichosteleum papillosum*. **S–U.** *Trichosteleum sentosum*. Scale bars: A, D, G, J, M, P, S = 200 µm; B, C, H, I, N, O, Q, R, T, U = 50 µm; F, K, L = 20 µm; E = 10 µm.

Teresinha, Serra da Jiboia; 12°55'55"S, 039°47'55"W; 800 m alt.; 03.IV.1999; E. Mello 2670; HUEFS 36594 – **Ceará** • Ubajara, Serra da Ibiapaba, São Luís; 03°58'45"S, 041°09'00"W; 924 m alt.; 24.VIII.2010; C. Bastos 5360; ALCB 97126 – **Maranhão** • Mirador; 30.VI.2008; E.S. Brito & G.M. Conceição leg.; HABIT 574 – **Pernambuco** • Reserva Ecológica de Garajau, Mata do encruamento; 06.VI.2000; S.R. Germano leg.; UFP 32934.

Identification. Leaves lanceolate or oblong-lanceolate, convave, rarely plane, apex acuminate; cells linear, flexuose, thick-walled, smooth, not porose at base; alar cells not curved to insertion, inflated, oblong; supra-alar cells quadratic, 1–2 rows. Capsules cylindric; exostome teeth triangular, bordered, cross-striolate below, trabeculate at back; endostome with a high basal membrane, segments papillose, perforate or not, cilia none or rudimentary.

Sematophyllum succedaneum (Hook.f. & Wilson)

Mitt.

Figure 3D–F

Material examined. BRAZIL – **Bahia** • Santa Teresinha, Serra da Jiboia; 12°55'55"S, 039°47'55"W; 800 m alt.; 13.IX.1997; E. Melo 2217; HUEFS 29954.

Identification. Leaves oblong-lanceolate, plane or concave, apex acuminate, margin entire to subentire; cells rhomboidal, flexuose, thick-walled, smooth, not porose at base; alar cells not curved to insertion, inflated or not, oblong; supra-alar cells quadratic, 1 row. Capsules cylindric; exostome teeth triangular, not bordered, cross-striolate below, trabeculate at back; endostome with high basal membrane, segments papillose, perforate, cilia single.

Trichosteleum brachydictyon (Besch.) A.Jaeger

Figure 3G–I

Material examined. BRAZIL – **Bahia** • Eunápolis, Estação Ecológica Veracruz, fazenda Malacarne; 16°36'00"S, 039°16'00"W; C. Bastos & S.B.V. Bôas-Bastos 2129; ALCB 44918.

Identification. Leaves lanceolate or oblong-lanceolate, concave, apex long-acuminate, rarely twisted, margins serrulate; cells linear, flexuose, thick-walled, unipapillose in ½ of leaf, porose at base; alar cells not curved to insertion, enlarged, inflated, oblong; supra-alar cells quadratic, 1–2 rows. Capsules cylindric; exostome teeth triangular, bordered, cross-striolate below, trabeculate at back; endostome with a high basal membrane, segments papillose, perforate, cilia single.

Trichosteleum glaucinum (Besch.) A.Jaeger

Figure 3J–L

Material examined. BRAZIL – **Bahia** • Nova Esperança, Estação Ecológica Wenceslau Guimarães, Trilha Água Vermelha; 13°35'43"S, 039°43'10"W; 583 m alt.; 29.IX.2017; R.R. Ferminano 102; ALCB 124284.

Identification. Leaves lanceolate or oblong-lanceolate, concave or plane, apex short-acuminate, rarely twisted, margins serrulate; cells linear to rhomboidal,

think-walled, unipapillose in ¾ of leaf, porose in the insertion; alar cells not curved to insertion, inflated, ovate to oblong; supra-alar cells not differentiated. Capsule not observed.

Trichosteleum glaziovii W.R.Buck

Figure 3M–O

Material examined. BRAZIL – **Bahia** • Miguel Calmon, Parque Estadual das Sete Passagens; 11°38'14"S, 040°52'25"W; 1200 m alt.; 22.IV.2003; C. Bastos & S.B.V. Bôas-Bastos 3699; ALCB 51783.

Identification. Leaves falcate, lanceolate, concave, apex acuminate, margins serrulate; cells linear, flexuose, thick-walled, unipapillose, sometimes papillae are inconspicuous, porose at base; alar cells not curved to insertion, inflated, oblong; supra-alar cells quadratic 1 row. Capsule not observed.

Trichosteleum papillosum (Hornsch.) A.Jaeger

Figure 3P–R

Material examined. BRAZIL – **Bahia** • Boa Nova, Parque Nacional de Boa Nova, Recanto dos Pássaros; 14°39'20"S, 040°14'96"W; 27.VI.2013; A.M. Souza 678; HUEFS 204984 • Una, Reserva Ecológica do Mico-Leão; 14°47'00"S, 039°03'17"W; 100 m alt.; 26.IV.1981; B.M. Boom 794; CEPLAC 49719 • Nova Esperança, Estação Ecológica Wenceslau Guimarães, Trilha Água Vermelha; 13°35'43"S, 039°43'10"W; 583 m alt.; 29.IX.2017; L.C. Reis 629; ALCB 96976 – **Pernambuco** • Recife, Mata de Dois Irmãos, perto da Sede do Reservatório; 08°05'38"S, 034°56'26"W; 03.VIII.1998; O. Yano et al. 23412; UB 204790 – **Sergipe** • Itabaiana, Serra de Itabaiana; 10°41'50"S, 037°25'29"W; 190 m alt.; 13.I.1981; D. Andrade-Lima 6744; UB 204680.

Identification. Leaves lanceolate or oblong-lanceolate, plane or concave, apex acuminate to crispatate, margins serrulate; cells fusiform to linear fusiform, thick-walled, unipapillose in ½ of leaf, porose at base; alar cells not curved to insertion, inflated, oblong; supra-alar cells quadrate, 1–2 rows. Capsule not observed.

Trichosteleum sentosum (Sull.) A.Jaeger

Figure 3S–U

Material examined. BRAZIL – **Bahia** • Boa Nova, Parque Nacional de Boa Nova, Recanto dos Pássaros; 14°39'20"S, 040°14'16"W; 27.VI.2013; A.M. Souza 608; HUEFS 204907 • Igrapiúna, Reserva Ecológica da Michelini; 13°47'19"S, 039°11'51"W; 383 m alt.; 12.VIII.2006; S.B.V. Bôas-Bastos 2224; ALCB 77065 • Nova Esperança, Estação Ecológica Wenceslau Guimarães, Trilha Água Vermelha; 13°35'43"S, 039°43'10"W; 583 m alt.; 29.VII.2019; R.R. Ferminano; ALCB.

Identification. Leaves lanceolate or oblong-lanceolate, concave or plane, apex long-acuminate, flexuose and twisted, margins serrate; cells linear, thick-walled, unipapillose in ¾ of leaf, porose at base; alar cells not curved to insertion, inflated, oblong; supra-alar cells quadrate,

1–2 rows. Capsules cylindric; exostome teeth triangular, bordered, cross-striolate below, trabeculate at back; endostome with a high basal membrane, segments papillose, perforate, cilia single rarely in pairs.

Discussion

According to Câmara and Carvalho-Silva (2022), Sematophyllaceae s.lat. is represented in the Northeast Region of Brazil by 23 species, five of which currently belong to the family Pylaisiadelphaceae Goffinet & W.R.Buck. Hence, 18 species of Sematophyllaceae s. str. were known from this region (Carvalho-Silva et al. 2017). In Brazil, 38 species of Sematophyllaceae s. str. are recognized (Câmara and Carvalho-Silva 2022), and with 25 species, in the Northeast Region has 65.7% of the total reported for the country. All species were found in the state of Bahia (Evangelista-dos-Santos et al. 2021).

The species reported from the Northeast Region are similar to the other Brazilian regions, especially the South and Southeast Regions, according to the published literature (i.e., Visnadi 2006; Amélio 2019; Faria et al. 2020; Koga and Peralta 2021; Remor et al. 2021). The South and Southeast Regions are predominantly within the Atlantic Forest domain, which has the largest number of species of the family in Brazil, which justifies the similarity (Costa and Peralta 2015).

With our new data from previously unexplored locations and new records from the Caatinga or Cerrado domain, the number species known from the Northeast Region is considerably increased. The occurrence of the relatively larger number of Sematophyllaceae s. str. species found in this region can be explained by the variety of habitats, floristic heterogeneity, and environmental conditions (Giulietti et al. 2005; Moraes et al. 2020). Greater availability of water, which provides ideal microhabitats for bryophytes may also be of importance (Vanderpoorten and Goffinet 2009).

Although cited in the literature, some species have not since been recorded in certain places, such as *Aptychopsis estrellae* cited for the Serra de Itabaiana in Sergipe (Silva and Pôrto 2015), *Jirivanaea caespitosa* (Hedw.) U.B.Deshmukh & Rathor for the Serra do Cariri in Pernambuco (Yano 1981), and *Trichosteleum sentosum* for Pernambuco with no defined location (Costa et al. 2011; Yano 2011). These are likely misidentifications. The specimens originally identified as *A. estrellae* were found to be *Sematophyllum beyrichii*.

Sematophyllum adnatum (Michx.) E.Britton is frequently cited for the Northeast Region (Alvarenga et al. 2007; Yano et al. 2009; Costa et al. 2011; Yano 2011; Evangelista-dos-Santos et al. 2019), but it is excluded here, as specimens identified as this species instead belong to *S. beyrichii*. *Donnellia commutata* requires a detailed analysis of the sporophytic characteristics for its identification. In the absence of sporophytes, the gametophytes on leaves resemble some forms of *Brittonodoxa subpinnata* (Evangelista-dos-Santos et al. 2021).

Jirivanaea cuspidifera (Mitt.) U.B. Deshmukh & Rathor is often confused with *Sematophyllum beyrichii*.

The species analyzed in this study occur preferentially in the Atlantic Forest, and 10 occur exclusively in this domain (Fig. 1). In contrast, only *Donnellia lageniformis* (Müll.Hal.) W.R.Buck was found exclusively in the Caatinga, in campo rupestre, growing on decaying plant material. The presence of this species is known in the South and Southeast regions of the country, such as riparian forest formations or gallery and mixed ombrophilous forest of the Atlantic region (Câmara and Carvalho-Silva 2022). Its identification was possible thanks to the presence of the sporophyte that made it possible to distinguish between *D. commutata* (Buck 1994; Evangelista-dos-Santos et al. 2021).

According to Gradstein and Pócs (1989) and Frahm (1990), most bryophytes colonize the trunks of living trees, and pH and a rough cortex help in their establishment. Few species grow on only one type of substrate, such as *Acroporium pungens* (Hedw.) Broth. on the bases of living tree trunks, *Colobodontium vulpinum* on rocks, *Trichosteleum papillosum* on decaying plant material, and *Trichosteleum glaziovii* on rocks.

Our results show the importance of analyzing herbarium collections in better understanding species. Many of the new occurrences records come from materials originally identified at the family or genus level. Based on our results we highlight the important contribution of taxonomic study of herbarium collections.

Authors' Contributions

Conceptualization: MES. Data curation: MES. Formal analysis: MES. Investigation: MES. Methodology: MES. Project administration: MES. Resources: MES. Software: MES. Supervision: EBV. Validation: MES. Visualization: MES. Writing – original draft: MES. Writing – review and editing: MES.

Acknowledgements

This study was funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the scholarship granted to the first author. We thank the Programa de Pós-Graduação em Botânica of Universidade Estadual de Feira de Santana for the award of doctoral degree, the personnel at the herbaria ALCB, ASE, HUESPI, HUEFS, HUNEB, HABIT, NY, NHM, PACA, UB, and UFP for their support during our visits and for loans of specimens, and Dr. Paulo Eduardo Aguiar Saraiva Câmara for his help in the identification of some species.

References

- Alvarenga LDP, Silva MPP, Oliveira JRPM, Pôrto KC (2007) Novas ocorrências de briófitas para Pernambuco, Brasil. Acta Botanica Brasiliensis 21 (2): 349–360. <https://doi.org/10.1590/S0102-33062007000200009>

- Amélio LA, Peralta DF, Carmo DM (2019) Briófitas do Parque Estadual de Campos do Jordão, estado de São Paulo, Brasil. Hoehnea 46 (2): e962018. <https://doi.org/10.1590/2236-8906-96/2018>
- Araújo FS, Martins FR (1998) Fisionomia e organização da vegetação do Carrasco no planalto da Ibiapaba, estado do Ceará. Acta Botânica Brasiliensis 13 (1): 1–13. <https://doi.org/10.1590/S0102-33061999000100002>
- Bates JW (2009) Mineral nutrition and substratum ecology. In: Goffinet B, Shaw AJ. Bryophyte biology. Cambridge University Press, Cambridge, UK, 299–356.
- Buck WR (1998) Pleurocarpous mosses of the West Indies. Memoirs of the New York Botanical Garden 82: 1–400.
- Buck WR (1994) A synopsis of the American species of *Donnellia* (Sematophyllaceae). Hikobia 11: 377–385.
- Câmara PEAS, Carvalho-Silva M, Buck WR (2016) The genus *Acroporium* (Sematophyllaceae) in the neotropics. Journal of Bryology 37 (4): 284–291. <https://doi.org/10.1179/1743282014Y.0000000126>
- Câmara PEAS, Carvalho-Silva M (2020). Sematophyllaceae. Flora do Brasil 2020. Jardim Botânico do Rio de Janeiro. <http://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB96943>. Accessed on: 2022-04-10.
- Carvalho-Silva M, Stech M, Soares-Silva LH, Buck WR, Wickett NJ, Liu Y, Câmara P EAS (2017) A molecular phylogeny of the Sematophyllaceae s.l. (Hypnales) based on plastid, mitochondrial and nuclear markers, and its taxonomic implications. Taxon 66 (4): 811–831. <https://doi.org/10.12705/664.2>
- Costa DP, Peralta DF (2015) Bryophytes diversity in Brazil. Rodriguésia 66 (4): 1063–1071. <https://doi.org/10.1590/2175-7860201566409>
- Costa DP, Pôrto KC Luizi-Ponzo AP, Ilkiu-Borges AL, Bastos CJP, Câmara PEAS, Peralta DF, Bôas-Bastos SBV, Imbassahy CAA, Henriques DK, Gomes HCS, Rocha L M, Santos ND, Siviero TS, Vaz-Imbassahy TF, Churchill SP (2011) Synopsis of the Brazilian moss flora: checklist, distribution and conservation. Nova Hedwigia 93: 277–334. <https://doi.org/10.1127/0029-5035/2011/0093-0277>
- Deshmukh UB, Rathor OS (2021) *Jirivanaea*, a new name for *Vitalia* P.E.A.S. Câmara, Carv.-Silva & W.R.Buck, hom. illeg. (Sematophyllaceae, Bryophyta) versus *Vitalia* Batista & Ciferri (Chaetothyriales, Ascomycota). Phytotaxa 500 (1): 055–056. <https://doi.org/10.11646/phytotaxa.500.1.9>
- Drumond MA, Kill LHP, Lima PCF, Oliveira MC, Oliveira VR, Albuquerque SG, Nascimento CES, Cavalcanti J (2000) Avaliação e identificação de ações prioritárias para a conservação, utilização sustentável e repartição de benefícios da biodiversidade do bioma Caatinga. <https://www.infoteca.cnptia.embrapa.br/infoteca/bitstream/doc/134000/1/usosustentavel.pdf>. Accessed on: 2022-04-20.
- Evangelista-dos-Santos M, Valente EB, Bastos CJP, Bôas-Bastos SBV (2019) Musgos (Bryophyta) da Estação Ecológica Wenceslau Guimarães, Bahia, Brasil. Hoehnea 46 (4): e092019. <https://doi.org/10.1590/2236-8906-09/2019>
- Evangelista-dos-Santos M, Almeida GSS (2019) Briófitas de um fragmento de floresta ombrófila densa, Fazenda Patioba, Alagoinhas, Bahia, Brasil. Pesquisas, Botânica 74: 325–336.
- Faria ALA, Valente DV, Silva AL, Cunha MJ, Amorim ET, Peralta DF (2020) Briófitas do Parque Estadual do Forno Grande, Espírito Santo – Mata Atlântica, Brasil. Pesquisas, Botânica 74: 283–302.
- Fhahm JP (1990) The effect of light and temperature on the growth of the bryophytes of tropical forest. Nova Hedwigia 51: 151–164.
- Florschutz PA (1964) The moss of Suriname. E.J. Brill, Leiden, 271 pp.
- Fiaschi P, Pirani JR (2009) Review of plant biogeographic studies in Brazil. Journal of Systematics and Evolution 47: 477–496. <https://doi.org/10.1111/j.1759-6831.2009.00046.x>
- Giuliette AM, Harely RM, Queiroz LP, Wanderley MGL, Berg CVD (2005) Biodiversity and conservation of plants in Brazil. Conservation Biology 19 (3): 632–639. <https://doi.org/10.1111/j.1523-1739.2005.00704.x>
- Gradstein SR, Pócs T (1989) Bryophytes. In: Lieth H, Wagner MJA (Eds.) Tropical rain Forest Ecosystems. Elsevier Science Publisher, Amsterdam, the Netherlands, 311–315.
- IBGE (Instituto Brasileiro de Geografia e Estatística IBGE) (2012) Manual técnico da vegetação brasileira, 2º ed. Série Manuais Técnicos em Geociências, Rio de Janeiro, Brazil, 271 pp.
- Koga ML, Peralta DF (2021) Bryophytes of Rio Turvo State Park (SP), Brazil: integrating floristics, geographical distribution, reproduction and ecological traits to support the conservation of the Atlantic Forest Fragment. Acta Botanica Brasiliensis 35 (3): 389–417. <https://doi.org/10.1590/0102-33062020abb0377>
- Moraes B, Razgour O, Souza-Alves JP, Boulli JP, Bezerra B (2020) Habitat suitability for primate conservation in north-east Brazil. Oryx 54 (6): 803–813. <https://doi.org/10.1017/S0030605319001388>
- Oliveira RR, Medeiros DL, Oliveira HC, Conceição GM (2018) Briófitas de área sob o domínio fitogeográfico do Cerrado e novas ocorrências para o Maranhão e região Nordeste do Brasil. Iheringia, Série Botânica 73 (2): 191–195. <https://doi.org/10.21826/2446-8231-201873211>
- Queiroz R, Cordeiro LS, Sampaio VS, Ribeiro RTM, Loiola MIB (2018) A região Nordeste. In: Coradin L, Camillo J, Pareyn FGC (Eds.). Espécies Nativas da Flora Brasileira de Valor Econômico Atual ou Potencial – Plantas para o futuro: Região Nordeste. Ministério do Meio Ambiente, Brasília, Brazil, 75–104.
- Ramsay HP, Schofield WB, Tan BC (2004) The family Sematophyllaceae (Bryopsida) in Australia, Part 2. *Acroporium*, *Clastobryum*, *Macrohymentium*, *Meiotheciella*, *Meiothecium*, *Papillidiopsis*, *Radulina*, *Rhaphidorrhynchium*, *Trichosteleum*, and *Warburgiella*. Journal of the Hattori Botanical Laboratory 95: 1–69.
- Remor D, Pastore JFP, Peralta DF (2021) Levantamento das briófitas da trilha do Passegueirinho, Curitibanos, estado de Santa Catarina, Brasil. Hoehnea 48: 1–14. <https://doi.org/10.1590/2236-8906-118/2020>
- Robins RG (1952) Bryophyte ecology of a dune area in New Zealand. Acta Geobotanica 4: 1–31.
- Sharp AJ, Crum H, Eckel PM (1994) The moss flora of Mexico. Memoirs of the New York Botanical Garden 69: 1–1113.
- Souza MEB, Oliveira HC (2019) Briófitas de uma área ecotonal de Caatinga/Cerrado no estado do Piauí, Brasil. Encyclopédia Biosfera 29 (16): 1796–1808.
- Species-Link (2022) Sematophyllaceae. <https://splink.cria.org.br/>. Accessed on 2022-04-10.
- Vanderpoorten A, Goffinet B (2009) Introduction to bryophytes. Cambridge University Press, New York, USA, 294 pp.
- Varão LF, Cunha IPR, Peralta DF (2011) Levantamento de briófitas do distrito Bananal, município Governador Edison Lobão, Maranhão, Brasil. Revistas de Biologia e Ciências da Terra 11 (2): 88–92.
- Visnadi RS (2006) Sematophyllaceae da Mata Atlântica do nordeste do Estado de São Paulo. Hoehnea 33 (4): 455–484.
- Yano O (1981) A checklist of Brazilian mosses. The Journal of the Hattori Botanical Laboratory 50: 279–456.
- Yano O (1984) Briófitas. In: Fidalgo O, Bononi VLR (Eds.). Técnicas de coleta, conservação e herborização do material botânico. Instituto de Botânica, São Paulo, Brazil, 27–30.
- Yano O (1989) An additional checklist of Brazilian bryophytes. The Journal of the Hattori Botanical Laboratory 66: 371–434.
- Yano O (1994) Briófitas da Serra de Itabaiana, Sergipe, Brasil. Acta Botanica Brasiliensis 8(1): 45–57.
- Yano O (2011) Catálogo de Musgos Brasileiros: literatura original, básiônimo, localidade-tipo e distribuição geográfica. Instituto de Botânica de São Paulo, São Paulo, Brazil, 180 pp.
- Yano O, Bordin J, Peralta DF (2009) Briófitas dos estados do Ceará, Maranhão, Paraíba, Piauí e Rio Grande do Norte (Brasil). Hoehnea 36 (3): 387–415.