
Shape, size, and polymer dependent effects of microplastics on *Daphnia magna*

Michael Schwarzer^{*1}, Julian Brehm², Martina Vollmer³, Julia Jasinski⁴, Chengzhang Xu⁵, Shakir Zainuddin⁶, Thomas Fröhlich⁷, Matthias Schott⁸, Andreas Greiner⁹, Thomas Scheibel¹⁰, and Christian Laforsch^{†11}

¹Michael Schwarzer – University of Bayreuth, Germany

²Julian Brehm – University of Bayreuth, Germany

³Martina Vollmer – University of Bayreuth, Germany

⁴Julia Jasinski – University of Bayreuth, Germany

⁵Chengzhang Xu – University of Bayreuth, Germany

⁶Shakir Zainuddin – University of Bayreuth, Germany

⁷Thomas Fröhlich – LMU Munich, Germany

⁸Matthias Schott – University of Bayreuth, Germany

⁹Andreas Greiner – University of Bayreuth, Germany

¹⁰Thomas Scheibel – University of Bayreuth, Germany

¹¹Christian Laforsch – University of Bayreuth, Germany

Abstract

The effects of microplastic (MP) pollution on organisms are gaining increasing attention. To date, a variety of polymers of different shapes and sizes are used in polystyrene (PS) is the predominant polymer type used in effect studies, it is still unclear whether the observed effects derive from the polymer itself or from a certain particle shape and size. To elucidate whether the effects are polymer specific, we conducted a systematic study on *Daphnia*

Keywords: Microplastics, Polymer dependent effect, *Daphnia*, Hazard assessment, Control particles

*Speaker

†Corresponding author: Christian.Laforsch@uni-bayreuth.de