

In the format provided by the authors and unedited.

Complex evolving patterns of mass loss from Antarctica's largest glacier

Jonathan L. Bamber^{ID}* and Geoffrey J. Dawson

Bristol Glaciology Centre, University of Bristol, Bristol, UK. *e-mail: j.bamber@bristol.ac.uk

Supplementary Information for

Complex, evolving patterns of mass loss from Antarctica's largest glacier

Jonathan L. Bamber*, Geoffrey Dawson
Bristol Glaciology Centre, University of Bristol, Bristol, UK

*Correspondence to: j.bamber@bristol.ac.uk

The supplementary Information contains one table.

Data	Temporal Coverage	Spatial resolution	Temporal resolution	Subset used	Satellite
MEaSUREs Annual Antarctic Ice Velocity Maps, Version 1.	2005-2017	1 km x 1 km	1 Year	N/A	ALOS, ENVISAT, LANDSAT-8, RADARSAT-1/2, SENTINEL-1 A/B, TDX, TSX.
Global Land Ice Velocity Extraction from Landsat 8 (GoLIVE).	2013/05/01 - 2017/04/30	300 m x 300 m	16 day	Path 1,229-233 Row 112-114	LANDSAT-8

Supplementary Table 1: Ice velocity products discussed in Section 1.4.