

Oreocharis polyneura, a new species of Gesneriaceae from southern Yunnan, China

Yan-Xiong Gong¹, Hong-Bo Ding^{2,3,4}, Xiang-Shuai Yan¹,
Fang Wen^{5,6}, Yao-Hua Tian¹, Yun-Hong Tan^{2,3,4}

1 Yunnan Institute of Tropical Crops, Jinghong, CN-666100, Yunnan, China **2** Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences, Menglun, Mengla, CN-666303, Yunnan, China **3** Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, Mengla, CN-666303, Yunnan, China **4** Center of Conservation Biology, Core Botanical Gardens, Chinese Academy of Sciences, Menglun, Mengla, CN-666303, Yunnan, China **5** Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin, CN-541006, Guangxi, China **6** Gesneriad Committee of China Wild Plant Conservation Association, National Gesneriaceae Germplasm Resources Bank of GXIB, Gesneriad Conservation Center of China, Guilin Botanical Garden, Chinese Academy of Sciences, Guilin, CN-541006, Guangxi, China

Corresponding authors: Yao-Hua Tian (tyhyhf@126.com), Yun-Hong Tan (tyh@xtbg.org.cn)

Academic editor: Eberhard Fischer | Received 23 August 2022 | Accepted 13 October 2022 | Published 22 November 2022

Citation: Gong Y-X, Ding H-B, Yan X-S, Wen F, Tian Y-H, Tan Y-H (2022) *Oreocharis polyneura*, a new species of Gesneriaceae from southern Yunnan, China. *PhytoKeys* 214: 7–15. <https://doi.org/10.3897/phytokeys.214.93901>

Abstract

Oreocharis polyneura, a new species from southern Yunnan, China, is described and illustrated. It is morphologically similar to *O. rhytidophylla* by having more obvious lateral veins forming a crosslinked network on the adaxial surface of leaf blades, but can be distinguished by having more lateral veins (12–15 pairs vs. 7–9 pairs) of leaf blades, shorter corolla tubes (the length ratio of corolla tube to corolla lobes = 1.2–2.9 vs. 3.4–6) and shorter pistils (6–8 mm long vs. 27–30 mm long). In addition, a detailed morphological description, a photographic illustration, the distribution and phenology of the new species are presented.

Keywords

flora of Yunnan, morphology, new taxon, *Oreocharis rhytidophylla*, taxonomy

Introduction

In recent years, the genus *Oreocharis* Benth. has been a hotspot for the research of the family Gesneriaceae. The newly circumscribed genus *Oreocharis* is recently enlarged

by incorporating ten other genera (*Ancylostemon* Craib, *Bournea* Oliv., *Dayaoshania* W.T.Wang, *Deinocheilos* W.T.Wang, *Isometrum* Craib, *Opithandra* B.L.Burtt, *Paraisometrum* W.T.Wang, *Thamnocharis* W.T.Wang, *Tremacron* Craib, *Briggsia* Craib s.str.) with high floral diversity (Möller et al. 2011; Weber et al. 2013), although *Bournea* was later reinstated as an independent genus (Chen et al. 2020a). Hitherto, *Oreocharis* s.l. contains ca. 150 taxa mainly distributed in southern and southwestern China (Wen et al. 2021), with a few of them (ca. 15 species) expanding to Bhutan, India, Japan, Myanmar, Thailand, and northern and central Vietnam (Möller et al. 2016, 2017, 2018; Cai et al. 2020a; Le et al. 2022). In China, southern Yunnan is rich in species diversity of the genus, with about 38 species recorded in this area (Li and Wang 2004; Möller et al. 2011; Chen et al. 2012, 2013, 2020b; Tan et al. 2013, 2015; Rossini and Freitas 2014; Cai et al. 2015, 2020; Cai and Dao 2020).

In 2021, during the expedition of the plant resources of tropical Yunnan in China, we collected an interesting plant of Gesneriaceae from Lancang county at anthesis (Figs 1, 2). Judging from the vegetative habit and floral characteristics, we considered it a member of *Oreocharis*. Upon careful comparisons of diagnostic morphological and anatomical features from the closely related species from China and adjacent regions, we determined that the plant is new to science and thus describe it herein. Its morphological characters are compared with the closely related species *Oreocharis rhytidophylla* C.Y. Wu ex H.W. Li (Li 1983; Zhang et al. 2019) (Fig. 3).

Materials and methods

Vouchers of the new species were collected from Lancang County, Yunnan Province, China. Photographs and phenology data were obtained during field expeditions. Morphological examinations and measurements of the new species were conducted on both living plants in the wild and herbarium specimens. All morphological characters are described according to the terminology presented by Wang et al. (1990, 1998). The preliminary conservation assessment for the new species was evaluated according to the guidelines of the IUCN Red List Categories and Criteria (IUCN 2019).

Taxonomic treatment

***Oreocharis polyneura* Y.H.Tan, F.Wen & Y.X.Gong, sp. nov.**

urn:lsid:ipni.org:names:77308755-1

Figs 1, 2

Diagnosis. *Oreocharis polyneura* is similar to *O. rhytidophylla* in having more obvious lateral veins forming crosslinked network on the adaxial surface of leaf blades, but can be distinguished by having more lateral veins (12–15 pairs vs. 7–9 pairs) of leaf blades, shorter corolla tubes (the length ratio of corolla tube to corolla lobes = 1.2–2.9 vs. 3.4–6) and shorter pistils (6–8 mm long vs. 27–30 mm long).

Type. CHINA. Yunnan Province, Puer City, Lancang County, Zhutang Town, on rocks or cliffs in evergreen broad-leaved forests in ravines, 22°45'44.28"N, 99°43'26.81"E, elev. 2137 m, 29 July 2021, *Y.X. Gong G344* (holotype: HITBC0079099; isotypes: HITBC, IBK). Fig. 2.

Description. Perennial herbs. *Rhizomes* subterete, short, straight, with numerous fibrous roots. *Leaves* basally forming a rosette, 4–8, petiolate; petioles terete, 1.2–7.0 cm long, 3–5 mm in diameter, densely rust-brown pannose, leaf blade thickly chartaceous, elliptic to ovate-elliptic, 4–13 × 2–7 cm, apex rounded, base rounded to subcordate, margin serrate to crenate, adaxially green, glabrous, abaxially whitish-green, densely rust-brown pannose along veins, lateral veins 12–15 on each side of the midrib, adaxially conspicuously sunken, abaxially conspicuously prominent, anastomosed forming on both surfaces. *Inflorescences* cymose, axillary, 2–4, 2–4-branched, 4–10-flowered; peduncles 4.5–9.0 cm long, ca. 1–3 mm in diameter, rust-brown lanate; bracts 3, verticillate, each 8.5–9.5 × 3.2–4.2 mm, lanceolate to elliptic, margins usually entire, occasionally 2–3-denticulate, adaxially glabrous, abaxially rust-brown pannose; pedicels 1.0–3.5 cm long, ca. 1 mm in diameter, sparsely puberulent. *Calyces* actinomorphic, 5-parted from the base, segments equal, linear-lanceolate, 7.5–9.0 × 1.5–2.5 mm, margin dentate, adaxially glabrous, abaxially rust-brown pannose. *Corollas* yellow, 1.8–2.2 cm long, outside glabrous, inside sparsely glandular-puberulent in the throat on adaxial lobes; tube narrowly infundibuliform, slightly bent near the throat, 1–1.5 cm long, 4–5.5 mm in diameter, proximally and distally almost equal in width; limb indistinctively bilabiate, adaxial lip 2-lobed, lobes obovate to elliptic, 5.8–6.2 × 3.8–4.8 mm, apex obtuse, margin entire, abaxial lip 3-lobed, lobes obovate to elliptic, almost equal, 6.2–8.2 × 5.2–6.2 mm, apex rounded, margin entire. *Stamens* 4, included, two pairs of stamens cohering at the anther tips, adnate to corolla tube 4–6 mm from base; adaxial stamens 3.5–4.5 mm long, abaxial stamens 5–6 mm long; filaments yellow, linear, glabrous; anthers basifixed, reniform, 2-locular, dehiscent transversely; staminode 1, 1.4 mm long, adnate to corolla tube 1.8 mm from the base. *Pistils* 6–8 mm long, glabrous; ovary 5–6.5 mm long, ca. 1.5 mm in diameter, glabrous, with two parietal placentae; style 1–1.5 mm long; stigma 1, disciform, retuse, edge flipped outwardly, 1.5–2.0 mm in diameter. *Discs* ring-shaped, ca. 2.3 mm high, margin slightly undulate with 5 irregularly and shallow lobes. *Capsules* indistinctively 4-angled to subterete, ca. 4 cm long, ca. 3 mm in diameter, glabrous.

Phenology. Flowering from July to October; fruiting from September to November.

Etymology. Greek *polys*, many, and *neuron*, nerve, alluding to abaxially conspicuous lateral leaf veins.

Habitat and distribution. Endemic to Zhutang town, Lancang county, Puer city, Yunnan Province, China (Fig. 4). It grows on rocks or cliffs under evergreen broad-leaved forests in ravines at elevations of 1900–2200 m.

Conservation status. *Oreocharis polyneura* is currently known only from its type locality. Based on our present knowledge and available data, its conservation status is assessed as “Data Deficient” (DD; IUCN 2019).




Figure 1. *Oreocharis polyneura* Y.H.Tan, F.Wen & Y.X.Gong, sp. nov. **A** habit **B** inflorescence **C** behind view of flower **D** fruit **E** whole plant **F** inflorescence with scale **G** adaxial and abaxial leaf surfaces **H** bracts **I, J** calyces **K** opened corolla showing stamens and staminode **L** pistil with disc and calyces **M** cross section of fruit. Photographed by Yan-Xiong Gong.



Figure 2. Type sheet of *Oreocharis polyneura* Y.H.Tan, F.Wen & Y.X.Gong, sp. nov.



Figure 3. Type sheet of *Oreocharis rhytidophylla* C.Y. Wu ex H.W. Li.

Vernacular name. The Chinese name of the new species is “Duō Mài Mǎ Líng Jù Tái” (多脉马铃薯苔). The first two words, Duō Mài mean the numerous veins of the leaf blades, and the rest four words mean the genus *Oreocharis* in mandarin.

Notes. Morphologically, *Oreocharis polyneura* is most similar to *O. rhytidophylla* in its leaves and inflorescences but is readily distinguishable by an array of characters (see Table 1).




Figure 4. Distribution of *Oreocharis polyneura* (black triangle) and *O. rhytidophylla* (black circle).

Table 1. Morphological comparison between *Oreocharis polyneura* and *O. rhytidophylla*.

Character	<i>Oreocharis polyneura</i>	<i>Oreocharis rhytidophylla</i>
Petiole	terete	plane
Leaf	lateral veins 12–15 pairs	lateral veins 7–9 pairs
Peduncle	4.5–9 cm long	up to 14 cm long
Corolla	1.8–2.2 cm long; tube narrowly infundibuliform	3.5–3.7 cm long; tube cylindrical
Length ratio of corolla tube to corolla lobes	1.2–2.9	3.4–6
Stamen	two pairs cohered at the anther tips, adnate to corolla tube 4–6 mm from base	adaxial stamens cohered in pair, abaxial stamens free, adnate to corolla tube 8–9 mm from base
Pistil	6.0–8.0 mm long; ovary 5.0–6.5 mm long	27–30 mm long; ovary 17–18 mm long

Acknowledgements

We are grateful to Dr. Eberhard Fischer and Dr. You-Pai Zeng for their valuable comments on the manuscript. We thank Shi-Wei Guo and Jian-Wu Li for providing scanned images of specimens. We also thank Bing-Fan Yang, Guo-Yun Chen and Yi-Gang Wei for their kind help in the fieldwork. This work was financially supported by Funds of Sci-Tech Innovation System Construction for Tropical crops Yunnan Province (RF2021-3), Xishuangbanna Rubber Ecosystem Research Station of Yunnan Province (202205AM070004), National Natural Science Foundation of China (31970223), the Project of the Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences (Y4ZK111B01), the Key Sci. & Tech. Research and Development Project of Guangxi (Guike AD20159091 & ZY21195050) and the capacity-building project of SBR of CAS (KFJ-BRP-017-68).

References

- Cai L, Chen RZ, Yin ZJ, Zhang GX, Chen WH, Shui YM (2015) *Tremacron hongheensis*, a new species of Gesneriaceae from Southeastern Yunnan, China. *Plant Diversity and Resources* 37(6): 733–736.
- Cai L, Dao ZL (2020) *Oreocharis argentifolia* (Gesneriaceae), a new species from the karst region in southeastern Yunnan, China. *Nordic Journal of Botany* 02699(8): njb.02699. <https://doi.org/10.1111/njb.02699>
- Cai L, Huang ZJ, Wen F, Dao ZL (2020) Two new species of *Oreocharis* (Gesneriaceae) from karst regions in Yunnan and notes on *O. tetraptera* and *O. brachypoda* from China. *PhytoKeys* 162: 1–12. <https://doi.org/10.3897/phytokeys.162.52174>
- Chen WH, Shui YM, Hua CL, Yu CY, Wen K (2012) *Ancylostemon dimorphosepalus* (Gesneriaceae), a new species from China. *Annales Botanici Fennici* 49(5–6): 391–394. <https://doi.org/10.5735/085.049.0612>
- Chen WH, Wang H, Shui YM, Möller M, Yu ZY (2013) *Oreocharis jinpingensis* (Gesneriaceae), a new species from Yunnan, China. *Annales Botanici Fennici* 50(5): 312–316. <https://doi.org/10.5735/086.050.0504>
- Chen WH, Zhang YM, Guo SW, Zhang ZR, Chen L, Shui YM (2020a) Reassessment of *Bournea* Oliver (Gesneriaceae) based on molecular and palynological evidence. *PhytoKeys* 157: 27–41. <https://doi.org/10.3897/phytokeys.157.55254>
- Chen WH, Zhang YM, He DM, Li YL, Shui YM (2020b) Four new species of *Oreocharis* (Gesneriaceae) in Yunnan province, China. *PhytoKeys* 157: 83–99. <https://doi.org/10.3897/phytokeys.157.32284>
- IUCN [Standards and Petitions Subcommittee] (2019) Guidelines for Using the IUCN Red List Categories and Criteria, Version 14. Prepared by the Standards and Petitions Subcommittee. <https://www.iucnredlist.org/resources/redlistguidelines> [Accessed 8 September 2021]
- Le KD, Nguyen TT, Nguyen PT, Hoang TT, Wen F, Do TV (2022) *Oreocharis phuongii* (Gesneriaceae), a new species from central Vietnam. *PhytoKeys* 193: 43–53. <https://doi.org/10.3897/phytokeys.193.77083>

- Li HW (1983) Notulae De Gesneraceis Yunnanensibus. Bulletin of Botanical Research 3(2): 1–55.
- Li ZY, Wang YZ (2004) Plants of Gesneriaceae in China. Henan Science & Technology Publishing House, Zhengzhou, 14–47.
- Möller M, Atkins HJ, Bramley GLC, Middleton DJ, Baines R, Nguyen VD, Bui HQ, Barber S (2018) Two new species of *Oreocharis* (Gesneriaceae) from northern Vietnam. Edinburgh Journal of Botany 75(3): 309–319. <https://doi.org/10.1017/S0960428618000148>
- Möller M, Middleton D, Nishii K, Wei YG, Sontag S, Weber A (2011) A new delineation for *Oreocharis* incorporating an additional ten genera of Chinese Gesneriaceae. Phytotaxa 23(1): 1–36. <https://doi.org/10.11646/phytotaxa.23.1.1>
- Möller M, Nampy S, Janeesha AP, Weber A (2017) The Gesneriaceae of India: Consequences of updated generic concepts and new family classification. Rheedia 27(1): 23–41. <https://doi.org/10.22244/rheedia.2017.27.1.5>
- Möller M, Wei YG, Wen F, Clark JL, Weber A (2016) You win some you lose some: Updated generic delineations and classification of Gesneriaceae - implications for the family in China. Guihaia 36(1): 44–60.
- Rossini J, Freitas J (2014) *Oreocharis yunnanensis*, a new name for the illegitimate *Oreocharis glandulosa* (Gesneriaceae) from China. Phytotaxa 163(3): 180–180. <https://doi.org/10.11646/phytotaxa.163.3.5>
- Tan YH, Li JW, Pan B, Wen B, Yin JT, Liu Q (2013) *Oreocharis glandulosa*, a new species of Gesneriaceae from southern Yunnan, China. Phytotaxa 131(1): 29–34. <https://doi.org/10.11646/phytotaxa.131.1.5>
- Tan YH, Li JW, Yin JT (2015) *Oreocharis tsaii*, a new species of Gesneriaceae from southern Yunnan, China. Phytotaxa 195(2): 188–192. <https://doi.org/10.11646/phytotaxa.195.2.9>
- Wang WT, Pan KY, Zhang ZY, Li ZY (1990) Gesneriaceae. In: Wang WT (Ed.) Flora Reipublicae Popularis Sinicae Vol. 69. Science Press, Beijing, 141–271.
- Wang WT, Pan KY, Li ZY, Weitzman AL, Skog LE (1998) Gesneriaceae. In: Wu ZY, Raven PH (Eds) Flora of China Vol. 18. Scrophulariaceae through Gesneriaceae. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis, 244–401.
- Weber A, Clark JL, Möller M (2013) A new formal classification of Gesneriaceae. Selbyana 31: 68–94.
- Wen F, Wei YG, Fu LF, Xin, ZB, Ge YZ (2021) The checklist of Gesneriaceae in China. <http://gccx.gxib.cn/cn/about-68.aspx> [Accessed 8 July 2021]
- Zhang YM, Guo SW, Chen WH, Shui YM (2019) Rediscovery and conformation of *Oreocharis rhytidophylla* (Gesneriaceae) with supplementary description of flowers. Guihaia 39: 569–573.