

Supplementary Appendix

This appendix has been provided by the authors to give readers additional information about their work.

Supplement to: Walter EB, Talaat KR, Sabharwal C, et al. Evaluation of the BNT162b2 Covid-19 vaccine in children 5 to 11 years of age. *N Engl J Med* 2022;386:35-46. DOI: 10.1056/NEJMoa2116298

SUPPLEMENTARY APPENDIX

List of Investigators	2
Inclusion and Exclusion Criteria.....	5
Ethical Conduct of the Study	5
Study Responsibilities.....	5
Phase 1 Review of Safety Data and Stopping Rules.....	6
Calculation of Immunogenicity Parameters.....	6
Determination of COVID-19 and MIS-C	7
Figure S1. Local reactions (A) and systemic events (B) reported in the phase 1 study in participants 5–11 years old.....	9
Figure S2. Immunogenicity of BNT162b2 at 7 days after dose 2 in the phase 1 study in participants 5–11 years old.....	11
Figure S3. Geometric mean titers and geometric mean fold rises of SARS-CoV-2 neutralizing titers for participants 5–11 and 16–25 years old.....	12
Table S1. Study populations	13
Table S2. Demographic characteristics of phase 1 participants 5–11 years old*	14
Table S3. Demographic characteristics of phase 2/3 participants 5–11 and 16–25 years old*	15
Table S4. Duration from first to last day of local reactions and systemic events	16
Table S5. Severity scale for local reactions and systemic events	18
Table S6. Phase 1 participants 5–11 years old reporting at least 1 adverse event	19
Table S7. Phase 2/3 participants 5–11 years old reporting at least 1 adverse event	20

List of Investigators

Name	Institute	Location
Acevedo, Armando	Acevedo Clinical Research Associates	Miami, FL, USA
Amisola, Rogelio	PriMed Clinical Research	Dayton, OH, USA
Anderson, Evan	Emory Children's Center Illness POD Emory University School of Medicine	Atlanta, GA, USA Atlanta, GA, USA
Arimany Montaña, Josep Lluís	Hospital Universitari General de Catalunya	Barcelona, Spain
Arora, Samir	Aventiv Research	Columbus, OH, USA
Baca Cots, Manuel	Grupo Pediátrico Uncibay	Málaga, Spain
Baker, Jeffrey	Clinical Research Prime	Idaho Falls, ID, USA
Barnett, Elizabeth	Boston Medical Center	Boston, MA, USA
Benun, Jacques	ARC Clinical Research at Wilson Parke	Austin, TX, USA
Boppana, Suresh	University of Alabama at Birmingham – School of Medicine Pediatrics	Birmingham, AL, USA
Butuk, David	Solaris Clinical Research	Meridian, ID, USA
Caserta, Mary	University of Rochester Medical Center	Rochester, NY, USA
Casey, Janet	Rochester Clinical Research	Rochester, NY, USA
Christensen, Shane	J. Lewis Research – Foothill Family Clinic South	Salt Lake City, UT, USA
Czajka, Hanna	Centrum Badań Klinicznych JCI	Krakow, Poland
Day, James Christopher (replaced Barbara Pahud)	Children's Mercy Hospital	Kansas City, MO, USA
De Valle, Oscar	West Houston Clinical Research Service	Houston, TX, USA
Dever, Michael	Clinical Neuroscience Solutions	Orlando, FL, USA
Domachowske, Joseph	State University of New York, Upstate University Hospital – Pediatric and Adolescent Infectious Disease/Immunology Center	Syracuse, NY, USA
Eder, Frank	Meridian Clinical Research	Binghamton, NY, USA
Englund, Janet	Seattle Children's Hospital	Seattle, WA, USA
Fergie, Jaime	Driscoll Children's Hospital	Corpus Christi, TX, USA
Finn, Daniel	Kentucky Pediatric/Adult Research	Bardstown, KY, USA
Fortuny Guasch, Clàudia	Hospital Sant Joan de Déu	Barcelona, Spain
Garcia-Diaz, Julia	Ochsner Clinic Foundation	New Orleans, LA, USA
Garcia-Sicilia Lopez, Jose	Hospital Universitario HM Montepríncipe	Madrid, Spain
Garza, Homero	Van Tran Family Practice Ventavia Research Group	Houston, TX, USA Houston, TX, USA
Hartman, Aaron	Virginia Research Center	Midlothian, VA, USA

Heller, Robert	Bayview Research Group	Valley Village, CA, USA
Jennings, Timothy	Clinical Research Professionals	Chesterfield, MO, USA
Kantele, Anu	Meilahti Vaccine Research Center (MeVac)	Helsinki, Finland
Karhusaari, Hanna	Tampere University/Pori Vaccine Research Clinic	Pori, Finland
Klein, Nicola	Kaiser Permanente Northern California	Oakland, CA, USA Sacramento, CA, USA Santa Clara, CA, USA
Klein, Terry	Alliance for Multispecialty Research	Wichita, KS, USA
Kokko, Satu	Tampere University/Oulu Vaccine Research Clinic	Oulu, Finland
Kopinska, Elzbieta	Nasz Lekarz Ośrodek Badań Klinicznych	Torun, Poland
Korbal, Piotr	IN VIVO	Bydgoszcz, Poland
Koski, Susanna	Tampere University/Helsinki South Vaccine Research Clinic	Helsinki, Finland
Kuchar, Ernest	Provita Poliklinika	Warszawa, Poland
Kulig, Karolina	Ośrodek Badań Klinicznych AppleTreeClinics	Lodz, Poland
Laajalahti, Outi	Tampere University/Seinäjoki Vaccine Research Clinic	Seinäjoki, Finland
Leonard, Daniel	Meridian Clinical Research	Hastings, NE, USA
Li, Simon	Rutgers University – Pediatric Research Center	New Brunswick, NJ, USA
Majda-Stanisławska, Ewa	GRAVITA Diagnostyka i Leczenie Nieplodności	Lodz, Poland
Maldonado, Yvonne	Stanford University – Clinical and Translational Research Unit	Palo Alto, CA, USA
Marshall, Gary	Novak Center for Children’s Health	Louisville, KY, USA
Martinón Torres, Federico	Hospital Clínico Universitario de Santiago de Compostela	Santiago de Compostela, Spain
Miranda Valdivieso, Mariano	Hospital de Antequera	Antequera, Spain
Munoz, Flor M.	Texas Children’s Hospital, Baylor College of Medicine	Houston, TX, USA
Nachman, Sharon	Advanced Specialty Care Clinical Research Center Stony Brook University	Commack, NY, USA East Setauket, NY, USA Stony Brook, NY, USA
Natalini Martínez, Silvina	Hospital HM Puerta del Sur	Móstoles, Spain
Ogbuagu, Onyema	Yale Center for Clinical Investigation	New Haven, CT, USA
Ohnmacht, Richard	Velocity Clinical Research – Providence	Warwick, RI, USA
Paavola, Pauliina	Tampere University/Kokkola Vaccine Research Clinic	Kokkola, Finland
Pajek, Barbara	Niepubliczny Zakład Lecznictwa Ambulatoryjnego Michalkowice, Jarosz i Partnerzy Spolka Lekarska	Siemianowice Śląskie, Poland
Patel, Nehali	St. Jude Children’s Research Hospital	Memphis, TN, USA
Paulsen, Grant	Cincinnati Children’s Hospital Medical Center	Cincinnati, OH, USA
Peltonen, Lauri	Tampere University/Järvenpää Vaccine Research Center	Järvenpää, Finland

Peterson, James	J. Lewis Research – Foothill Family Clinic North	Salt Lake City, UT, USA
Rey, M. Mildred	Center for Clinical Trials	Paramount, CA, USA
Riesenberg, Robert	Atlanta Center for Medical Research	Atlanta, GA, USA
Rojo, Pablo	Hospital Universitario 12 de Octubre	Madrid, Spain
Salamanca de la Cueva, Ignacio	Instituto Hispalense de Pediatría	Sevilla, Spain
Senders, Shelly	Senders Pediatrics	South Euclid, OH, USA
Seppa, Ilkka	Turku Vaccine Research Clinic	Turku, Finland
Sharp, Stephan	Clinical Research Associates	Nashville, TN, USA
Sher, Lawrence	Peninsula Research Associates	Rolling Hills Estates, CA, USA
Simões, Eric A.F.	Children’s Hospital Colorado	Aurora, CO, USA
Simonsen, Kari	Children’s Hospital & Medical Center Children’s Physicians Clinic, Spring Valley	Omaha, NE, USA Omaha, NE, USA
Sipila, Marjaana	Tampere Vaccine Research Clinic	Tampere, Finland
Slade, Lance	Meridian Clinical Research	Macon, GA, USA
Slechta, Stacy	Alliance for Multispecialty Research	Newton, KS, USA
Stryczyńska-Kazubska, Joanna	Rodzinne Centrum Medyczne Lubmed	Luboń, Poland
Talaat, Kawsar	Johns Hopkins University	Baltimore, MD, USA
Ukkonen, Benita	Tampere University/Espoo Vaccine Research Clinic	Espoo, Finland
Ulied Armiñana, Angels	EAP Centelles	Centelles, Spain
Vanchiere, John	Louisiana State University Health Shreveport – Clinical Trials Office	Shreveport, LA, USA
Walter, Emmanuel	Clinical Research Pickett Road Duke Human Vaccine Institute	Durham, NC, USA Durham, NC, USA
Wiedermann, Bernhard	Children’s National Medical Center	Washington, DC, USA
Wisman, Paul	Pediatric Associates of Charlottesville Pediatric Research of Charlottesville	Charlottesville, VA, USA
Zomcik, Anne	Allegheny Health and Wellness Pavilion	Erie, PA, USA

Inclusion and Exclusion Criteria

The study included healthy participants with pre-existing stable disease (ie, disease not requiring a significant change in therapy or hospitalization for worsening disease in the 6 weeks before enrollment). In the phase 1 portion of the study, exclusion criteria included past clinical (based on COVID-19 symptoms/signs alone if a SARS-CoV-2 nucleic acid amplification test [NAAT] result was unavailable) or virologic (based on COVID-19 symptoms/signs and a positive SARS-CoV-2 NAAT result) COVID-19 diagnosis, as well as known HIV or hepatitis C or B virus infection. Exclusion criteria in the phase 1 and 2/3 portions of the study included receipt of COVID-19 preventative treatments, history of multisystem inflammatory syndrome in children (MIS-C) or of severe adverse reaction with a vaccine or any component of the study intervention, and immunodeficiency, autoimmune disease, or conditions associated with prolonged bleeding.

Ethical Conduct of the Study

Conduct of this study was in accordance with the study protocol and with consensus ethical principles derived from international guidelines including the Declaration of Helsinki, Council for International Organizations of Medical Sciences Ethical Guidelines, International Conference on Harmonisation Guidelines for Good Clinical Practice, and applicable laws and regulations, including applicable privacy laws. The study protocol and any amendments, informed consent documents, and other relevant documents were approved by institutional review board/ethics committees (IRB/EC) before study initiation. Before any study activity, written informed consent was obtained from the participants' parents/legal guardians, and affirmative agreement from the participant was subsequently obtained when in the capacity to provide assent, as determined by the IRB/EC.

Study Responsibilities

Pfizer was responsible for study design and conduct; data collection, analysis, and interpretation; and writing of this manuscript. Both Pfizer and BioNTech manufactured the vaccine and placebo. BioNTech was the regulatory sponsor of the study and contributed to data interpretation and writing of the manuscript. All data were available to the authors, who vouch for its accuracy and completeness and to adherence of the study to the protocol.

Phase 1 Review of Safety Data and Stopping Rules

An Internal Review Committee (IRC) reviewed safety and immunogenicity data to permit dose level finding in each age group, and to select the dose level to proceed to phase 2/3. Data supporting these selections included safety, binding antibody levels, and neutralizing titers.

Stopping rules were applied to all phase 1 participants based on review of adverse event (AE) and reactogenicity data, until the start of phase 2/3 or 30 days after the last vaccine dose in phase 1. These data were continuously monitored by the investigators and sponsor to allow prompt identification of any event contributing to the stopping rule.

The stopping rule criteria for each BNT162b2 dose level included: (1) if a participant developed a serious AE as assessed by the investigator as possibly related to the vaccine and for which there was no alternative, plausible, or attributable cause; (2) if a participant developed a grade 4 local reaction or systemic event; (3) if any participant developed a fever $>40^{\circ}\text{C}$ for ≥ 1 daily measurement after vaccination; (4) if 2 participants within the same age group reported the same or similar severe AEs after vaccination as assessed by the investigator as possibly related to the vaccine and for which there was no alternative, plausible, or attributable cause; and (5) if any participant died or required intensive care unit admission because of SARS-CoV-2 infection.

If a stopping rule was met, the IRC was to consider all data; randomization and vaccine administration for all dose levels in the affected age group were to be paused; and the Data Monitoring Committee was to review all appropriate data. All other routine study conduct activities were to continue for all vaccinated participants.

Calculation of Immunogenicity Parameters

The geometric mean titers (GMTs) were derived by exponentiating the mean, and geometric mean ratios were derived by exponentiating the mean of the differences (5–11-year-olds minus 16–25-year-olds), in logarithmically transformed neutralizing titer values. Associated 2-sided 95% CIs were based on the Student *t* distribution and then exponentiating results. Geometric mean fold rises (GMFRs) were calculated for participants with non-missing values both before and after each vaccination by exponentiating the mean of the difference of logarithmically transformed assay results. Associated 2-sided 95% CIs were obtained using the Student *t* distribution for the mean difference and exponentiating the confidence limits.

Determination of COVID-19 and MIS-C

Surveillance for potential cases of COVID-19 and MIS-C will occur throughout a participant's involvement in the study. If a participant developed an acute illness, the participant was considered to potentially have COVID-19 illness. In this circumstance, assessments were to include nasal (anterior nares) swab sample collection, which were tested at a central laboratory using the reverse transcription–polymerase chain reaction (RT-PCR) test (Cepheid) or other equivalent nucleic acid amplification–based test (NAAT), to detect SARS-CoV-2. In addition, clinical information and results from local standard-of-care tests were assessed. The central laboratory NAAT result was used for the case definition. If no result was available from the central laboratory, a local NAAT result could be used if it was obtained using either the Cepheid Xpert Xpress SARS-CoV-2, Roche Cobas SARS-CoV-2 Real-Time RT-PCR test, or the Abbott Molecular/RealTime SARS-CoV-2 assay.

SARS-CoV-2–related cases, and SARS-CoV-2–related severe cases were documented (for both, the onset date of the case was the date that symptoms were first experienced by the participant; if new symptoms were reported within 4 days after resolution of all previous symptoms, they were considered as part of a single illness). The definition of confirmed COVID-19 included the presence of ≥ 1 symptom (ie, fever, new or increased cough, new or increased shortness of breath, chills, new or increased muscle pain, new loss of taste or smell, sore throat, diarrhea [≥ 3 stools per day], vomiting) and being SARS-CoV-2 NAAT-positive during, or within 4 days before or after, the symptomatic period (either at the central laboratory or at a local testing facility using an acceptable test).

Diagnosis of severe COVID-19 included confirmed COVID-19 and the presence of ≥ 1 of the following:

- (1) clinical signs at rest indicative of severe systemic illness (eg, respiratory rate and heart rate as shown in the following table, $SpO_2 \leq 92\%$ on room air or $>50\%$ FiO_2 to maintain $\geq 92\%$, or $PaO_2/FiO_2 < 300$ mmHg)

Participant age	Respiratory rate	Heart rate
4–6 years	>29	>131
6–<8 years	>27	>123
8–<12 years	>25	>115

- (2) respiratory failure (ie, needing high-flow oxygen, including continuous positive airway pressure, bilevel positive airway pressure, noninvasive ventilation, mechanical ventilation, or extracorporeal membrane oxygenation)
- (3) evidence of shock or cardiac failure (ie, systemic blood pressure $<70 + (\text{age in years} \times 2)$ for age up to 10 years, <90 for age ≥ 10 years; or requiring vasoactive drugs to maintain blood pressure in the normal range)


- (4) significant acute renal failure (ie, serum creatinine $\geq 2 \times$ upper limit of normal [ULN] for age or 2-fold increase in baseline creatinine)
- (5) significant gastrointestinal/hepatic failure (ie, total bilirubin ≥ 4 mg/dL or ALT $2 \times$ ULN for age)
- (6) significant neurological dysfunction (ie, Glasgow Coma Scale score ≤ 11 or acute change in mental status with a decrease in Glasgow Coma Scale score ≥ 3 points from abnormal baseline)
- (7) intensive care unit admission
- (8) death

Confirmed MIS-C was per the CDC MIS-C case definition (www.cdc.gov/mis-c/hcp/).

Figure S1. Local reactions (A) and systemic events (B) reported in the phase 1 study in participants 5–11 years old

Local reactions and systemic events were assessed within 7 days after administration of BNT162b2 at a 10- μ g, 20- μ g, or 30- μ g dose level. Of 16 participants who received 30 μ g BNT162b2 at dose 1, 4 received 30 μ g BNT162b2 at dose 2, and 12 received 10 μ g BNT162b2 at dose 2. Severity scales for reactogenicity events are summarized in **Table S5**. Fever categories are designated in the key. The numbers above the bars show the number of participants in each group with the specified local reaction or systemic event. Error bars are the 95% CIs.

10 μ g=10 μ g BNT162b2; 20 μ g=20 μ g BNT162b2; 30/30 μ g=BNT162b2 administered at a 30- μ g dose level at dose 1 and dose 2; 30/10 μ g=BNT162b2 administered at a 30- μ g dose level at dose 1 and at a 10- μ g dose level at dose 2.



B




Figure S2. Immunogenicity of BNT162b2 at 7 days after dose 2 in the phase 1 study in participants 5–11 years old

Results are for participants in the evaluable immunogenicity population (**Table S1**) who had no serological or virological evidence of SARS-CoV-2 infection (before the 7-day post-dose 2 blood collection) and no medical history of COVID-19. Dots represent individual titers. GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titers and the corresponding CIs (based on the Student *t* distribution). No assay results were below the lower limit of quantitation.

GMT=geometric mean titer; SARS-CoV-2=severe acute respiratory syndrome coronavirus 2.




Figure S3. Geometric mean titers and geometric mean fold rises of SARS-CoV-2 neutralizing titers for participants 5–11 and 16–25 years old

Results are for participants in the immunobridging subset of the evaluable immunogenicity population (**Table S1**) who had no serological or virological evidence of SARS-CoV-2 infection before the 1-month post-dose 2 blood sample collection, and who had no COVID-19 medical history. GMTs, GMFRs, and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titers or fold rises and the corresponding CIs (based on the Student *t* distribution). Assay results below the LLOQ were set to $0.5 \times$ LLOQ.

GMFR=geometric mean fold rise; GMT=geometric mean titer; LLOQ=lower limit of quantitation; SARS-CoV-2=severe acute respiratory syndrome coronavirus 2.




Table S1. Study populations

Population	Description
Enrolled	All participants who provided informed consent
Randomized	All participants who were assigned a randomization number
Immunobridging subset	A subset of participants 5–11 years old from this study (approximately the first 6 participants from every site) and randomly selected participants 16–25 years old from the ongoing pivotal study selected for immunogenicity analyses.
Evaluable immunogenicity	All eligible randomized participants who received 2 vaccine doses to which they were randomized (with dose 2 received within the predefined window), had ≥ 1 valid and determinate immunogenicity result from the blood sample collected within an appropriate window, and had no important protocol deviations as determined by the clinician
All-available immunogenicity	All randomized participants who received ≥ 1 dose of the study intervention and had ≥ 1 valid and determinate immunogenicity result after vaccination
Safety	All randomized participants who received ≥ 1 dose of the study intervention
Evaluable efficacy	All eligible randomized participants who received 2 vaccine doses to which they were randomized within the predefined window, and who had no other important protocol deviations as determined by the investigator
All-available efficacy	Dose 1: all randomized participants who received at least 1 dose Dose 2: all randomized participants who received both doses

Table S2. Demographic characteristics of phase 1 participants 5–11 years old*

Characteristic	BNT162b2 10 µg (N=16)	BNT162b2 20 µg (N=16)	BNT162b2 30/30 µg† (N=4)	BNT162b2 30/10 µg† (N=12)	Total (N=48)
Male, n (%)	5 (31.3)	10 (62.5)	3 (75.0)	6 (50.0)	24 (50.0)
Race, n (%)‡					
White	11 (68.8)	13 (81.3)	4 (100.0)	10 (83.3)	38 (79.2)
Black or African American	3 (18.8)	0	0	0	3 (6.3)
Asian	2 (12.5)	2 (12.5)	0	1 (8.3)	5 (10.4)
Multiracial, American Indian or Alaska Native	0	1 (6.3)	0	1 (8.3)	2 (4.2)
Ethnicity, n (%)‡					
Hispanic/Latinx	2 (12.5)	0	2 (50.0)	0	4 (8.3)
Non-Hispanic/non-Latinx	14 (87.5)	16 (100.0)	2 (50.0)	12 (100.0)	44 (91.7)
Age, years					
Mean, ± standard deviation	7.9±1.89	8.0±1.97	6.8±2.36	8.3±1.76	7.9±1.90
Median (range)	8.5 (5–11)	8.5 (5–10)	6.0 (5–10)	8.5 (5–10)	8.0 (5–11)
Country, n (%)					
United States	16 (100.0)	16 (100.0)	4 (100.0)	12 (100.0)	48 (100.0)

* Results are for the safety population (**Table S1**).

† Of 16 participants who received 30 µg at dose 1, 4 received 30 µg at dose 2 and 12 received 10 µg at dose 2.

Percentages may not total 100 because of rounding.

‡ Race and ethnic group were reported by the participants or their parents/guardians.

Table S3. Demographic characteristics of phase 2/3 participants 5–11 and 16–25 years old*

Characteristic	BNT162b2		Placebo	
	10 µg 5–11 years old (N=311)	30 µg 16–25 years old† (N=286)	5–11 years old (N=156)	16–25 years old‡ (N=49)
Male, n (%)	161 (51.8)	141 (49.3)	89 (57.1)	18 (36.7)
Race, n (%)§				
White	247 (79.4)	217 (75.9)	127 (81.4)	31 (63.3)
Black or African American	20 (6.4)	33 (11.5)	6 (3.8)	12 (24.5)
Asian	23 (7.4)	17 (5.9)	15 (9.6)	3 (6.1)
Multiracial	17 (5.5)	12 (4.2)	6 (3.8)	1 (2.0)
Other/not reported	4 (1.3)	7 (2.4)	2 (1.3)	2 (4.1)
Ethnicity, n (%)§				
Hispanic/Latinx	55 (17.7)	102 (35.7)	27 (17.3)	14 (28.6)
Age at vaccination, years				
Mean, ± standard deviation	8.3±1.88	20.9±3.03	8.2±2.06	20.8±3.05
Median (range)	8.0 (5–11)	21.0 (16–25)	9.0 (5–11)	22.0 (16–25)
Obese, n (%)§	30 (9.6)	48 (16.8)	20 (12.8)	15 (30.6)
Baseline SARS-CoV-2 status**				
Positive	24 (7.7)	14 (4.9)	14 (9.0)	1 (2.0)
Negative	287 (92.3)	271 (94.8)	142 (91.0)	48 (98.0)
Missing	0	1 (0.3)	0	0

* Results are for the all-available immunogenicity population of the immunobridging subset (**Table S1**).

† Randomly selected from participants in all countries in the ongoing pivotal trial.

‡ Race and ethnic group were reported by the participants or their parents/guardians.

§ Body mass index ≥95th percentile according to the growth chart

(https://www.cdc.gov/growthcharts/html_charts/bmiagerev.htm) for 5–11-year-olds, or ≥30 kg/m² for 16–25-year-olds.

** A positive severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) status required a positive N-binding antibody result at vaccination visit 1, a positive nucleic acid amplification test result at vaccination visit 1, or a medical history of coronavirus disease 2019.

Table S4. Duration from first to last day of local reactions and systemic events

Phase 1				
Local reaction or systemic event	BNT162b2 10 µg	BNT162b2 20 µg	BNT162b2 30/30 µg*	BNT162b2 30/10 µg*
Dose 1, median (range), days [n]				
Redness	3.5 (2, 5) [2]	NE [0]	2.0 (1, 4) [4]	3.5 (3, 4) [2]
Swelling	1.0 (1, 1) [3]	1.0 (1, 1) [1]	1.0 (1, 1) [2]	2.0 (2, 2) [1]
Injection site pain	2.0 (1, 5) [14]	2.0 (1, 6) [15]	3.0 (2, 5) [4]	2.0 (1, 5) [10]
Fever	1.0 (1, 1) [1]	1.0 (1, 1) [1]	NE [0]	1.0 (1, 1) [4]
Fatigue	2.0 (1, 6) [8]	2.0 (1, 6) [11]	1.0 (1, 5) [4]	1.5 (1, 3) [6]
Headache	1.5 (1, 5) [4]	2.0 (1, 5) [5]	2.0 (2, 5) [3]	1.0 (1, 1) [4]
Chills	NE [0]	1.0 (1, 5) [4]	2.5 (1, 4) [2]	1.0 (1, 1) [2]
Vomiting	NE [0]	1.0 (1, 1) [1]	NE [0]	1.0 (1, 1) [1]
Diarrhea	1.0 (1, 1) [1]	1.0 (1, 1) [1]	NE [0]	NE [0]
Muscle pain	1.5 (1, 2) [2]	1.0 (1, 3) [4]	1.0 (1, 4) [4]	NE [0]
Joint pain	1.0 (1, 1) [1]	1.0 (1, 1) [1]	3.0 (3, 3) [1]	1.0 (1, 1) [1]
Dose 2, median (range), days [n]				
Redness	1.0 (1, 2) [6]	1.0 (1, 1) [3]	3.0 (2, 3) [3]	1.0 (1, 1) [2]
Swelling	2.0 (1, 3) [5]	1.0 (1, 5) [3]	1.0 (1, 1) [2]	NE [0]
Injection site pain	2.0 (1, 5) [14]	2.0 (1, 6) [12]	2.5 (2, 4) [4]	2.0 (1, 3) [11]
Fever	1.0 (1, 1) [2]	1.0 (1, 1) [3]	1.0 (1, 1) [4]	NE [0]
Fatigue	1.0 (1, 6) [11]	1.0 (1, 5) [10]	1.0 (1, 9) [4]	1.0 (1, 4) [9]
Headache	1.0 (1, 5) [8]	2.0 (1, 6) [9]	1.0 (1, 4) [3]	1.0 (1, 1) [4]
Chills	1.0 (1, 4) [5]	1.0 (1, 3) [7]	1.0 (1, 2) [3]	1.0 (1, 1) [4]
Vomiting	NE [0]	NE [0]	1.0 (1, 1) [1]	1.0 (1, 1) [1]
Diarrhea	1.0 (1, 1) [1]	NE [0]	1.0 (1, 1) [2]	NE [0]
Muscle pain	NE [0]	1.0 (1, 6) [3]	5.0 (1, 9) [2]	1.0 (1, 1) [1]
Joint pain	NE [0]	NE [0]	9.0 (9, 9) [1]	NE [0]
Phase 2/3				
Dose 1, median (range), days [n[†]]	BNT162b2 10 µg		Placebo	
Redness	1.0 (1, 8) [176]		1.0 (1, 10) [89]	
Swelling	1.0 (1, 9) [125]		1.0 (1, 6) [53]	
Injection site pain	2.0 (1, 10) [915]		2.0 (1, 10) [437]	
Fever	1.0 (1, 3) [29]		1.0 (1, 3) [19]	
Fatigue	1.0 (1, 21) [501]		1.0 (1, 10) [241]	
Headache	1.0 (1, 22) [359]		1.0 (1, 19) [160]	
Chills	1.0 (1, 9) [77]		1.0 (1, 10) [28]	
Vomiting	1.0 (1, 3) [26]		1.0 (1, 5) [18]	
Diarrhea	1.0 (1, 6) [85]		1.0 (1, 8) [35]	
Muscle pain	1.0 (1, 8) [133]		1.0 (1, 9) [55]	
Joint pain	1.0 (1, 7) [60]		1.0 (1, 5) [31]	

Dose 2, median (range), days [n[†]]		
Redness	2.0 (1, 11) [211]	1.0 (1, 8) [105]
Swelling	2.0 (1, 12) [164]	2.0 (1, 8) [83]
Injection site pain	2.0 (1, 12) [849]	2.0 (1, 9) [431]
Fever	1.0 (1, 5) [79]	1.0 (1, 3) [28]
Fatigue	1.0 (1, 12) [527]	1.0 (1, 14) [243]
Headache	1.0 (1, 51) [379]	1.0 (1, 9) [179]
Chills	1.0 (1, 8) [131]	1.0 (1, 8) [48]
Vomiting	1.0 (1, 5) [17]	1.0 (1, 2) [17]
Diarrhea	1.0 (1, 28) [88]	1.0 (1, 9) [26]
Muscle pain	1.0 (1, 9) [155]	1.0 (1, 8) [75]
Joint pain	1.0 (1, 7) [73]	1.0 (1, 18) [33]

n=number of participants reporting the specified reaction or event on any of the 7 days after dose 1 or dose 2;

NE=not estimable.

Results are for the safety population (**Table S1**).

Duration was calculated in days as the difference from the start of the first reported reaction or event to the resolution of the last reported reaction or event, inclusive. For symptoms that were ongoing at the time of the next dose, the stop date was computed as the next dose date. Reactions and events were recorded in the electronic diary and unscheduled clinical assessments from Day 1 through Day 7 after each dose. The resolution date for reactions lasting longer than 7 days was recorded on the participant's case report form.

* Of 16 participants who received 30 µg at dose 1, 4 received 30 µg at dose 2 and 12 received 10 µg at dose 2.

† Includes participants with reactions or events of unknown duration.

Table S5. Severity scale for local reactions and systemic events

	Mild	Moderate	Severe	Grade 4
Local reaction				
Pain	Does not interfere with activity	Interferes with activity	Prevents daily activity	ER visit or hospitalization
Redness	1–4 caliper units (0.5–2.0 cm)	5–14 caliper units (>2.0–7.0 cm)	>14 caliper units (>7.0 cm)	Necrosis or exfoliative dermatitis
Swelling	1–4 caliper units (0.5–2.0 cm)	5–14 caliper units (>2.0–7.0 cm)	>14 caliper units (>7.0 cm)	Necrosis
Systemic event				
Vomiting	1–2 times/24 hours	>2 times/24 hours	Requiring IV hydration	ER visit or hospitalization
Diarrhea	2–3 loose stools/24 hours	4–5 loose stools/24 hours	≥6 loose stools/24 hours	ER visit or hospitalization
Headache	Does not interfere with activity	Some interference with activity	Prevents daily routine activity	ER visit or hospitalization
Fatigue	Does not interfere with activity	Some interference with activity	Prevents daily routine activity	ER visit or hospitalization
Chills	Does not interfere with activity	Some interference with activity	Prevents daily routine activity	ER visit or hospitalization
Muscle pain	Does not interfere with activity	Some interference with activity	Prevents daily routine activity	ER visit or hospitalization
Joint pain	Does not interfere with activity	Some interference with activity	Prevents daily routine activity	ER visit or hospitalization

ER=emergency room; IV=intravenous.

For the fever scale, refer to the legend and scale of **Figure 2B** and **Figure S1B**.

Table S6. Phase 1 participants 5–11 years old reporting at least 1 adverse event

	BNT162b2 10 µg (N†=16) n‡ (%)	BNT162b2 20 µg (N†=16) n‡ (%)	BNT162b2 30/30 µg* (N†=4) n‡ (%)	BNT162b2 30/10 µg* (N†=12) n‡ (%)
Adverse event				
Any event (from dose 1 to 1 month after dose 2)	7 (43.8)	5 (31.3)	2 (50.0)	3 (25.0)
Related§	4 (25.0)	2 (12.5)	2 (50.0)	3 (25.0)
Severe	0	1 (6.3)	0	0
Life-threatening	0	0	0	0
Any serious adverse event (from dose 1 through data cutoff)	0	0	0	0
Any adverse event leading to discontinuation (from dose 1 through data cutoff)	0	0	0	0
Death (from dose 1 through data cutoff)	0	0	0	0

Results are for the safety population (**Table S1**).

The data cutoff date was July 16, 2021.

* Of 16 participants who received 30 µg at dose 1, 4 received 30 µg at dose 2 and 12 received 10 µg at dose 2.

† Number of participants in the specified group. This value is the denominator for the percentage calculations.

‡ Number of participants reporting ≥ 1 occurrence of the specified event category. For ‘any event’, n=the number of participants reporting ≥ 1 occurrence of any event.

§ Assessed by the investigator as related to the investigational product.

Table S7. Phase 2/3 participants 5–11 years old reporting at least 1 adverse event

Adverse event	BNT162b2 10 µg (N*=1518) n† (%)	Placebo (N*=750) n† (%)
Any adverse event (from dose 1 to 1 month after dose 2)	166 (10.9)	69 (9.2)
Related‡	46 (3.0)	16 (2.1)
Severe	2 (0.1)	1 (0.1)
Life-threatening	0	0
Any serious adverse event (from dose 1 through data cutoff)	1 (0.1)	1 (0.1)
Related‡	0	0
Severe	1 (0.1)	1 (0.1)
Life-threatening	0	0
Any adverse event leading to discontinuation (from dose 1 through data cutoff)	0	0
Death (from dose 1 through data cutoff)	0	0

Results are for the safety population (**Table S1**).

The data cutoff date was September 6, 2021.

* Number of participants in the specified group. This value is the denominator for the percentage calculations.

† Number of participants reporting ≥ 1 occurrence of the specified event category. For ‘any event’, n=the number of participants reporting ≥ 1 occurrence of any event.

‡ Assessed by the investigator as related to the investigational product.