



CODEN [USA]: IAJ PBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Review Article

LABORATORY SCREENING FOR H.Pylori AMONG HIGH RISK POPULATION

¹Mohammed Ahmed Alshehri, ²Ahmed Mohamed Youssef, ³Faraj Ali Al-Asbali, ⁴Mohamed Issa Hassan, ⁵Abdul Rahman Mufreh Ali, ⁶Ali Mufreh Amer, ⁷Hassan Ali Alshehri, ⁸Amer Mohammed Albarqi, ⁹Ahmed Mohamed Ahmed, ¹⁰Abdu Yahya Mohamed

¹Laboratory specialist, Mjaredah General Hospital
(speaker)

²laboratory technician, Al-Qahma General Hospital

³laboratory technician, Al-Qahma General Hospital

⁴laboratories technician, Al-Qahma sector

⁵laboratory technician, Al-Qahma General Hospital

⁶laboratory technician, Al-Qahma General Hospital

⁷Laboratory technician, Mjaredah General Hospital

⁸Laboratory technician, Mjaredah General Hospital

⁹laboratory technician, Bariq sector

¹⁰laboratory technician, Mahayel General Hospital

Article Received: October 2022

Accepted: November 2022

Published: December 2022

Abstract:

Helicobacter pylori is a highly motile bacteria with numerous unipolar flagella that generates urease. H. pylori's virulence components consist of its flagella and urease. H. pylori frequently causes a chronic infection in the stomach, which can result in gastric and duodenal ulcers, gastric malignancies, and gastric lymphomas, as well as other gastrointestinal illnesses. For H. pylori, there are numerous invasive and noninvasive clinical laboratory testing. Laboratory testing is not recommended for asymptomatic patients and should only be considered if H. pylori infection therapy is planned. If an endoscopy is done on the patient, invasive testing for H. pylori, including as tissue histology, culture, and fast urease assays, are utilized. Patients whose symptoms do not require endoscopy are advised to undergo noninvasive tests for H. pylori, such as enzyme antibody and urea breath tests. Therefore, comprehensive search through literature using electronic databases; PubMed and Embase to search all relevant studies to our review topics, that were published in English language and limited to human subjects up to June 2022.

Corresponding author:

Mohammed Ahmed Alshehri,

QR code



Please cite this article in press Mohammed Ahmed Alshehri et al, *Laboratory Screening for H.Pylori Among High Risk Population., Indo Am. J. P. Sci, 2022; 09(12).*

INTRODUCTION:

Helicobacter pylori is likely the most successful human pathogen, as it has colonized the stomachs of roughly half of the world's population [1]. In several regions of the world, the incidence is even higher, such as 80% or more in portions of China and some Eastern European and South American countries, although it is decreasing in the United States and Western European nations [2,3]. Less-developed areas are more likely to have high *H. pylori* prevalence and higher rates of recurrence of infection after treatment [4]; rates of infection are correlated with socioeconomic characteristics such as level of education and living standards.

Despite a reducing prevalence, gastric cancer was the fifth most prevalent cancer and the third leading cause of cancer-related deaths globally in 2012 [5], accounting for over three-quarters of a million fatalities annually. The current accepted model for stomach carcinogenesis is an extension of the original model published in 1975 by Correa *et al.* [6]. This model proposes that gastric cancer is the end result of a series of mutations that begin with an unknown environmental trigger in early life, now known to be infection with *H. pylori*, resulting in a superficial gastritis, followed by chronic non-atrophic gastritis, gastric atrophy, and achlorhydria [7]. The development of gastric intestine metaplasia, which assumes increasingly basic forms, is followed by cell change, dysplasia, and ultimately cancer (Figure 1) [6].

H. pylori generates a pattern of gastritis known as acute-on-chronic inflammation, which leads to chronic progressive gastric damage and, eventually, gastric atrophy [8]. Stomach cancer is a malignancy associated with inflammation in which the infection directly and indirectly causes progressive genetic damage to the gastric epithelium, which may finally develop to gastric adenocarcinoma. In *H. pylori*-infected patients, the risk of developing gastric cancer can be calculated based on the degree and extent of mucosal damage and atrophy, identified as metaplastic epithelia with or without intraepithelial neoplasia or dysplasia. Eradication of *H. pylori* reduces the risk of gastric cancer, which is proportional to the extent of genetic and epigenetic changes existing at the time of *H. pylori* eradication [9,10]. The translation of this basic and clinical information into public health intervention through population-wide screening and eradication of *H. pylori*, which would ultimately prevent the development of stomach cancer, is, however, lacking. First, we need up-to-date information on the worldwide disease burden of *H.*

pylori infection and stomach cancer, and we must better define the target groups for screening and eradication initiatives. Second, the implementation of mass screening requires consideration of the choice of non-invasive test, when to proceed to endoscopy, whether and how to test for eradication efficacy and how to survey those with advanced atrophic gastritis, how to implement this programmed, how to treat asymptomatic *H. pylori*-infected subjects for gastric cancer prevention, and how to identify subjects at higher risk of gastric cancer for endoscopic examination [10,11].

DISCUSSION:

Numerous studies have demonstrated that *H. pylori* infection is a key factor in the etiology of peptic ulcers, gastric ulcers, noncardia gastric cancer, low-grade mucosa-associated lymphoid tissue (MALT) lymphoma of the stomach, and other diseases, including non-ulcer dyspepsia and gastrointestinal reflux diseases. *H. pylori* is a gram-negative spiral bacterium that colonizes the gastric epithelium [12,13], but not the duodenal epithelium. The organism possesses specific, crucial characteristics: A) *H. pylori* create high quantities of urease, which is essential for colonization, pathogenicity, and neutralizing the effects of an acidic environment via the production of ammonia [14,15]. B) *In vivo*, *H. pylori* promotes epithelial cell growth and apoptosis. Nevertheless, infection with bacteria of the CagA genotype results in greater proliferation than apoptosis [16]. C) The production of a vacuolating toxin and the presence of the CagA gene, which encodes virulence genes involved in the stimulation of epithelial chemokine response, are the primary determinants of *H. pylori* [16]. The body of the stomach becomes highly inflamed at initial infection with *H. pylori*, and acid output is blocked [12]. With *H. pylori* infection, the inflammation migrates to the antrum, resulting in antral gastritis and increased acid output. Reduced bicarbonate secretion, intestinal inflammation, and ulceration are observed in the duodenum [12,16]. Those infected with *H. pylori* who develop acid hypersecretion are susceptible to developing duodenal ulcers. If duodenal ulcer is evident, eradication of the *H. pylori* bacteria from the gastric mucosa is curative [12]. Most infected individuals may develop a symbiotic connection with the *H. pylori* organism. Nonetheless, a few patients with illness in the stomach body may have normal or reduced acid secretion and develop gastric ulcers [14,15]. *H. pylori* is implicated in the pathophysiology and development of both adenocarcinoma and stomach MALT lymphoma, according to substantial evidence. *H. pylori*'s mechanism and significance in the etiology of gastric

cancer and lymphoma remain undefined. *H. pylori* infection is a risk factor. However, only a small fraction of *H. pylori*-infected patients actually

develops cancers. Therefore, *H. pylori* must interact with additional environmental and genetic cofactors [15,16].




FIG. 1: A model for the development of gastric cancer [6].

Indications for Diagnostic Testing:

Currently, the diagnostic gold standard for *H. pylori* infection necessitates the histologic inspection of two specifically stained stomach antral biopsy specimens (28,29). Sampling and observer error have the potential to restrict the utility of routine histology in normal clinical practice. Currently, there are no clearly and reliably defined indications or guidelines for determining whether a child or an adult should undergo definitive testing for *H. pylori* infection [17,18]. *H. pylori* testing should only be deemed appropriate if treatment is planned. Currently, there is no evidence that *H. pylori* eradication is beneficial for asymptomatic children, adults, or adults [19]. Therefore, the physician should utilize a meticulously documented history of gastrointestinal complaints and symptoms to determine *H. pylori* diagnosis indicators (TABLE.1). If *H. pylori* is detected by a credible test, the infection is presumed to be treated [19].

General Diagnostic Testing

The reference method (gold standard) for the diagnosis of active *H. pylori* infection is esophago-gastro-duodenoscopy with gastric biopsies. At present, there are numerous other accurate detection assays, which can be categorized as invasive or noninvasive. The primary invasive technique is endoscopy with biopsy, which enables histological examinations to identify the microorganisms directly. Most of the noninvasive techniques rely on the detection of biochemical properties of *H. pylori* (the ability to hydrolyze urea), or the response of the immune system with the production of specific antibodies [19].

Histological testing:

For the diagnosis of *H. pylori* infections, histological staining of stomach biopsies is still regarded as the

gold standard. Two specifically coloured gastric antral biopsy specimens are examined [20,21]. In the hands of professionals, Giemsa, Warthin-Starry, and Genta stains can be as accurate as 98%, according to studies [18,21]. These two stains increase the percentage of organisms that can be positively identified. Histological investigation necessitates expensive intrusive procedures, such as endoscopy. In regular clinical practice, sampling error is cited as a source of inaccurate histological diagnosis; hence, some authors have suggested that additional specimens be collected from the lesser curvature and/or the stomach body to improve diagnostic accuracy [19,21].

Culture as testing method:

H. pylori is a microaerophilic bacterium that must be cultured as a testing method. Culture of biopsies is the most specific approach for diagnosing *H. pylori* infection [18,19]. Due to the organism's sensitivity, however, the culture method is exceedingly insensitive; hence, routine culture cannot be considered an appropriate gold standard for general clinical practice.

In tissue culture, *H. pylori* grows slowly and requires 2–5 days to become positive. Identification is based on Gram stain morphology in addition to positive urease, catalase, and oxidase reactions.

Due to the organism's sensitivity, the culture yield may decrease if the stomach biopsies are not performed correctly. Consequently, a negative culture result does not exclude *H. pylori* infection. With the culture approach, experienced laboratories have demonstrated sensitivities of at least 90 percent [22,23].

The major advantage of utilizing culture as a

diagnostic method is that culture and isolation of the *H. pylori* bacterium can help select the choice of antibiotic [23].

Rapid Urease Tests

Owen et al. [24] reported in 1985 that *H. pylori* exhibited a quick urease hydrolysis reaction that differentiated it from other bacteria. Several diagnostic kits based on the urease reaction have been developed.

The test requires the addition of a gastric mucosal biopsy to a urea substrate and a pH-sensitive marker in order to detect the ammonia produced by the urease reaction during pH rise [19].

At 1 hour, the test endpoint is measured. When read after more than one hour, there are several false-positive results, and specificity drops to 68% [25]. Rapid urease tests are straightforward to administer, however they require endoscopic biopsies.

The sensitivity of highly sensitive fast urease tests was shown to be comparable to that of culture regardless of testing time, but lower than that of histology a few months after treatment [26]. However, the very

sensitive fast urease test was accurate in verifying treatment success [26] when utilized many months following therapy.

Polymerase Chain Reaction-Based Molecular Diagnosis (PCR):

Molecular assays, such as PCR, may be utilized for the exact diagnosis of *H. pylori* infections. Testing bacteria with molecular techniques does not require that they be alive. Numerous PCR techniques are highly accurate at diagnosing *H. pylori* from clinical biopsies. It is widely recognized that PCR adds little to other procedures when used to biopsy samples [27]. The combination of culture and histology detects *H. pylori* in about the same proportion of patients as PCR [28,29]. Gastric juice, which can be obtained through nasogastric catheter, demonstrates the principal advantage and utility of PCR. Westblom et al. [30], utilizing PCR on specimens of 5 mL gastric juice, detected *H. pylori* infection with a sensitivity of 96% and a specificity of 100%. With the use of a highly sensitive semi-nested PCR assay on stomach juice samples taken with capsulated strings, Yoshida et al. [31] found 97% of cases of recurrent infection of *H. pylori* within 8 wk following antimicrobial therapy.

TABLE. 1: Diagnostic tests for *Helicobacter pylori*

Tests	Advantages	Disadvantages
Non-invasive		
Serology	Widely available, relatively inexpensive	Cannot be used to confirm eradication after treatment; does not
Urea breath test	High sensitivity and specificity; no specific transport conditions; good test to use to confirm eradication of <i>H. pylori</i> infection after treatment	permit antimicrobial sensitivity Cannot do antimicrobial sensitivity; use ¹⁴ C radioactive-labeled material
Stool <i>H. pylori</i> antigen test	Useful to confirm eradication of <i>H. pylori</i> a few weeks after treatment	Collection of stool specimen
Invasive		
Histology	Can estimate presence of <i>H. pylori</i> and extent of inflammation and damage; can do retrospective examination	Endoscopy to obtain samples; performance depends on experience of pathologist; cannot do antimicrobial susceptibility studies
Culture	100% specific; allows testing for antimicrobial sensitivity; permits typing of strains	Endoscopy needed to obtain samples
Rapid urease test	Close to 100% specific; results within serial hematoxylin-eosin-stained tissue sections 1–2 hours	Endoscopy needed; no antimicrobial sensitivity studies can be done
Molecular methods (PCR)	High sensitivity and specificity; can do retrospective analysis; applicable to special samples such as oral and gastric fluids, stool, environmental samples	Does not permit antimicrobial susceptibility testing; requires skills in testing personnel

CONCLUSION:

It is estimated that around fifty percent of the world's population is infected with *H. pylori*. Humans appear to be the infection's natural host and source. For in vivo proliferation, *H. pylori* requires gastric-type mucosa, and eating appears to be the most prevalent route of infection. Numerous investigations have shown that *H. pylori* infection has a significant role in the etiology and pathophysiology of gastric and duodenal ulcers, gastric cancer, gastric MALT lymphoma, and other gastrointestinal illnesses. There are numerous diagnostic tests available for *H. pylori* infections (Table 1). In general, laboratory testing in asymptomatic children and adults is not indicated. Testing should only be considered if *H. pylori* infection therapy is planned. The most crucial factor is whether or not the patient will undergo endoscopy.

REFERENCES:

1. Pounder, R. E. & Ng, D. The prevalence of *Helicobacter pylori* infection in different countries. *Aliment. Pharmacol. Ther.* 9 (Suppl. 2), 33–39 (1995).
2. Roberts, S. E. et al. Review article: the prevalence of *Helicobacter pylori* and the incidence of gastric cancer across Europe. *Aliment. Pharmacol. Ther.* 43, 334–345 (2016).
3. Graham, D. Y. et al. Epidemiology of *Helicobacter pylori* in an asymptomatic population in the United States. *Gastroenterology* 100, 1495–1501 (1991).
4. Yan, T. L., Hu, Q. D., Zhang, Q., Li, Y. M. & Liang, T. B. National rates of *Helicobacter pylori* recurrence are significantly and inversely correlated with human development index. *Aliment. Pharmacol. Ther.* 37, 963–968 (2013).
5. Torre, L. A. et al. Global cancer statistics, 2012. *CA Cancer J. Clin.* 65, 87–108 (2015).
6. Correa, P., Haenszel, W., Cuello, C., Tannenbaum, S. & Archer, M. A model for gastric cancer epidemiology. *Lancet* 306, 58–60 (1975).
7. Correa, P. The gastric precancerous process. *Cancer Surv.* 2, 437–450 (1983).
8. Wong BC-Y, Lam SK, Wong WM, et al. *Helicobacter pylori* eradication to prevent gastric cancer in a high-risk region of China: a randomized controlled trial. *JAMA* 2004;291:187–94.
9. Choi IJ, Kook M-C, Kim Y-I, et al. *Helicobacter pylori* therapy for the prevention of metachronous gastric cancer. *N Engl J Med* 2018;378:1085–95.
10. Choi IJ, Kim CG, Lee JY, et al. Family history of gastric cancer and *Helicobacter pylori* treatment. *N Engl J Med* 2020;382:427–36.
11. Lee Y-C, Chiang T-H, Chou C-K, et al. Association between *Helicobacter pylori* eradication and gastric cancer incidence: a systematic review and meta-analysis. *Gastroenterology* 2016;150:1113–24.
12. Walker MM, Crabtree JE. *Helicobacter pylori* infection and the pathogenesis of duodenal ulceration. *Ann N Y Acad Sci* 1998;859:96–111.
13. Axon ATR, Forman D. *Helicobacter gastro-duodenitis*: a serious infectious disease. *Br Med J* 1997;314:1430–1431.
14. Veldhuyzen van Zanten SJO, Lee A. The role of *Helicobacter pylori* infection in duodenal and gastric ulcer. *Curr Topics Microbiol Immunol* 1999;241:47–56.
15. Mirkhopadhyay P. Gastric cancer and lymphoma. *Curr Topics Microbiol Immunol* 1999;241:57–69.
16. Moss SF. *Helicobacter pylori* and apoptosis. *Yale J Biol Med* 1998;71:53–61.
17. Leu J, O'Moran C. Consensus or confusion: a review of existing guidelines on *Helicobacter pylori* related disease. *Eur J Gastroenterol Hepatol* 1997;8:531–541.
18. Metz DC, Furth EE, Faigel DO, et al. Realities of diagnosing *Helicobacter pylori* infection in clinical practice: a case for non-invasive indirect methodologies. *Yale J Biol Med* 1998;71:81–90.
19. Westblom TU, Bhatt BD. Diagnosis of *Helicobacter pylori* infection. *Curr Topics Microbiol Immunol* 1999;241:215–235.
20. Genta RM, Robason GR, Graham DY. Simultaneous visualization of *Helicobacter pylori* and gastric morphology: a new strain. *Hum Pathol* 1994;25:221–226.
21. Genta RM, Graham DY. Comparison of biopsy sites for the histopathologic diagnosis of *Helicobacter pylori*: a topographic study of *H. pylori* density and distribution. *Gastrointest Endosc* 1994;40:342–345.
22. Deltenre M, Glupezynski Y, DePrez C, et al. The reliability of urease tests, histology, and culture in the diagnosis of *Campylobacter pylori* infection. *Scand J Gastroenterol* 1989;160(Suppl):19–24.
23. Nichols L, Sughayer M, DeGerolani PC, et al. Evaluation of diagnostic methods for *Helicobacter pylori* gastritis. *Am J Clin Pathol*

- 1991;95:769–773.
24. Owen RJ, Martin SR, Borman P. Rapid urea hydrolysis by gastric *Campylobacter*. *Lancet* 1985;1:111.
 25. Yousfi MM, el-Zimaity HM, Cole RA, Genta RM, Graham DY. Comparison of agar gel (CLO test) on reagent strip (Pylori Tek) rapid urease test for detection of *Helicobacter pylori* infection. *Am J Gastroenterol* 1997;92:997–999.
 26. Murata H, Tsuji S, Kawano S. Possible availability of rapid urease test for diagnosis of *H. pylori* eradication: comparative study with culture and histology. *Nippon Rinsho* 1999;57:97–100.
 27. Ashton-Key M, Diss TC, Isaacson PG. Detection of *Helicobacter pylori* in gastric biopsy and resection specimens. *J Clin Pathol* 1996;49:107–111.
 28. Len SY, Jeng YS, Wang CK, et al. Polymerase chain reaction diagnosis of *Helicobacter pylori* in gastrointestinal diseases: comparison with culture and histopathological examinations. *J Gastroenterol Hepatol* 1996;11:286–289.
 29. Lage AP, Godfried E, Fauconnier A, et al. Diagnosis of *Helicobacter pylori* infection by PCR: comparison with other invasive techniques and detection of *CagA* gene in gastric biopsy specimens. *J Clin Microbiol* 1995;33:2752–2756.
 30. Westblom TU, Phadnis S, Yang P, Czinn SJ. Diagnosis of *Helicobacter pylori* infection by means of a polymerase chain reaction for gastric juice aspirates. *Clin Infect Dis* 1993;16:367–371.
 31. Yoshida H, Maeda S, Ogura K. PCR-monitoring of gastric juice obtained with the capsulated string for evaluation of *H. pylori* infection. *Nippon Rinsho* 1999;57:107–110.