



## DESIGN, SYNTHESIS AND BIOLOGICAL ACTIVITY OF 1, 4-BENZOXAZINONE DERIVATIVES AS ANTI-MICROBIAL AGENTS

P. Aravinda Reddy<sup>1</sup>, Ramya Sri. S<sup>2</sup>

<sup>1</sup>Department of Pharmaceutical analysis, Vaageswari College of Pharmacy, Karimnagar, Telangana-505527, India

<sup>2</sup>Department of Pharmacy, University College of Technology, Osmania University, Hyderabad, Telangana, 500007, India

Received on: 22-12-2017; Revised and Accepted on: 31-12-2017

### ABSTRACT

Column chromatography is used to separate the components. Analytical and spectral data (IR, <sup>1</sup>H NMR, and Mass) are used to identify and describe all novel compounds. Synthesized chemicals are tested for their ability to kill bacteria and fungi. The synthesised benzoxazinone derivatives were evaluated and found to be moderately to weakly antimicrobial. The antibacterial activity of the fused or combined moieties was found to be moderate to weak against Gram (+ve) and Gram (-ve) bacteria in screening experiments. Various biological activities have been attributed to the substituted benzoxazinone derivatives.

**Keywords:** benzoxazinone, derivatives, spectral data.

### INTRODUCTION

The subject of drugs is as old as disease. Illness has been man's heritage from the beginning of his existence, along with globalization many new diseases are arising in the world and the search for remedies combat it is perhaps equally old. The sincere attempt by man to control and cure diseases has led to search of new drugs or suitable derivatives of existing drugs. The earlier sources of drugs were from plant, animal and mineral sources, but due to the lack of potential action and definitive cure and sometimes more toxicity, the discovery of new drugs that are more potential and less toxic is essential. The synthesis of derivatives has been an important part and is aimed at modifying the action of drugs, particularly to reduce the side effects and to potentiate the drug action. Today more than 60% drugs used in practice are synthesized derivatives and day-by-day the scope of synthetic medicinal chemistry is broadening. Drugs are chemicals that prevent disease or assist in restoring health to the diseased individuals as such they play an indispensable role in modern medicine. Medicinal chemistry is the branch of science that provides these drugs either through discovery or through design. In the last century, the classical drugs were primarily discovered either by alteration of natural

substances or entirely by chemical synthesis. In the recent years, an ever-increasing understanding of pathophysiology of diseases has increasingly led to novel opportunities to design, synthesis and evaluation of candidate drug molecules.<sup>1</sup>

During the later decades of 20th century traditional dividing lines between biological, chemical and physical sciences were erased and in the present millennium, new borderline investigation such as ,molecular biology, molecular pharmacology, biomedicine, bio metallic chemistry, cellular biology, genetics, bioinformatics and others have begun to capture the interest of medicinal scientist. With the race to discover novel drug molecules, the drug discovery program has become a multidisciplinary approach. Thus, the activity of medicinal chemist essentially encompass biological (understanding of the enzymology and pharmacology) and physicochemical aspects (understanding of thermodynamic quantum mechanics, spectroscopy, crystallography etc.) these aspects have acquired increasingly mechanistic underpinnings with the development and quantification of the enzyme/receptor concept and the availability of receptor-based assays. Lead discovery and lead optimization are the two important concepts of drug design in medicinal chemistry and the process of lead structure identification and refinement, now manifest further sophistication in accordance with developments in structural and molecular biology and the combinatorial chemical synthesis and automated screening techniques like high throughout screening.<sup>1</sup>

#### \*Corresponding author:

P. Aravinda Reddy  
Department of Pharmaceutical analysis  
Vaageswari College of Pharmacy, Karimnagar  
Telangana-505527  
India

Email: [surapharmalabs1@gmail.com](mailto:surapharmalabs1@gmail.com)

DOI:

BIOLOGICAL ACTIVITY METHODS

ANTIBACTERIAL ACTIVITY

The compounds synthesized during the present investigation were screened for their antibacterial activity. The antibacterial activity of the compounds was assessed by cup and plate method. The antibacterial tests were conducted on six common microorganisms which are the representative types of gram

**The results of screening were included in the below table-1.**

Gram positive bacteria	Gram negative bacteria
Bacillus subtilis	Pseudomonas aeruginosa
Staphylococcus aureus	Escherichia coli
Staphylococcus epidermidis	Klebsiella pneumoniae

#### Preparation of nutrient agar media:

The nutrient agar media was prepared by using the following ingredients.

Peptone (Bacteriological): 20 g.

Beef extract (Bacteriological): 5 g.

Sodium chloride (Bacteriological): 5 g.

Agar (Bacteriological): 20 g.

Distilled water up to: 1000 ml.

Weighed quantities of peptone, beef extract were dissolved in distilled water by gentle warming, and then the specified amount of agar was dissolved by heating on boiling water bath. Then the pH of the above solution is adjusted by adding sodium chloride and the volume of final solution is made up to 1000 ml with distilled water. Then the above prepared nutrient agar media is sterilized by autoclave at 121 oC for 20 minutes at 15 lbs/in<sup>2</sup> pressure.

#### Preparation of test solution:

20 mg of the test compound was dissolved in 20 ml of DMF, from this stock solution, 1 ml of solution was taken and further diluted to required concentration with DMF. These sample solution were made in suitably labeled sterilized test tubes.

Preparation of standard solution:

The standard drug used for the comparison are Penicillin and Streptomycin, the solutions were prepared from sterile water soluble.

Method of testing:

The above prepared nutrient agar media is cooled to 45 oC with gentle shaking to bring about uniform cooling. To this 0.5–0.6 ml of 18-24 h old culture was injected aseptically and mixed well by gentle shaking. This was poured onto the petri dishes and was allowed to set for 1 h.

Thereafter the cups were made by punching into the set agar with a sterile cork borer and scooping out the punched part of the agar. The diameter of each cup was 6 mm. To these cups 50 µl of the test compound was put, which was prepared in DMF. After adding the drug solution, it was allowed to diffuse for about 45 minutes, at room temperature. Then the plates were incubated at 37 oC for 24 h in an incubator.

The minimum inhibitory concentration (MIC) is taken as a parameter of antibacterial activity.

Standard: Penicillin, Streptomycin.

Note: 20 µg/ml and above poor activity, 14-20 µg/ml moderate activity and 4-13 µg/ml significant activity.

#### ANTIFUNGAL ACTIVITY

##### ANTIFUNGAL ACTIVITY:

##### Procedure:

The anti-fungal activity of all compounds was determined on potato dextrose agar medium against *Rhizopus oryzae*, *Aspergillus niger*, *Aspergillus flavus* *Candida albicans*, and

**Table No-2: Antibacterial activity of synthesized compounds (3A-F)**

Compound Code	Minimum Inhibitory Concentration (µg/ml)					
	Gram Positive Organism			Gram Negative Organism		
	B.Subtilis	S.aureus	S.epidermis	E.coli	P.aeruginosa	K.pneumoniae
3A	23.15	150	17.10	75	150	18.0
3B	18.60	75	22.50	14.55	13.0	75
3c	24.75	4.65	75	75	17.75	16.75
3D	24.50	14.25	1.16	75	75	29.35
3E	75	30.15	75	150	150	75
3F	21.60	75	85	13.50	75	150
Penicillin	1.526	6.25	3.125	7.81	12.50	6.25
Streptomycin	6.25	1.56	1.56	3.125	3.125	3.125

*Saccharomyces cerevisiae* organisms using Amphotericin-B as a standard and DMF was used as control. The sterile molten potato dextrose medium was cooled to 45 oC and inoculated with test organisms and mixed the contents thoroughly and poured into the sterile petri dishes under aseptic conditions. All the inoculated petri dishes were incubated at 28 oC for 4 days and the extent diameter of inhibition was measured as the zone of inhibition in millimeters and the results were shown in Table No-3.

#### Preparation of sub-culture:

Peeled potato: 200-300 gm

Dextrose: 5 gm

Distilled water up to: 1000 ml

Peeled potato were cut into pieces and boiled for 30 min to get extract. The extract is filtered through muslin cloth. The dextrose was added to the filtrate and final volume is adjusted to 1000 ml with distilled water. Then it was sterilized by autoclave at 121 oC for 20 min.

Note: Two days before testing the culture is prepared by inoculating the fungus from master culture into potato dextrose medium and incubated for 48 h at room temperature

#### Preparation of fungal medium:

The potato dextrose medium is prepared by dissolving.

Peeled potato: 200-300 gm

Dextrose : 5 gm

Agar: 20 gm

Distilled water up to: 1000 ml

Peeled potato were cut into pieces and boiled for 30 min to get the extract. The extract is filtered through muslin cloth. The dextrose and agar were added to the filtrate and the final volume is adjusted to 1000 ml with distilled water. Then it was sterilized by autoclave at 121 oC for 20 min.

#### Preparation of test solution:

10 mg of the compound was dissolved in 10 ml of DMF, from this stock solution further required concentration solutions were prepared by dilution using DMF.

#### Preparation of standard anti-fungal solution:

Amphotericin-B was used as standard anti-fungal for comparison and solution were prepared by using sterile water, so that the concentrations of the solution were 100 µg/ml.

#### Method of testing:

The method of testing for fungicidal activity is the same as that of antibacterial testing. DMF was used as a solvent control.

**Table No-3 Antifungal activity of synthesized compounds (3<sub>A-F</sub>)**

Compound (100 µg/ml)	Zone of Inhibition(mm)				
	R. oryzae	A.niger	A.flavis	C.albicans	S.cerevisiae
3 <sub>A</sub>	24	15	-	-	16
3 <sub>B</sub>	17	16	12	14	22
3 <sub>C</sub>	16	14	18	12	12
3 <sub>D</sub>	15	14	12	12	10
3 <sub>E</sub>	28	16	14	-	-
3 <sub>F</sub>	14	-	21	15	16
Amphotericin-B	24	25	24	23.5	22


Standard: Amphotericin-B

Note:- '-' denotes no activity, 10-14 mm poor activity, 15-17 mm moderate activity, 18-25 mm significant activity.

## RESULTS AND DISCUSSION

## SPECTRAL DATA

## Compound (1)

2*H*-benzo[*b*][1,4]oxazin-3(4*H*)-one<sup>1</sup>H NMR Spectra (CDCl<sub>3</sub>, δ ppm):


Value in δ ppm	Nature of segments	No of protons	Type of proton
8.98	Singlet	1H	1H of NH
6.8-7.2	Multiplet	4H	4H of Ar-H
4.64	Singlet	2H	2H of CH <sub>2</sub>

## MASS Spectra (m/z):

Molecular weight of the compound is 149; the molecular ion peak appeared at 150 as M+1.

Type of Vibration	Group frequency in Wave number (cm <sup>-1</sup> )
NH stretching	3133
Ar-CH=CH Stretching	2982 and 2902
C=O Absorption band	1709

## Compound 2


4-(3-Bromo-propyl)-4*H*-benzo[1,4]oxazin-3-one<sup>1</sup>H NMR Spectra (CDCl<sub>3</sub>, δ ppm):

Value in δ ppm	Nature of segments	No of protons	Type of proton
7.01-7.07	Multiplet	4H	4H of Ar-H
4.60	Singlet	2H	CH <sub>2</sub>
4.09	triplet	2H	2H of CH <sub>2</sub>
2.25	Pentet	2H	2H of CH <sub>2</sub>
3.40	Triplet	2H	2H of CH <sub>2</sub>

## MASS Spectra (m/z):

Molecular weight of the compound is 269; the molecular ion peak appeared at 270 as M+1.

## Compound

4-(3-Morpholin-4-yl-propyl)-4*H*-benzo[1,4]oxazin-3-one<sup>1</sup>H NMR Spectra (CDCl<sub>3</sub>, δ ppm):

Value in δ ppm	Nature of segments	No of protons	Type of proton
6.99-7.09	Multiplet	4H	4H of Ar-H
4.60	Singlet	2H	CH <sub>2</sub>
4.00	triplet	2H	2H of CH <sub>2</sub>
1.89	Pentet	2H	2H of CH <sub>2</sub>
2.48	Triplet	6H	CH <sub>2</sub> , CH <sub>2</sub> (morpholine)
3.72	Triplet	4H	CH <sub>2</sub> (morpholine)


## MASS Spectra (m/z):

Molecular weight of the compound is 276; the molecular ion peak appeared at 277 as M+1.

IR Spectra (cm<sup>-1</sup>):

Type of Vibration	Group frequency in Wave number (cm <sup>-1</sup> )
=CH stretch	2922
=CH bend	823
CH <sub>2</sub> stretch	2851
Alkane CH <sub>2</sub> stretch	2816
C=O stretch	1606

## Compound 3B

4-{3-[4-(4-Trifluoromethyl-phenyl)-piperazin-1-yl]-propyl}-4*H*-benzo[1,4]oxazin-3-one

**<sup>1</sup>H NMR Spectra (DMSO, δ ppm):**


Value in δ ppm	Nature of segments	No of protons	Type of proton
6.90-7.48	Multiplet	8H	8H of Ar-H
4.60	Singlet	2H	CH <sub>2</sub>
4.00	Triplet	2H	2H of CH <sub>2</sub>
1.92	Pentet	2H	2H of CH <sub>2</sub>
3.28	Triplet	4H	CH <sub>2</sub>
2.62	Triplet	4H	CH <sub>2</sub>
2.50	Triplet	2H	CH <sub>2</sub>

**MASS Spectra (m/z):**

Molecular weight of the compound is 419; the molecular ion peak appeared at 420 as M+1

**IR Spectra (cm<sup>-1</sup>):**

Type of Vibration	Group frequency in Wave number (cm <sup>-1</sup> )
=CH stretch	2922
=CH bend	823
CH <sub>2</sub> stretch	2851
Alkane CH <sub>2</sub> stretch	2816
C=O stretch	1606

**Compound 3C****4-{3-[4-(4-Methoxy-phenyl)-piperazin-1-yl]-propyl}-4H-benzo[1,4]oxazin-3-one****<sup>1</sup>H NMR Spectra (CDCl<sub>3</sub>, δ ppm):**

Value in δ ppm	Nature of segments	No of protons	Type of proton
6.83-7.11	Multiplet	8H	8H of Ar-H
4.60	Singlet	2H	CH <sub>2</sub>
4.00	triplet	2H	2H of CH <sub>2</sub>
1.92	Pentet	2H	2H of CH <sub>2</sub>
3.11	Triplet	4H	CH <sub>2</sub>
2.62	Triplet	4H	CH <sub>2</sub>
2.49	Triplet	2H	CH <sub>2</sub>
3.77	singlet	3H	CH <sub>3</sub>

**MASS Spectra (m/z):**

Molecular weight of the compound is 381; the molecular ion peak appeared at 382 as M+1.

**IR Spectra (cm<sup>-1</sup>):**

TYPE OF VIBRATION	GROUP FREQUENCY IN WAVE NUMBER(cm <sup>-1</sup> )
C=O stretch	1606
=CH stretch	2957
=CH bend	860
Alkane CH <sub>2</sub> stretch	2817

**BIOLOGICAL EVALUATION**

From the literature survey, it reveals that novel benzoxazinones have been reported for number of pharmacological activities and some molecules have shown significant activities and some compounds shows moderate and good activities. All the synthesized compounds (**3A-F**) were screened them for their anti-bacterial and anti-fungal activities and the results are as follows:

**ANTIMICROBIAL STUDIES****ANTIBACTERIAL ACTIVITY:**

All the synthesized compounds were screened for antibacterial activity studies against *Bacillus subtilis*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* by cup-plate technique on nutrient agar media, Penicillin and Streptomycin used as standard against Gram positive and Gram-negative bacteria.

The data in the **Table No-1** indicate that,

- ❖ Compound **3c** show significant activity and compound **3d** shows moderate activity against *staphylococcus aureus*.
- ❖ Compound **3d** also shows moderate activity against *staphylococcus epidermidis*.
- ❖ Compound **3b** shows moderate activity against *Escherichia coli*.
- ❖ Compound **3b** and Compounds **3c** shows moderate activity against *Pseudomonas aeruginosa*.
- ❖ Compound **3a** and Compounds **3c** shows moderate activity against *Klebsiella pneumoniae*.


And rest of the compounds were found to exhibit poor activity when compared to the standard Penicillin and Streptomycin.

Table No-4 SPECTRAL DATA OF THE SYNTHESISED COMPOUNDS

C.N O	IUPAC Name	MOLECULAR FORMULA	MOL. WT	<sup>1</sup> H NMR in CDCl <sub>3</sub> +DMSO (JCAMP), δ, J(Hz)	MS (m/z)	IR (KBr disc in cm <sup>-1</sup> )
I	4H- Benzo[1,4]oxazin-3- one	C <sub>8</sub> H <sub>7</sub> NO <sub>2</sub>	149	8.98(s,1H,NH),4.64(s, 2H, CH <sub>2</sub> ), 6.80- 7.20(m, 4H, Ar-H)	150  (M+H)	3133 NH- Strech 2982 =CH strech 2856 CH <sub>2</sub> strech 1606 C=Ostrech 888 =CH bend
II	4-(3-Bromo- propyl)-4H- benzo[1,4]oxazin- 3-one	C <sub>11</sub> H <sub>12</sub> O <sub>2</sub> N Br	269	2.25(p, 2H, CH <sub>2</sub> ), 4.60(s, 2H, CH <sub>2</sub> ), 3.40(t, 2H, CH <sub>2</sub> ), 4.09(t, J=7.554, 2H, CH <sub>2</sub> ), 7.01- 7.07(m, 4H, Ar-H).	270  (M+H)	-----
3A	4-(3-Morpholin-4- yl-propyl)-4H- benzo[1,4]oxazin- 3-one	C <sub>15</sub> H <sub>20</sub> O <sub>3</sub> N <sub>2</sub>	276	6.99-7.09(m, 4H, Ar-H), 4.60(s, 2H, CH <sub>2</sub> ), 4.00(t, J=7.365, 2H, CH <sub>2</sub> ),1.89(p, 2H, CH <sub>2</sub> ),2.48(6H,CH <sub>2</sub> ), 3.72(d,4H,CH <sub>2</sub> (Morpholine))	277  (M+H)	2922 =CH strech 2851 CH <sub>2</sub> stretch 2816 alkane CH <sub>2</sub> str 1606 C=O stretch 823 =CH bend
3B	4-{3-[4-(4- Trifluoromethyl- phenyl)-piperazin-1- yl]-propyl}-4H- benzo[1,4]oxazin-3- one	C <sub>22</sub> H <sub>24</sub> O <sub>2</sub> N <sub>3</sub> F <sub>3</sub>	419	6.90-7.48(m,8H,Ar-H), 4.60(s, 2H, CH <sub>2</sub> ), 4.00(t, J=7.176, 2H, CH <sub>2</sub> ), 3.28(d,4H,CH <sub>2</sub> ) ,2.62(d,4H,CH <sub>2</sub> ) ,2.50(t, J=6.987, 2H, CH <sub>2</sub> ),1.92(p, 2H, CH <sub>2</sub> )	420  (M+H)	2923 =CH stretch 2856 CH <sub>2</sub> stretch 1683 C=O stretch
3C	4-{3-[4-(4- Methoxy-phenyl)- piperazin-1-yl]- propyl}-4H- benzo[1,4]oxazin- 3-one	C <sub>22</sub> H <sub>27</sub> O <sub>3</sub> N <sub>3</sub>	381	6.83-7.11(m,8H,Ar-H), 4.60(s, 2H, CH <sub>2</sub> ), 4.00(t, J=7.554, 2H, CH <sub>2</sub> ), 3.11 (d,4H,CH <sub>2</sub> ),3.77(s,3H,CH <sub>3</sub> ),1.91(p, 2H, CH <sub>2</sub> ), 2.49(t, J=6.798, 2H, CH <sub>2</sub> ), 2.62(d,4H,CH <sub>2</sub> )	382  (M+H)	1606 C=O stretch 2817 alkane CH <sub>2</sub> str 2957 =CH stretch 860 =CH bend




**Fig No.1 Antibacterial activity of synthesized compounds (3A-F)**



7b



6b




#### ANTIFUNGAL ACTIVITY:

The antifungal activities of the test compounds were tested by agar diffusion method (cup-plate method) taking drug at a concentration of 150µg/50 µL against five fungal organisms. The area of Zone of Inhibition (ZOI) was taken as a parameter of antifungal activity. The ZOI of the compound was compared to that of the standard drug i.e. Amphotericin-B.

The compounds were screened at 150µg/50 µL showed different ZOI against five fungal organisms (Table 2). Among the compounds screened the maximum ZOI was observed against *Saccharomyces cerevisiae*, *Aspergillus niger*, *Rhizopus oryzae*, *Candida albicans* and *C. rugosa*.

- 3A and 3E show significant activity and compounds 3B and 3C show moderate activity against *Rhizopus oryzae*

- compound 3D show moderate activity against *Aspergillus niger* and
- Compound 3F show significant activity and compound 3C show moderate activity against *Aspergillus flavus* and
- compound 3F show moderate activity against *Candida albicans* and
- Compound 3B show significant activity and compounds 3A and 3F show moderate activity against *Saccharomyces cerevisiae* and rest of the compounds were found to exhibit poor activity when compared to the standard Amphotericin-B.



#### CONCLUSION

The compounds are purified by column chromatography. All new compounds are characterized by the analytical and spectral (IR, <sup>1</sup>H NMR and Mass) data. The compounds synthesized are evaluated for antibacterial, antifungal activity

It is clearly concluded that the synthesized benzoxazinone derivatives were found to be moderate to weak antibacterial agents. When the two moieties are fused or combined and screened for antibacterial studies they showed moderate to weak antibacterial activity against Gram (+ve) and Gram (-ve) bacteria.

The substituted benzoxazinone derivatives are already known for different biological activity.

Further the detail structure activity relationship studies are required along with the molecular manipulation i.e. molecular modeling may give better drugs and further toxicological study is needed. Molecules prepared for the biological testing do not always turn out as potential new molecules, but may be intended to serve as models for evaluation of the hypothesis.

#### ACKNOWLEDGEMENT

The Authors are thankful to the Management and Principal, Department of Chemistry, Vaageswari College of Pharmacy, Karimnagar, for extending support to carry out the research work. Finally, the authors express their gratitude to the Sura Labs, Dilsukhnagar, Hyderabad, for providing research equipment and facilities.

**BIBLIOGRAPHY**

- 1) Abraham, D.J., In *Burgers Medicinal Chemistry and Drug Discovery*, Vol 1, 5th Ed, John Wiley and Sons, 1995, 3-4.
- 2) Sugimoto Y, Otani T, Oie S, Wierzba K, Yamada Y. Mechanism of action of a new macromolecular antitumor antibiotic, C-1027. *J Antibiot* 1990; 43: 417-21.
- 3) Frechette R, Beach M. Ortho Pharm Corp. 2-Substituted amino and thioalkylbenzoxazine antimicrobial agents. WO Patent 9728167. 1997 Aug 7.
- 4) Frechette R, Weindner WMA. Ortho Pharm Corp. Benzoxazine antimicrobial agents. WO Patent 9717333. 1997 May 15.
- 5) Miles DH, Petrovna KO, Naser S, Yurjevich SS, Goun EA, Michailovich SV. 2-Oxo-1, 4-benzoxazine compounds for treatment of tuberculosis. US Patent 6,649,610. 2003 Nov 18.
- 6) Hori M, Watanabe I, Ohtaka H, Harada K, Maruo J, Morita T et al. Kanebo Ltd. 1,4-Benzoxazine derivative, pharmaceutical composition containing the same and use thereof. EP Patent 0719766. 2000 Apr 12.
- 7) Bhuniya D, Das SK, Madhavan GR, Iqbal J, Chakrabarti R. DR. Reddy's Lab Ltd. Benzoxazine and benzothiazine derivatives and pharmaceutical compositions containing them. WO Patent 03033481, 2003 Apr 24.
- 8) Fleury MB, LARGERON M, Lestage P, Lockhart B. Adir Et Compagnie. Novel 8-amino-1,4-benzoxazine compounds, preparation method and pharmaceutical compositions containing them. WO Patent 062889. 1999 Dec 9.
- 9) Holscher P, Jautelat R, Rehwinkel H, Jaroch S, Sulzle D, Hillmann M et al. Schering Aktiengesellschaft. Benzoxazine derivatives and benzothiazine derivatives having NOS-inhibitory and antioxidant properties. WO Patent 081324. 2001 Nov 1.
- 10) Hoelscher P, Rehwinkel H, Jaroch S, Sulzle D, Hillmann M, Burton GA et al. Schering Aktiengesellschaft. Benzoxazine and benzothiazine derivatives and their use in medicines. WO Patent 017173. 2000 Mar 30.
- 11) Berryman KA, Downing DM, Dudley DA, Edmunds JJ, Narasimhan LS, Rapundalo ST. Warner Lambert Co. Benzoxazinones/benzothiazinones as serine protease inhibitors. WO Patent 050257. 1999 Oct 7.
- 12) Bertani B, Borriello M, Bozzoli A, Bromidge SM, Granci E, Leslie C et al. Glaxo Group Ltd. Benzoxazinone derivatives, preparation thereof and uses in the treatment of CNS and other disorders. WO Patent 046124. 2004 Jun 3.
- 13) Belliotti T, Wise LD, Wustrow DJ. Warner Lambert Co. Benzoxazinone dopamine D4 receptor antagonists. WO Patent 045419. 1997 Dec 4.
- 14) Kikelj D, Suhadolc E, Rutar A, Pecar S, Puncuh A, Urleb U et al. Univerza V Ljubljani, Fakulteta Za Naravoslovje in Tehnologijo, Oddelek Za Farmacijo. Novel heterocyclic acyldipeptides, processes for the preparation thereof and pharmaceutical compositions containing the same. Eur Patent 0695308. 1998 Jul 1.
- 15) Stefanic P, Simoncic Z, Breznik M, Plavec J, Anderluh M, Addicks E et al. Conformationally tailored N-[(2-methyl-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-2-yl)carbonyl]proline templates as molecular tools for the design of peptidomimetics. Design and synthesis of fibrinogen receptor antagonists. *Org Biomol Chem* 2004; 2: 1511-7.
- 16) Anderluh PS, Anderluh M, Ilas J, Mravljak J, Dolenc MS, Stegnar M et al. Toward a novel class of antithrombotic compounds with dual function. Discovery of 1,4-benzoxazin-3(4H)-one derivatives possessing thrombin inhibitory and fibrinogen receptor antagonistic activities. *J Med Chem* 2005; 48: 3110-3.
- 17) R. H. R. White and O. D. Standen. Piperazine in the Treatment of Threadworms in Children (1953). *British Medical Journal*, 2, 755-757.
- 18) <http://www.drugbank.ca/drugs/DB00592>
- 19) M. Amelia Santos, Sergio M. Marques, Tiziano Tuccinardi, Paolo Carelli, Laura Panelli and Armando Rossello, *Bioorg. Med. Chem. Lett.*(2006), 14, 7539-7550.
- 20) Upadhayaya RS, Sinha N, Jain S, Kishore N, Chandra R, Arora SK, *Bioorg. Med. Chem.* 12, (2004), 2225.



**How to cite this article:**

**P. Aravinda Reddy. et al., DESIGN, SYNTHESIS AND BIOLOGICAL ACTIVITY OF 1, 4-BENZOXAZINONE DERIVATIVES AS ANTI-MICROBIAL AGENTS . J Pharm Res, 2017; 06(12): 244- 252 DOI:**

**Conflict of interest:** The authors have declared that no conflict of interest exists.

**Source of support:** Nil