

Hawking temperature and the quantum pressure of the Schwarzschild black hole

Kapil P. Chandra*

Department of physics and astronomy, University of Wuerzburg, D - 97074 Wuerzburg, Germany

Abstract

There is no term for pressure (PdV) in the first law of black hole thermodynamics. To address this question, we study the first law of black hole thermodynamics and derive an expression for it. We report that this pressure corresponds to the Hawking temperature and is inversely proportional to the quartic of the Schwarzschild radius. It implies that a lighter and smaller black hole exerts more pressure on its surrounding environment. It might shed light on the other thermodynamic aspects of the black hole.

Black holes are the most enigmatic objects in physics, and physicists are working hard to fully comprehend their behavior. The classical black hole was an object that absorbed everything and from which nothing, not even a photon, could escape.

Bekenstein and other physicists have recently made advances in this area, demonstrating that black holes have entropy. Hawking's contributions to this field have demonstrated that its temperature is inversely correlated with the black hole's surface gravity. In this context, the concept of quantum effect or quantum-gravitational effect has emerged [1] [2] [3] [4]. This work laid the foundation for the theory of black hole thermodynamics [5] where for a classical black hole (Schwarzschild black hole without charge and spin) it's first law states,

$$dM = TdS \quad (1)$$

here the black hole mass M corresponds to the total energy of the system, or enthalpy. T is the Hawking temperature, and S is the entropy. A remarkable aspect of this expression is that it

does not incorporate the pressure terms PdV , i.e. the corresponding change in volume V at pressure P , in contrast to the laws of thermodynamics of physical systems. It is important in the sense that it may help to understand the properties of a black hole by making an analogy with everyday things

This problem has been addressed, and it has been proposed that the variable cosmological constant might be the thermodynamic pressure in the first law of black hole thermodynamics [6] [7]. However, the question is, how does the cosmological constant incorporate itself into the particular individual black hole's pressure given that it is independent of the black hole's mass or other aspects? We thus consider the following fact: How do the fundamental properties of a black hole, such as mass, size, volume, and others, incorporated into the black hole's thermodynamical pressure? By deriving a mathematical expression for a black hole's pressure in terms of its attributes, we try to provide a solution to this intriguing question.

*kapil.chandra@gov.in

Now, as we discussed earlier, the Hawking temperature is an integral and basic part of the first law of black hole thermodynamics. And this temperature is correlated with the properties of a black hole. Furthermore, this temperature corresponds to the cosmological constant as written below [8],

$$\frac{R^2 c^7}{h G^2} = \frac{h c^5}{G^2 m^2} \quad (2)$$

where its first term is the mathematical expression of the cosmological constant [9], and RHS is Hawking radiation power without a numerical factor [2]. We wrote it in the form of force because the desired expression of thermodynamic pressure is simply force per area, so this representation would be convenient for us.

Now that it has been discussed, the cosmological constant corresponds to the pressure in the first law of thermodynamics [6] [7]. Therefore, if we set the first terms of Eq.(2) as pressure P for the black hole thermodynamics, However, by taking R as the Schwarzschild radius, the RHS will turn into the following,

$$P = \frac{h c^9}{G^4 m^4} \quad (3)$$

by taking Schwarzschild radius $R \sim \frac{Gm}{c^2}$ this Eq.(3) corresponds to,

$$P = \frac{h c}{R^4} \quad (4)$$

one can see that it denotes the pressure of the black hole; however, this expression contain the Planck constant thus we coined its name “quantum pressure of the black hole” for convenience, and hereafter we call it by this name in this paper. The origin of this quantum pressure might be the quantum effect of black holes or the quantum gravitational effect such as Hawking temperature.

It can be observed from Eq.(3) that the numerical amount of pressure is only depends on the mass of the black hole or Schwarzschild radius of black hole thus in only terms of mass, the pressure is inversely proportional to the quartic of mass, however, a lighter black hole

exerts more pressure on the surrounding environment.

From this description it is clear that the black hole’s pressure and energy density of vacuum is theoretically equivalent. Previously, this equivalence was discussed in refs [6] [7], and the possibility of similar kind of pressure in refs [10]. In previous work, we discussed it in a wider context [11]. However, its correspondence and congruence with the Hawking temperature and cosmological constants show it’s consistency with existing theories.

In summary, quantum pressure exists in the environment surrounding black holes. Theoretically, this quantum pressure corresponds to the Hawking temperature and vice versa. This quantum pressure is remarkable in that it is only a function of mass and is inversely proportional to the quartic of mass of the black hole; thus, a lighter black hole exerts more pressure on the environment. For the study of black hole thermodynamics, it might present a fresh perspective.

Acknowledgements: I would like to say thanks to the anonymous referee for his comment on the manuscript to make it better.

References

- [1] Jacob D. Bekenstein. Black holes and entropy. *Phys. Rev. D*, 7:2333–2346, Apr 1973. URL: <https://link.aps.org/doi/10.1103/PhysRevD.7.2333>, doi:10.1103/PhysRevD.7.2333.
- [2] S. W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43(3):199–220, 1975. doi:10.1007/BF02345020.
- [3] T Padmanabhan. Thermodynamical aspects of gravity: new insights. *Reports on Progress in Physics*, 73(4):046901, mar 2010. URL: <https://doi.org/10.1088/0034-4885/73/4/046901>, doi:10.1088/0034-4885/73/4/046901.

- [4] Igor Novikov. Black holes. In *Stellar Remnants*, pages 237–334. Springer, 1997.
- [5] J. M. Bardeen, B. Carter, and S. W. Hawking. The four laws of black hole mechanics. *Communications in Mathematical Physics*, 31(2):161–170, 1973. doi:10.1007/BF01645742.
- [6] Brian P Dolan. Pressure and volume in the first law of black hole thermodynamics. *Classical and Quantum Gravity*, 28(23):235017, nov 2011. URL: <https://doi.org/10.1088/0264-9381/28/23/235017>, doi:10.1088/0264-9381/28/23/235017.
- [7] David Kubizňák and Robert B. Mann. P - v criticality of charged AdS black holes. *Journal of High Energy Physics*, 2012(7), jul 2012. URL: [https://doi.org/10.1007/JHEP07\(2012\)033](https://doi.org/10.1007/JHEP07(2012)033), doi:10.1007/JHEP07(2012)033.
- [8] Kapil Chandra. Does Hawking predict the correct temperature of a black-hole? *Open Astron.*, 29(1):56–58, 2020. doi:10.1515/astro-2020-0008.
- [9] Ya. B. Zel’dovich, Andrzej Krasinski, and Ya. B. Zeldovich. The Cosmological constant and the theory of elementary particles. *Sov. Phys. Usp.*, 11:381–393, 1968. doi:10.1007/s10714-008-0624-6.
- [10] Xavier Calmet and Folkert Kuipers. Quantum gravitational corrections to the entropy of a schwarzschild black hole. *Phys. Rev. D*, 104:066012, Sep 2021. URL: <https://link.aps.org/doi/10.1103/PhysRevD.104.066012>, doi:10.1103/PhysRevD.104.066012.
- [11] kapil chandra. Anomalous nature of gravity at quantum realm. *open science foundation*, Sep 2019. doi:10.31219/osf.io/arns3.