



Short communication

Physicochemical analysis of *Terminalia catappa* (Almond) seed nuts grown in Benin city, Nigeria

Osaro Iyekowa* , Justina E. Ukpebor and Emmanuel Ehigimotor

Department of Chemistry, University of Benin, Benin City, Nigeria

*Author to whom correspondence should be addressed

Received: 26-12-2022, Revised: 09-01-2023, Accepted: 12-01-2023, Published: 31-03-2023

Copyright © 2023 Iyekowa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HOW TO CITE THIS

Iyekowa et al. (2023) Physicochemical analysis of *Terminalia catappa* (Almond) seed nuts grown in Benin City, Nigeria. *Mediterr J Pharm Pharm Sci*. 3 (1): 6-10. <https://doi.org/10.5281/zenodo.0000000>

Keywords: Minerals, Nigeria, proximate analysis, *Terminalia catappa*

Abstract: The need for natural medicine and nutrients from natural product sources cannot be overstressed. Plant parts like leaves, stem, root, seed, nut and flowers supply enormous minerals, phytochemicals and other bioactive constituents apart from the primary metabolites. *Terminalia Catappa* (*T. catappa*) is a combretaceous plant often found in tropical and subtropical regions. The plant has many uses as an ingredient of various drugs such as anti-oxidants, anti-inflammatory, anti-cancer, and anti-diarrheal. Thus, the main objective of this study was to determine the proximate, mineral and physicochemical analysis of *T. catappa* seed nuts. The proximate composition was determined according to the previous standard methods and elemental analysis by Atomic Absorption Spectrophotometry and Flame Photometer. Proximate composition revealed a moisture content of 02.34%, ash 03.89%, fibre content 08.50%, crude protein 29.66%, fat 21.98% and carbohydrate 33.63%. Elemental analysis of the plant have indicated the presence of iron, zinc, calcium, magnesium, potassium, sodium, copper and chromium in the aqueous extract. Physicochemical parameters of the plant revealed a specific gravity of 0.9182, viscosity 33.62 mm²/s, refractive index 1.4642, acid value of 0.102 mg KOH/g, saponification value of 151.28 mg KOH/g, iodine value 47.19 g I₂/100 g and peroxide value 8.93 meq/kg. In conclusion, the findings from this study confirmed that almond nut oil has a long shelf life which may explain its applications in pharmaceutical, non-pharmaceutical and personal care industries.

Introduction

Worldwide, fruits and sometimes fruit nuts of many plants have been consumed for their nutritional benefits but, apart from that, they could be of medical and pharmaceutical importance use to humans. *T. catappa* (*Terminalia catappa*, almond plant) from *Combretaceae* family is a large tree in the Leadwood

tree family that grows mainly in tropical regions of Africa, Asia and Australia [1]. In Southern Nigeria, it is referred to as “Ebelebo” among the Binis, “Baushe” in Hausa and “Igi-furutu” in Yoruba [2]. The juice obtained from its fresh leaves has been used in preparation of medicinal lotion which is

effective against scabies, leprosy, stomach ache and headache [3]. The tree provides several other dyes, tannins, timbers, carbohydrates, fuel plant and many other medicinal purposes [2]. *T. catappa* is a well-known herb in Ayurveda traditions in Indian. The ethno-medical uses of *T. catappa* includes modulatory activity of methanol extract of the leaves, wound healing activity of the chloroform extract of the bark [4], anticancer potency of the ethanol extract of the leaves [5] and anti-oxidant, hepato-protective, anti-microbial, analgesic, anti-inflammatory, anti-diabetic and anti-aging activity [3]. Recent scientific investigations revealed that the methanol extract of *T. catappa* leaf extract exhibits the dosage-dependent increase in inhibitory effect on α -glucosidase enzyme and α -amylase enzyme [6]. *T. catappa* exhibits anti-tumor activity with high anti-oxidant levels and the anti-oxidant defense may be due to the presence of phenolic and flavonoid components [5].

Phytochemical screening of the tree bark and leaves contains saponins, alkaloids, glycoside, terpenes, volatile oils, steroid and phenols. While, the seeds are rich in oleic-linoleic acid group, steroids, glycosides, phenolics and terpenes [7]. *T. catappa* has been observed to possess the anti-oxidant activity in a dose-dependent manner by DPPH assay, nirtic oxide assay, reducing power assay and H₂O₂ assay [8]. The plant also reverses the lipid levels to normal range and this shows anti-tumor and anti-lipidemic activities [9]. The leaves extract also inhibit the expression and activities of matrix metalloproteinase-9 (MMP-9) by the assessment of mRNA levels in hepato-cellular carcinoma [10]. This research is aimed at determining the proximate composition, mineral elements and physicochemical parameters of *T. catappa* seed extract.

Materials and methods

Materials: All the chemicals and materials used in this work were of analytical grade and were obtained from brand company of Merck and Sigma Aldrich, Modderfontein Johannesburg 1645, South Africa.

Sample collection and treatment: The fruits of *T. catappa* were collected from tree grown in Benin City, Edo State, Nigeria. The fresh epicarps of the *T. catappa* fruit were peeled to expose the seeds. The exposed seeds were then air dried for seven days and cracked open to collect the nuts. The seed nuts were air-dried under normal laboratory conditions for four weeks and finally pulverized. The powder seed nuts were then used for the proximate analysis and the determination of mineral composition [11].

Proximate analysis: The proximate composition of the seed kernel was analyzed following standard methods [11], crude fat by Soxhlet method and moisture by vacuum oven, crude protein by Kjeldahl method and ash by ignition. Carbohydrate content was calculated by subtracting the values of all the other proximate analysis from 100 [11].

Mineral element determination: For the elemental analysis of the powder seeds, one gram of the sample was digested with concentrated HNO₃ and HClO₄. The sample was then filtered and made up to 50 ml with distilled water [11]. The metal concentrations were determined using an Atomic Absorption Spectrophotometer (Buck Scientific model 210) and a Flame Spectrophotometer (Sherwood, model 410). The mineral elements investigated were Iron (Fe), Zinc (Zn), Calcium (Ca), Magnesium (Mg), Potassium (K), Nickel (Ni), Sodium (Na), Cadmium (Cd), Cupper (Cu), Chromium (Cr) and Lead (Pb).

Physico-chemical analysis: The physicochemical analysis including acid value, saponification value, peroxide value and iodine value were determined according to standard methods [1].

Results

The results of the proximate analysis, mineral elements and physicochemical analysis of the seed oil extract of *T. catappa* seed nut are shown in **Tables 1, 2** and **3**, respectively. Thus, a wide range of percentage among the constituents of *T. catappa* seed nut was found as shown in **Table 1**. Thus, carbohydrate, fat and crude protein were found in high percentage (33.3, 21.9 and 29.6% respectively),

whereas, fibre, ash and moisture contents were found in very low percentage (08.5, 03.8 and 02.3%, respectively).

Table 1: Results of proximate composition of *T. catappa* seed nut

Parameter	Value (%)
Moisture	02.34 ± 0.2
Ash	03.89 ± 0.1
Fibre content	08.50 ± 0.1
Crude protein	29.66 ± 0.3
Fat	21.98 ± 0.1
Carbohydrate	33.63 ± 0.2

In **Table 2**, different contents of mineral elements of the extract are given. Potassium, magnesium and zinc contents represent the highest, whereas, other elements are presented in a very low contents. However, lead (Pb) content was not detected in this extract.

Table 2: Mineral elements detected in hexane extract of *T. catappa* seed nut

Element	Content (mg/kg)
Fe	06.40 ± 0.00
Zn	39.00 ± 0.42
Ca	29.00 ± 2.60
Mg	30.00 ± 3.13
K	54.7 ± 1.50
Ni	0.00 ± 0.00
Na	2.10 ± 0.18
Cd	0.00 ± 0.00
Cu	0.30 ± 0.01
Cr	0.18 ± 0.01
Pb	0.00 ± 0.00

Table 3 shows acid, saponification, iodine and peroxide values of the crude oil hexane extract of *T. catappa* seed nut. A large variation was found among them with saponification value represents the high value and acid value was the lowest value.

Table 3: Chemical parameters of the crude oil hexane extract of *T. catappa* seed nut

Parameter	Mean ± S.E.
Acid value [mg KOH/g of oil]	0.102 ± 0.01
Saponification value [mg KOH/g of oil]	151.28 ± 2.63
Iodine value [g I ₂ /100 g of oil]	47.19 ± 1.44
Peroxide value [meq/kg of oil]	8.93 ± 0.86

Discussion

In this study, the moisture content obtained for the almond seed nuts was to be 02.34%. Moisture content helps to determine storage time. Low moisture content is important for storage quality and shelf life of seeds. The moisture content indicates that almond seed can thus be kept for a long time. Furthermore, most researches have inferred that low moisture in seeds reduces the microbial activities and decreases unwarranted fermentation [1]. The lower moisture content recorded may be due to the fact that the almond seeds were air dried for 28 days. The ash content of a sample is an indication of the level of minerals present [12]. The values of ash content reported from different locations in Nigeria appear to fall within a similar range of 02.00% - 05.00%. According to the previous report [13], low ash content of seeds could be credited to the fact that during maturation, inorganic ions migrate from different parts of the plant to the region of active growth. This value was also lower than those obtained for almond seeds from other countries like Ivory Coast (04.60%) but higher than 03.78% reported in the seeds from Malaysia [14]. The crude fiber content obtained in this work was 08.50% and higher than 03.11% obtained by Akpabio [12]. The high amount of protein content obtained in this work suggests that almond nut can be used to supplement other dietary sources. Therefore, it is recommended that almond nut be added to the meals of individuals in order to increase the protein content. From other study, the seed contains carbohydrates, protein, fat,

fiber, iron, ascorbic acid, arachidic acid and β -carotene are in good proportion [15].

Amongst the minerals determined, the concentration of potassium was the highest (54.7 mg/kg) and Cr had the least quantity (0.18 mg/kg) while Ni, Cd and Pb were beyond detection limit. Iron had a quantity of 6.40 mg per kg. On comparison with the work of Mandloi [15], phosphorus, potassium, niacin, riboflavin and thiamin were also detected in the seed. Therefore, almond seed could be recommended as a dietary supplement for people who need essential minerals and iron. The World health Organization (WHO) recommended daily allowance for zinc is 11 mg. Therefore, almond nut when properly processed will be a good source for zinc due to its high value. Potassium had a quantity of 54.7 mg per kg which plays a role in many body functions including transmission of nerve signals, muscle contractions, fluid balance and various chemical reactions.

The chemical parameters of the oil extract shown below in this study indicated an acid value of 0.102 mg KOH per g of oil which is attributable to its low free fatty acid value of 0.051. This value was low when compared to 0.787 mg KOH per g of almond nut oil obtained from India [12]. However, Ogbeide and others [16] reported a much higher level acid value of 2.94 mg KOH per g. The saponification value obtained showed a value of 151.278 mg KOH per g which less than the value 326.08 mg KOH per

g is obtained [16]. The saponification value obtained in this work could suggest that the almond seed has a few proportions of low molecular weight fatty acids and the difference observed could be as a result of method of extraction of the oil from the seed. Other report, however, indicated that almond nut oil contains higher proportions of low molecular weight fatty acids [1]. Rancidity begins to occur in oil when the peroxide value ranges from 20.0 mg per g oil to 40.0 mg per g oil. The peroxide value obtained in this work was 8.93 meq per kg and this low values of peroxide value are indicative of low levels of oxidative rancidity of the oils and suggest high levels of antioxidants [1]. It has been asserted that oils with iodine value less than 100 mg I₂ per 100 g are non-drying oils, and consequently, the lesser the number of unsaturation the lower the susceptibility of such oil to oxidative rancidity [17]. The iodine value obtained in this work was 47.186 g per 100 g which is less than 131.37 g per 100 g obtained by [16] and 121.19 g per 100 g obtained by [1]. The iodine value obtained was less than 100 g per 100 g, thus, almond nut oil cannot be placed in the class of drying oil.

Conclusion: *Terminalia catappa* seed nuts contain essential minerals required for nutritional and medicinal purposes, and the low moisture content of the oil indicates that products or formulations from the nuts will possess longer shelf life.

Author contribution: OI conceived, designed the study, performed the analysis and drafted the manuscript. JEU & EE collected and analyzed the data. All authors have approved the final version of the manuscript and agreed to be accountable for its contents.

Conflict of interest: The authors declare absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Ethical issues: Including plagiarism, informed consent, data fabrication or falsification and double publication or submission have completely been observed by authors.

Data availability statement: The raw data that support the findings of this article are available from the corresponding author upon reasonable request.

References

1. Akpakpan AE, Akpabio UG (2012) Evaluation of proximate composition, mineral element and anti-nutrient in almond (*Terminalia catappa*) seeds. *Research Journal of Applied Sciences*. 7 (9): 489-493. doi: 10.3923/rjasci.2012.489.493.
2. Aigbokhan EI (2014) Annotated checklist of vascular plants of Southern Nigeria- a quick reference guide to the vascular plants of southern Nigeria: a systematic approach. Uniben Press, Benin City. doi: 10.13140/RG.2.1.1604.0808.
3. Anand AV, Divya N, Kotti PP (2015) An updated review of *Terminalia catappa*. *Pharmacognosy Reviews*. 9 (18): 93-98. doi: 10.4103/0973-7847.162103.
4. Khan AA, Kumar V, Singh BK, Singh R (2014) Evaluation of wound healing property of *Terminalia catappa* on excision wound models in Wistar rats. *Drug Research (Stuttg)*. 64 (5): 225-228. doi: 10.1055/s-0033-1357203.
5. Pandya NB, Tigari P, Dupadahalli K, Kamurthy H, Nadendla RR (2013) Antitumor and antioxidant status of *Terminalia catappa* against Ehrlich ascites carcinoma in Swiss albino mice. *Indian Journal of Pharmacology*. 45 (5): 464-469. doi: 10.4103/0253-7613.117754.
6. Divya N, Vijaya A (2014) Phytochemical investigation and in vitro anti-diabetic activity of *Terminalia catappa* leaves. *International Journal of Phytopharmacy*. 4 (5): 132-134. doi.org/10.7439/ijpp.v4i5.111.
7. Mininel FJ, Leonardo-Junior CS, Espanha LG, Resende FA, Varanda EA, Leite COF, Vilegas W, Dos Santos LC (2014) Characterization and quantification of compounds in the hydroalcoholic extract of the leaves from *Terminalia catappa* Linn. (Combretaceae) and their mutagenic activity. *Evidence-Based Complementary and Alternative Medicine*. 67-69. 2014: 676902. doi: 10.1155/2014/676902.
8. Kotti PP, Anand AV (2014) Phytochemical analysis and in vitro antioxidant activity of *Terminalia catappa*. *World Journal of Pharmaceutical Sciences*. 2 (11): 1495-1498.
9. Naitik P, Prakash T, Kotresha D, Rao NR (2012) Effect of *Terminalia catappa* on lipid profile in transplanted fibrosarcoma in rats. *Indian Journal of Pharmacology*. 44 (3): 390-392. doi: 10.4103/0253-7613.96345.
10. Yeh CB, Yu YL, Lin, CW, Chiou HL, Hsieh MJ, Yang SF (2014) *Terminalia catappa* attenuates urokinase-type plasminogen activator expression through Erk pathways in hepatocellular carcinoma. *BMC Complementary and Alternative Medicine*. 14: 141. doi: 10.1186/1472-6882-14-141.
11. Association of Official Analytical Chemists AOAC (2015) Official methods of analysis of the association of official analytical chemists, 5th Ed. Association of official analytical chemists, Virginia, USA. ISBN: 0-935584-42-0.
12. Atsu-Barku VY, Nyarko HD, Dordunu P (2012) Studies on the physicochemical characteristics, microbial load and storage stability of oil from Indian almond nut (*Terminalia catappa*). *Food Science and Quality Management*. 8: 9-17. Corpus ID: 73648051.
13. Monnet YT, Gbogouri A, Koffi, PKB, Kouamé LP (2012) Chemical characterization of seeds and seed oils from mature *Terminalia catappa* fruits harvested in Côte d'Ivoire. *International Journal of Bioscience*. 2: 110-124. Corpus ID: 86147763.
14. Siew NG, Ola L, Muhammad KS, Hussain N, Sulaiman R (2015) Physicochemical properties of Malaysian-grown tropical almond nuts (*Terminalia catappa*). *Journal of Food Science and Technology*. 52 (10): 6623-6630. doi: 10.1007/s13197-015-1737-z.
15. Mandloi S, Mishra R, Varma R, Varughese B, Tripathi J (2013) A study on phytochemical and antifungal activity of leaf extracts of *Terminalia catappa*. *International Journal of Pharm and BioScience*. 4: 1385-1393. Scopus ID: 13835580.
16. Ogbeide OK, Eze OF, Akaeze DA, Akhigbe IU, Omoruyi U, Iyekowa O, Owolabi BJ (2021) Physico-chemical properties, chemical composition, biodiesel production and antibacterial potential of *Terminalia catappa* seed oil. *ChemSearch Journal*. 12 (2): 70-80.
17. Aremu MO, Ibrahim H, Bamidele TO (2015) Physicochemical characteristics of the oils extracted from some Nigerian plant foods - A review. *Chemical and Process Engineering Research*. 32: 36-52.