

Active afforestation of drained peatlands is not a viable option under the EU Nature Restoration Law

Gerald Jurasinski¹, Kenneth Byrne², Bogdan H. Chojnicki³, Jesper Riis Christiansen⁴, Vytas Huth⁵, Hans Joosten¹, Radosław Juszczak³, Sari Juutinen⁶, Åsa Kasimir⁷, Leif Klemedtsson⁷, Wiktor Kotowski⁸, Ain Kull⁹, Mariusz Lamentowicz¹¹, Amelie Lindgren⁷, Rita Linkevičienė¹², Annalea Lohila¹³, Ülo Mander⁹, Michael Manton¹⁴, Kari Minkkinen¹⁵, Jan Peters¹⁶, Florence Renou-Wilson¹⁷, Jūratė Sendžikaitė^{18,19}, Rasa Šimanauskienė^{12,20}, Franziska Tanneberger²¹, Rudy van Diggelen²², Harri Vasander¹⁵, David Wilson²³, Dominik H. Zak²⁴, John Couwenberg¹

¹Peatland Science, University of Greifswald, partner in the Greifswald Mire Centre, Greifswald, Germany

²Faculty of Science and Engineering, University of Limerick, Limerick, Ireland

³Laboratory of Bioclimatology, Poznan University of Life Sciences, Poznan, Poland

⁴Department of Geoscience and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark

⁵Faculty of Agricultural and Environmental Sciences, University of Rostock, Rostock, Germany

⁶Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

⁷Department of Earth Sciences, University of Gothenburg, Gothenburg, Sweden

⁸Department of Plant Ecology and Environmental Conservation, University of Warsaw, Warsaw, Poland

⁹Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

¹¹Climate Change Ecology Research Unit, Adam Mickiewicz University, Poznań, Poland

¹²Laboratory of Climate and Water Research, Nature Research Centre, Vilnius, Lithuania

¹³Institute for Atmospheric and Earth System Research, University of Helsinki, Finland

¹⁴Faculty of Forest Science and Ecology, Vytautas Magnus University, Akademija, Lithuania

¹⁵Department of Forest Sciences, University of Helsinki, Finland

¹⁶Michael Succow Foundation, partner in the Greifswald Mire Centre, Greifswald, Germany

¹⁷School of Biology and Environmental Science, University College Dublin, Dublin, Ireland

¹⁸Laboratory of Flora and Geobotany, Institute of Botany, Nature Research Centre, Vilnius, Lithuania

¹⁹Foundation for Peatland Restoration and Conservation, Vilnius, Lithuania

²⁰Institute of Geosciences, Faculty of Chemistry and Geosciences, Vilnius university, Vilnius, Lithuania

²¹Greifswald Mire Centre, Greifswald, Germany

²²Geobiology Research Group, Department of Biology, University of Antwerp, Belgium

²³Earthy Matters, Ireland

²⁴Institute of Ecoscience, Aarhus University, Aarhus, Denmark

Executive Summary

Several EU Member States argue that active afforestation of degraded peatlands should be recognized as a restoration measure under the Nature Restoration Law (NRL). In this policy brief, we discuss the scientific evidence on the greenhouse gas fluxes of peatlands under forestry and its limitations, uncertainties and evidence gaps. In our opinion:

- Afforestation of drained peatlands, while maintaining their drained state, is not equivalent to ecosystem restoration. This approach will not restore the peatland ecosystem's flora, fauna, and functions.
- Currently, there is insufficient evidence to support the long-term climate change mitigation benefits of active afforestation of drained peatlands.
- Most studies only focus on the short-term gains in standing biomass and rarely explore the full life cycle emissions associated with afforestation of drained peatlands. Thus, it is unclear whether the CO₂ sequestration of a forest on drained peatland can offset the carbon loss from the peat over the long term.
- In some ecosystems, such as abandoned or certain cutaway peatlands, afforestation may provide short-term benefits for climate change mitigation compared to taking no action. However, this approach violates the concept of sustainability by sacrificing the most space-effective carbon store of the terrestrial biosphere, the long-term peat store, for a shorter-term, less space-effective, and more vulnerable carbon store, namely tree biomass.
- Consequently, active afforestation of drained peatlands is not a viable option for climate mitigation under the EU Nature Restoration Law.
- To restore degraded peatlands, hydrological conditions must first be improved, primarily through rewetting.

1. Introduction

The EU Nature Restoration Law is critical in reaching climate mitigation targets, as the most recent IPCC report very clearly shows (IPCC AR6 SYR 2023). The carbon dioxide (CO₂) sinks necessary to reach climate neutrality and then net cooling during the second half of this century will, for the most part, rely on the land use sector (LULUCF). Sinks should not be used to compensate for avoidable sources and emission reduction remains the utmost priority.

Several EU member states pursue the recognition of active afforestation of drained and degraded peatlands (without rewetting) as a measure under the Nature Restoration Law (NRL). This is, however, problematic from a scientific perspective. Firstly, afforestation of drained peatlands, while keeping them drained, will not restore the peatland ecosystem with its flora, fauna and functions. Secondly, as long as a peatland remains drained, it will degrade further, carbon stored in the peat is lost to the atmosphere, and downstream systems may be severely polluted with nutrients and dissolved organic matter (Zak & McInnes 2022). Typically, much more carbon is stored in the peat than in the forest biomass (Tanneberger et al. 2021), and long-term losses from the drained soil will likely be larger than storage in the forest biomass (Dunn & Freeman 2011), and the biomass carbon will be lost back to the atmosphere if a forest is used for production purposes.

Presently, there is little evidence to support the claim that afforestation on drained peatlands could be beneficial for climate change mitigation in the long-term. In most cases, CO₂ release from peat

soil degradation will likely exceed carbon sequestration in forest biomass when considering full growth cycles, as was also concluded by IPCC in the 2013 Wetlands Supplement (IPCC 2014). In contrast, ecosystem-scale flux measurements show that rewetting/restoration¹ of forestry-drained peatlands can reduce soil CO₂ emissions and even restore the CO₂ sink function within a few years/decades (Hambley et al. 2019).

Some recent studies suggesting afforestation as the best possible restoration measure (*e.g.*, Butlers et al. 2021, Butlers et al. 2022a/b, Samariks et al. 2023) either did not measure all relevant greenhouse gases and carbon fluxes or only covered very short time periods. Potential lateral losses of C as dissolved organic or inorganic carbon and as particulate organic carbon are rarely considered but can be substantial (Billett et al. 2004). To provide reliable data, measurement periods should cover the entire land use cycle—from site preparation to sowing or planting, growth of the biomass, thinning or ditch cleaning, through to harvest, the fate of the biomass (whether used for long-lived products or not) and the fallow time before renewed site preparation. For cropping agriculture, this cycle commonly takes a single year, in forestry it takes much longer. Longer, comprehensive greenhouse gas monitoring studies are currently only available from the boreal region and even here, the authors call for longer-term studies that include the entire life cycle (*e.g.*, Bjarnadottir et al. 2021). Such studies do not exist to date and so we can only approach the true climate balance of peatland forestry by using space-for-time substitution and stitching together studies that cover different stages of the forestry life cycle.

Given the lack of conclusive evidence, it would be wrong to accept active afforestation of drained peatlands as a viable option for climate mitigation under the NRL. This is all the more true given additional negative effects that may arise from peatland forestry: Forestry on drained peatlands further lowers the water table by interception and increased transpiration of the trees, is more susceptible to wildfires, which are becoming more frequent and severe due to climate change and droughts in the boreal zone leading to increased carbon losses (*e.g.*, Zhao et al. 2021, Zheng et al. 2023, Liu et al. 2023), and also negatively impacts water storage capacity, water quality, and nutrient runoff, leading to greater variations in stream flow and water quality for downstream aquatic ecosystems (*e.g.*, Evans et al. 2016, Flynn et al. 2022).

In the following, we discuss the role of peatlands in relation to the climate with respect to forestry-drained peatlands in particular.

2. Peatlands and climate

Plants absorb CO₂ and store the carbon in their biomass. When they die, they are decomposed and the CO₂ is released again. In intact peatlands—called mires—water saturation of the soil effectively excludes oxygen, inhibiting the full decomposition of the dead plant material, which then accumulates as peat. In this way, peatlands have sequestered huge amounts of carbon over thousands of years. Globally, peatlands store approximately 600 Gt of carbon (Yu et al. 2010), which is more than is contained in global forest above-ground biomass (Santoro et al. 2021), and have cooled the planet by approximately 0.6°C over the past 10,000 years (Frolking & Roulet 2007, Joosten et al. 2016). Forests and peatlands are fundamentally different in terms of C cycling over

¹ *Rewetting* and *restoration* are often used interchangeably but they are not. *Rewetting* describes the deliberate action of raising the water table on drained soils to re-establish water saturated conditions. In contrast, *restoration* refers to the full re-establishment of all ecosystem functions which cannot be done directly but only be aimed at by taking facilitating measures like, for instance, rewetting, removing degraded topsoil, re-introducing mire-typical vegetation, etc.

time, in that pristine mature forests are generally in balance, whereas peatlands continue to accumulate carbon year after year.

To understand the climate impact of peatlands, it is necessary to briefly review the relevant processes. While intact peatlands (mires) sequester carbon from the atmosphere, incomplete decomposition under water-saturated oxygen-free conditions results in the production of methane (CH₄). The amount of CH₄ released from intact, wet peatlands varies strongly depending on environmental conditions like pH, temperature and dominant vegetation (*e.g.*, Lai 2009). Methane is a short-lived greenhouse gas. On average, it stays in the atmosphere for less than 12 years. If there is a steady emission of CH₄, a dynamic equilibrium will establish over time in which in a certain year exactly as much CH₄ disappears from the atmosphere as is added, and the CH₄ concentration in the atmosphere and the climate impact do not increase any further (Frolking & Roulet 2007). Natural, undrained peatlands almost always release CH₄, but the net uptake of CO₂ overcompensates the CH₄ losses in the long-term. In mires, *i.e.*, wet peat-forming peatlands, formation and emission of nitrous oxide (N₂O, a potent greenhouse gas) is negligible.

When peatlands are drained, the upper soil layers are no longer water saturated, oxygen enters the peat, and decomposition of organic matter becomes much more efficient, leading to mineralisation of the peat and, thus, high CO₂ (Ojanen et al. 2010, Jovani-Sancho et al. 2018) and N₂O emissions (IPCC 2014, Klemetsson et al. 2005, Leppelt et al. 2014, Minkkinen et al. 2020). In contrast, CH₄ production and emissions decrease because of water table drawdown, while drainage ditches may remain a major source of CH₄ emissions (Minkkinen et al. 1997, Köhn et al. 2021, Rissanen et al. 2023). Thus, draining undisturbed peatlands reduces soil CH₄ emissions but increases the soil CO₂ and N₂O emissions. In the case of forestry, drainage also increases tree stand carbon stock, until the stand is cut. This may lead to short-term cooling, but in most cases, the long-term effect will be warming (Laine et al. 1996, Ojanen & Minkkinen 2020).

In order to determine the climate-effectiveness of rewetting drained peatlands, we need to compare CO₂, CH₄, and N₂O emissions and lateral losses of C and N both in the drained and the rewetted situations. Rewetting of drained peatlands is always a choice between continued emissions of (a) long-lived GHGs (CO₂ and N₂O) and (b) a short-lived GHG (CH₄). In the short term, rewetting will cause a warming impact because of increased CH₄ emissions, but in the long-term (decades to centuries) the result is cooling (Ojanen and Minkkinen 2020, Wilson et al. 2016/2022, Günther et al. 2020). Especially for cutover peat bogs (Rocheffort et al. 2003, González & Rocheffort 2014) but also after top-soil removal on formerly drained grassland on bog peat (Huth et al. 2022) it has been shown that re-introduction of mire-typical vegetation can speed up the process of restoration including a strong reduction of total GHG emissions after rewetting.

3. Forestry on peatlands

Peatland forestry is concentrated in the boreal zone, but also occurs across the temperate climate zone. It is common in many European countries, mainly in the north (*i.e.*, Finland, Sweden, and Norway), but is also of national importance in the Baltic countries, the United Kingdom, Ireland, Poland and Germany (see Figure 1, data for non-EU countries are not included). Peatlands used for forestry are typically drained, which leads to simultaneously increasing CO₂ emissions through peat decomposition (see above) and CO₂ sequestration through the growing tree stand.

4. Available evidence regarding peatland forestry and climate

Although soil CO₂ emissions increase after drainage, several studies on drained boreal peatland forests, made with micrometeorological or combined chamber-efflux-litter-production-methods, suggest that carbon sequestration in the tree biomass can exceed the carbon loss from the decomposition of the peat (Lindroth et al. 2007, Meyer et al. 2013, Ojanen et al. 2013; Hommeltenberg et al. 2014, Uri et al. 2017, Minkkinen et al. 2018, Bjarnadottir et al., 2021). In most cases, however, soil C stocks decrease over time, which is the deciding factor when whole rotation climate impacts are considered. Nutrient-poor sites in the boreal zone may accumulate C in soils (Ojanen et al. 2013), but in the absence of a high water table, the fate of this C is unclear over a production cycle.

The most common way to measure greenhouse gas fluxes in treeless peatlands has been to place airtight chambers on the soil and to measure the change in the concentration of gases inside the chamber. This commonly applied chamber technique cannot be used in a forest stand as the trees do not fit inside the chambers. Therefore, the soil carbon balance has been estimated by subtracting measured litter production from heterotrophic soil respiration (*e.g.*, Ojanen et al. 2012, 2013, Jovani-Sancho et al. 2021, Uri et al. 2017) where the latter has been measured from trenched plots in which plants and tree roots have been killed. Simply placing chambers on the forest soil will not provide reliable measurements of fluxes from soil degradation, because the roots of the trees emit CO₂ as well (*i.e.*, autotrophic soil respiration). Intricate chamber set-ups are required to distinguish between emissions from the soil (*i.e.*, heterotrophic soil respiration) and from the living tree roots (*i.e.*, autotrophic respiration) (Mäkiranta *et al.* 2008, Hermans et al. 2022).

The eddy covariance technique (EC), which uses highly sophisticated, fast measuring gas analysers, requires large flat and homogeneous areas that are rarely available also because forestry areas on peatlands are often organised in fairly small blocks. EC is the standard method for ecosystem-level measurements. The technique has a typical accuracy of about $\pm 20\%$ (SEM) and is technically challenging. The method also allows for the estimation of the soil C sink/source by subtracting the biomass increment from the measured net ecosystem exchange. This can be done by biomass measurements and modelling, which needs accuracy, but is doable. Other soil C balance estimation methods include the estimation of soil subsidence through pollen or C isotope profiles or by consecutive thickness measurements (Minkkinen et al. 1999, Hooijer et al., 2012, Simola et al. 2012, Sloan et al., 2019). These integrate longer time periods. Getting accurate estimates is, thus, challenging.

The above-mentioned studies consider forestry on drained peatlands, which is based on natural tree stands in the boreal zone. In the temperate zone, sites have more often been drained for agriculture or peat extraction and only later been afforested. Most of the studies do not describe the situation after afforestation of agricultural fields or cutover peatlands, but this difference in soil management history may lead to different outcomes. In addition, the above-mentioned gas exchange studies only give a temporary GHG balance for the study period. ***They do not consider the whole forestry cycle, which includes harvesting of the wood (i.e., the removal of sequestered biomass C), decomposition of wood products, and the time needed for stand regeneration. Harvested sites, especially clear-cut sites are large C sources*** (Korkiakoski *et al.* 2019; Korkiakoski *et al.* 2023), until a new stand has been regenerated. In addition, only a small part of wood products are long-lived, most of the C in wood products is lost to the atmosphere in a few years after cuttings (Soimakallio et al. 2016).




Figure 1: Areal share of peatlands under agriculture, peat extraction, forestry, and other uses in countries of the European Union. Colours refer to different land use types, the size of the circles reflects total peatland area. Map prepared by Cosima Tegetmeyer.

One recent, relatively comprehensive and well-measured study from Iceland (Bjarnadottir et al. 2021) showed no negative effects of peatland forestry compared to rewetted or wet peatlands. The study area, a poorly drained site, was afforested with a very productive species (Black cottonwood, *Populus trichocarpa*). According to the authors, only further long-term studies and life-cycle assessments will show whether forestry on drained peatlands can really be more climate friendly than wet and healthy peatlands. A recent meta-analysis on the climate effects of forestry on *shallow* organic soils (<40-50cm peat depth) in Scotland showed no negative climate effect (Vanguelova et al. 2018). However, meta-analyses on the effects of forestry on *deep* peat soils in the temperate climate region showed considerable net emissions of CO₂ to the atmosphere (Hommeltenberg et al. 2014, Jovani-Sancho et al. 2021). The latter finding confirms the emission factors derived in the 2013 IPCC Wetland Supplement (IPCC 2014). The considerably lower figures for carbon loss from the soil in Hermans et al. (2022) show that there can be strong variation across sites suggesting that more research on the matter is needed. With respect to the harvesting stage of peatland forestry, it

was shown that selective cutting instead of clear-cutting can lower GHG emissions in a forestry drained peatland in the boreal climate zone (Korkiakoski *et al.* 2023). Although the site with partial harvest transformed into a CO₂ sink five years after harvest, peat decomposition continued, releasing almost the same amount of carbon into the atmosphere as was fixed by the trees. Most studies in drained peatland forests show similar results: the ecosystem may be a C sink, whereas the soil is a C source. Therefore, production forestry, where biomass C is harvested and rapidly lost back to the atmosphere (Soimakallio *et al.* 2016), will likely result in net C losses in the long run. Thus, ***none of the studies discussed above provides a basis to include active afforestation on drained peatlands, especially if managed for production forestry, as a viable option under the NRL.***

Some recent studies from Latvia do seem to support the idea that afforestation of drained peatlands could be better for the climate than rewetting. However, these studies are inconclusive, have major flaws and biases and cannot be verified and validated because the methods used are error prone and descriptions often lack clarity. For instance, Samariks *et al.* (2023) claim that they show that afforestation of peat extraction sites can result in net carbon removals. However, they did not measure all elements of the carbon cycle. Using chamber measurements, they failed to clearly distinguish between soil emissions and emissions from tree roots. Instead of analysing, they simply *assumed* that the soil fluxes made up less than half of the measured total flux, independent of changes in management or site conditions, which is unrealistic. Moreover, the appropriate comparison would be with a rewetted peat extraction site. The study of Bārdule *et al.* (2022) suggests that emissions from wet peatlands are not lower than in drained peatlands. However, they measured only once per month over a period of only four months, during only one vegetation period. Their experimental set-up allowed only the CO₂ flux going out of the system (soil and litter degradation plus plant respiration) to be measured, but not the sequestration of CO₂ from the atmosphere during photosynthesis. Moreover, they neglected to account for the carbon export from the drained sites. ***Results of studies that only address parts of the GHG exchange in afforested peatlands are not appropriate to draw conclusions on whether afforestation of drained peatlands is the best choice for the climate.***

In another example, Butlers *et al.* (2021) report larger N₂O and CH₄ emissions from ‘naturally wet’ sites than from ‘drained’ forests. These findings are not surprising considering that their ‘naturally wet’ site was only slightly less dry (water tables more than 60 cm below the soil surface in summer!). Again, a full GHG balance can only be assessed when CO₂ exchange is included as well. The same authors claim to have shown that CO₂ emissions from ‘naturally wet’, nutrient-rich organic forest soils can be larger than those from drained sites (Butlers *et al.* 2022a). However, they again used a technique that fails to depict net-CO₂ exchange appropriately, as it excludes photosynthesis by the ground vegetation and lacks the quantification of CO₂ release from the tree roots. The authors state that the “study results demonstrate that drainage does not have significant effect on CO₂ emissions”, a conclusion not supported by their data. ***Without inclusion of the contribution of the trees and without a full life cycle assessment, no sensible conclusions on the climate effect of wet vs. drained forested peatlands can be drawn.***

Butlers *et al.* (2022a) attempt to get a better understanding of the whole GHG balance by looking at different stages of the harvest cycle and even including the input of litter. Again, the same error-prone methods are used. No distinction is made between root- and soil derived emissions in the measurements, but instead a regression equation is used that deems slightly more than half of the measured flux to be related to decomposition of litter and soil. The used equation presents a

broad relationship that may be helpful to constrain large scale estimates, but it was not made to infer site-specific flux values, as was stressed by the original authors (Bond-Lamberty et al. 2004). Again, the conclusions are based on a very short time series of measurements. This falls far short of a full life cycle assessment that is essential to provide reliable data on the effects of forest growth on drained peatlands in comparison to the situation in rewetted or pristine peatlands. In addition, there were no measurements conducted in intact or rewetted peatlands to provide an appropriate baseline for comparison. Thus, ***the study results are unsuitable to support far reaching generalisations regarding afforestation of peatlands as viable options to achieve climate goals under the NRL.***

In addition to GHG exchange, also the change in albedo and the release of aerosols as well as the lateral exchange of carbon and nitrogen add to the total climate effect of ecosystems (Billett et al. 2010). Increased tree cover decreases albedo compared to a treeless mire (Lohila et al. 2010), which leads to local warming (Gao et al. 2014). In contrast, forests are large sources of biogenic volatile organic compounds (BVOCs) and thus may have a considerable cooling impact on the climate (Tunved et al. 2006). In the boreal zone, this impact is similar in magnitude, but opposite to that of albedo (Kalliokoski et al. 2020).

Most importantly, however, the felling of trees and the subsequent fate of the carbon sequestered in the wood needs to be considered (Ciais et al. 2008). As little as 50% of actual tree biomass may be extracted during harvest, the remainder is left to decompose on site and within a few years returns as CO₂ to the atmosphere (Leturq 2020; Korkiakoski et al. 2019). After the tree is felled, a fair amount of its carbon can be stored in wood products (there will always be some parts lost during production). Yet, a simple carbon balance does not tell the whole story; unlike the carbon stock in soil and peat, not everything made out of wood is climate neutral (Leturq 2020). As a rule, for a long-lasting product made out of 80 year old wood to be climate cooling, the wood should not be discarded and burnt for at least 40 years after harvest. A product made out of 40 year old wood would need to remain for at least 20 years after harvest (Guest et al. 2013; Galimshina et al. 2022). Long-lived harvested wood products are actually quite rare, and harvesting does reduce the amount of total carbon stored in the forest (Soimakallio et al. 2022). ***In assessing the GHG balance of peatland forestry, complete harvest cycles should be taken into account. Such data are simply absent at the moment.***

Since trees grow slowly, no measurements of full growth cycles are yet available and results spanning longer time periods are derived from the investigation of chronosequences. Certainly, more chronosequence work is needed, but measurements must be made over multiple years so that variations in weather and other environmental conditions can be integrated into the models. As Vanguelova et al. (2018) have expressed: “There is a clear need for long term studies using different planting ages (chronosequence studies) to ensure robust results when evaluating the impacts of afforestation and restocking on soil carbon stocks, as short-term impact studies are likely to provide misleading conclusions.”

Finally, it has been shown that peatland forestry on drained sites is more prone to wildfires (Kohlenberg et al. 2018), which will become more frequent and severe in times of climate change with more frequent and more intense droughts in the boreal zone (Walker et al. 2019). Boreal forests in North America have turned from a net sink to a net source of GHG in recent years (Zhao et al. 2021), primarily due to more frequent and more severe fires (Zheng et al. 2023). In addition to the carbon loss from burnt wood, substantial carbon losses from burnt and burning peat layers should be considered including the water-born carbon losses (Liu et al. 2023). The severity of peatland fires might also be high because in countries where the density of forest roads is relatively

high (e.g., in Finland), large drained peatland areas may usually still have lower density of roads compared to upland areas, making fires more difficult to be controlled. Having said that, Finland's land use sector seems to have gradually transitioned from being a CO₂ sink to a source already (Statistics Finland 2022b). Siljander et al. (2022) suggest that the fastest way to strengthen carbon sinks in Finland is to reduce logging and the Finnish nature panel recommends, among other things, the rewetting and restoring of wetlands and peatlands (Lång et al. 2022).

Aside from impacts to emissions, forestry on peatland may have negative impacts on water storage capacity, water quality and nutrients runoff, including loss of organic matter via fluvial pathways, which is subsequently mineralised and the carbon partially returned to the atmosphere (Evans et al 2016). It has to be mentioned, that also rewetting of drained peatlands can lead to considerable amounts of N and P leaching (e.g., Koskinen et al. 2017) but nutrient loss will decrease once the system is biogeochemically stable again. The loss in water storage capacity due to the loss of pore space caused by drainage can manifest itself in greater variations in stream flow and water quality for downstream aquatic ecosystems (Flynn et al. 2022). Intense rainfall events are predicted to increase under climate warming, which can cause mid-term flooding leading to the die-back of trees caused by increased hydrological instability. ***Indeed, research to evaluate the trade-off values of wood production, carbon sequestration and emissions and water storage of natural, drained and rewetted peatlands is urgently required*** (Stachowicz et al. 2021).

5. Conclusion

The most recent IPCC report (IPCC AR6 SYR 2023) outlines in clear language: We have to get to grips with proper natural climate solutions since they are our only option to avoid the gravest consequences of climate change and global warming. Therefore, we need an effective Nature Restoration Law (NRL) in the EU. ***The many open questions and lack of evidence for overall climate benefits of active afforestation on peatlands prohibit its inclusion as a viable climate change mitigation measure in the NRL.*** In addition, the NRL should foster true natural ecosystems wherever possible, and a forest that can only grow when the peat below the trees is drained does not comply with this requirement. ***Afforestation of drained peatlands is not restoration but afforestation. We cannot restore a peatland ecosystem, its flora, its fauna, its functions, by afforestation. The only solution to restore a peatland ecosystem is rewetting.*** The reintroduction of natural mire vegetation can potentially speed up the restoration process. If trees belong to such an ecosystem, they will regenerate naturally.

6. References

- Artz RRE, Chapman SJ, Saunders M, Evans CD, Matthews RB (2013) Comment on "Soil CO₂, CH₄ and N₂O fluxes from an afforested lowland raised peat bog in Scotland: implications for drainage and restoration" by Yamulki et al. *Biogeosciences* 10:7623–7630
- Bārdule A, Gerra-Inohosa L, Kļaviņš I, Kļaviņa Z, Biteniēks K, Butlers A, Lazdiņš A, Lībiēte Z (2022). Variation in the mercury concentrations and greenhouse gas emissions of pristine and managed hemiboreal peatlands. *Land* 11:1414
- Billett MF, Charman DJ, Clark JM, Evans CD, Evans MG, Ostle NJ, Worrall F, Burden A, Dinsmore KJ, Jones T, McNamara NP, Parry L, Rowson JG, Rose R (2010) Carbon balance of UK peatlands: current state of knowledge and future research challenges. *Climate Research* 45:13–29

- Bjarnadottir B, Sungur GA, Sigurdsson BD, Kjartansson BT, Oskarsson H, Oddsdottir ES, Gunnarsdottir GE, Black A (2021). Carbon and water balance of an afforested shallow drained peatland in Iceland. *Forest Ecology and Management* 482:118861
- Bond-Lamberty B, Wang C, Gower ST (2004) A global relationship between the heterotrophic and autotrophic components of soil respiration? *Global Change Biology* 2004(10):1756–1766
- Butlers A, Bārdule A, Spalva G, Muižnieks E (2021). N₂O and CH₄ emissions from naturally wet and drained nutrient-rich organic forest soils. In *Proceedings of the International Scientific Conference “Rural Development”* 196–200
- Butlers A, Lazdiņš A (2022) Case study on greenhouse gas (GHG) fluxes from flooded former peat extraction fields in central part of Latvia. *Research for Rural Development* 37:44–49
- Butlers A, Lazdiņš A, Kalēja S, Bārdule A (2022b) Carbon budget of undrained and drained nutrient-rich organic forest soil. *Forests* 13:1790
- Butlers A, Spalva G, Licite I, Purvina D (2022a) Carbon dioxide (CO₂) emissions from naturally wet and drained nutrient-rich organic forest soils. In *Proceedings of the 21st International Scientific Conference ‘Engineering for Rural Development’*, Jelgava, Latvia 25–27
- Ciais P, Schelhaas MJ, Zaehle S, Piao SL, Cescatti A, Liski J, Luysaert S, Le-Maire G, Schulze E-D, Bouriaud O, Freibauer A, Valentini R, Nabuurs GJ (2008) Carbon accumulation in European forests. *Nature Geoscience* 1:425–429
- Davidson SJ, Goud EM, Franklin C, Nielsen SE, Strack M (2020) Seismic line disturbance alters soil physical and chemical properties across boreal forest and peatland soils. *Frontiers in Earth Sciences* 8:281
- Dunn C, Freeman C (2011) Peatlands: our greatest source of carbon credits? *Carbon Management* 2:289–301
- Flynn R, Mackin F, McVeigh C, Renou-Wilson F (2021) Impacts of mature forestry plantation on blanket peatland runoff regime and water quality. *Hydrological Processes* 36:e14494
- Frolking S, Roulet NT (2007) Holocene radiative forcing impact of northern peatland carbon accumulation and methane emissions. *Global Change Biology* 13:1079–1088
- Galimshina A, Moustapha M, Hollberg A, Padey P, Lasvaux S, Sudret B, Habert G (2022). Bio-based materials as a robust solution for building renovation: A case study. *Applied Energy* 316:119102
- Gao Y, Markkanen T, Backman L, Henttonen HM, Pietikäinen J-P, Mäkelä HM, Laaksonen A (2014) Biogeophysical impacts of peatland forestation on regional climate changes in Finland. *Biogeosciences* 11:7251–7267
- González E, Rochefort L (2014) Drivers of success in 53 cutover bogs restored by a moss layer transfer technique. *Ecological Engineering* 68:279–290
- Guest G, Cherubini F, Strømman AH (2013) Global warming potential of carbon dioxide emissions from biomass stored in the anthroposphere and used for bioenergy at end of life. *Journal of Industrial Ecology* 17:20–30
- Günther A, Barthelmes A, Huth V, Joosten H, Jurasinski G, Koebsch F, Couwenberg J (2020). Prompt rewetting of drained peatlands reduces climate warming despite methane emissions. *Nature communications* 11:1644
- Hambley G, Andersen R, Levy P, Saunders M, Cowie NR, Teh YA, Hill TC (2019) Net ecosystem exchange from two formerly afforested peatlands undergoing restoration in the Flow Country of northern Scotland. *Mires and Peat* 2018/19:05
- He H, Jansson PE, Svensson M, Björklund J, Tarvainen L, Klemedtsson L, Kasimir Å (2016) Forests on drained agricultural peatland are potentially large sources of greenhouse gases – insights from a full rotation period simulation. *Biogeosciences* 13:2305–2318
- Hermans R, McKenzie R, Andersen R, Teh YA, Cowie N, Subke JA (2022) Net soil carbon balance in afforested peatlands and separating autotrophic and heterotrophic soil CO₂ effluxes. *Biogeosciences* 19:313–327
- Hommeltenberg J, Mauder M, Drösler M, Heidbach K, Werle P, Schmid HP (2014) Ecosystem scale methane fluxes in a natural temperate bog-pine forest in southern Germany. *Agricultural and Forest Meteorology* 198:273–284
- Hooijer A, Page S, Jauhiainen J, Lee WA, Lu XX, Idris A, Anshari G (2012) Subsidence and carbon loss in drained tropical peatlands. *Biogeosciences* 9:1053–1071

- Huth V, Günther A, Bartel A, Gutekunst C, Heinze S, Hofer B, Jacobs O, Koebsch F, Rosinski E, Tonn C, Ullrich K, Jurasinski G (2022) The climate benefits of topsoil removal and Sphagnum introduction in raised bog restoration. *Restoration Ecology* 30:e13490
- IPCC (2014) 2013 *Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands*. IPCC, Switzerland
- IPCC AR6 SYR (2023) *Synthesis report of the IPCC sixth assessment report (AR6) – Summary for Policymakers*
- Joosten H, Clarke D (2002). *Wise use of mires and peatlands*. International Mire Conservation Group and International Peat Society, 304
- Joosten H, Sirin A, Couwenberg J, Laine J, Smith P (2016) The role of peatlands in climate regulation. In: Bonn A, Allot T, Evans M, Joosten H, Stoneman R (eds) *Peatland Restoration and Ecosystem Services: Science, Policy and Practice*. Cambridge University Press, Cambridge 63–76
- Jovani-Sancho AJ, Cummins T, Byrne KA (2021). Soil carbon balance of afforested peatlands in the maritime temperate climatic zone. *Global Change Biology* 27:3681–3698
- Kalliokoski T, Bäck J, Boy M, Kulmala M, Kuusinen N, Mäkelä A, Minkkinen K, Minunno F, Paasonen P, Peltoniemi M, Taipale D, Valsta L, Vanhatalo A, Zhou L, Zhou P, Berninger F (2020) Mitigation impact of different harvest scenarios of Finnish forests that account for albedo, aerosols and trade-offs of carbon sequestration and avoided emissions. *Frontiers in Forests and Global Change* 3:562044
- Kasimir Å, He H, Coria J, Nordén A (2018) Land use of drained peatlands: Greenhouse gas fluxes, plant production, and economics. *Global Change Biology* 24:3302–3316
- Kasimir Å., He H, Jansson PE, Lohila A, Minkkinen K (2021) Mosses are important for soil carbon sequestration in forested peatlands. *Frontiers in Environmental Science* 9:680430
- Klemetsson L, von Arnold K, Weslien P, Gundersen P (2005) Soil CN ratio as a scalar parameter to predict nitrous oxide emissions. *Global Change Biology* 11:1142–1147
- Kohlenberg AJ, Turetsky MR, Thompson DK, Branfireun BA, Mitchell CPJ (2018) Controls on boreal peat combustion and resulting emissions of carbon and mercury. *Environmental Research Letters* 13:035005
- Köhn D, Welpelo C, Günther A, Jurasinski G (2021) Drainage ditches contribute considerably to the CH₄ budget of a drained and a rewetted temperate fen. *Wetlands* 41:71
- Korkiakoski M, Ojanen P, Penttilä T, Minkkinen K, Sarkkola S, Rainne J, Laurila T, Lohila A (2020) Impact of partial harvest on CH₄ and N₂O balances of a drained boreal peatland forest. *Agricultural and Forest Meteorology* 295:108168
- Korkiakoski M, Ojanen P, Tuovinen JP, Minkkinen K, Nevalainen O, Penttilä T, Aurela M, Laurila T, Lohila A (2023). Partial cutting of a boreal nutrient-rich peatland forest causes radically less short-term on-site CO₂ emissions than clear-cutting. *Agricultural and Forest Meteorology* 332:109361
- Korkiakoski M, Tuovinen JP, Penttilä T, Sarkkola S, Ojanen P, Minkkinen K, Rainne J, Laurila T, Lohila A (2019). Greenhouse gas and energy fluxes in a boreal peatland forest after clear-cutting. *Biogeosciences* 16:3703–3723
- Koskinen M, Tahvanainen T, Sarkkola S, Menberu MW, Laurén A, Sallantausta T, Marttila H, Ronkanen A-K, Parviainen M, Tolvanen A, Koivusalo H, Nieminen M (2017) Restoration of nutrient-rich forestry-drained peatlands poses a risk for high exports of dissolved organic carbon, nitrogen, and phosphorus. *Science of The Total Environment* 586:858–869
- Lång K, Aro L, Assmuth A, Haltia E, Hellsten S, Larmola T, Lempinen H, Lindfors L, Lohila A, Miettinen A, Minkkinen K, Nieminen M, Ollikainen M, Ojanen P, Sarkkola S, Sorvali J, Seppälä J, Tolvanen A, Vainio A, Wall A, T V (2022). Turvemaiden käytön vaihtoehdot hiilineutraalissa Suomessa (In Finnish). *Suomen ilmastopaneelin raportti2/2022*, Helsinki, Finland.
- Lepilinen D, Laurén A, Uusitalo J, Fritze H, Laiho R, Kimura B, Tuittila E-E (2021). Response of vegetation and soil biological properties to soil deformation in logging trails of drained boreal peatland forests. *Canadian Journal of Forest Research* 52:511–526

- Leppelt T, Dechow R, Gebbert S, Freibauer A, Lohila A, Augustin J, Drösler M, Fiedler S, Glatzel S, Höper H, Järveoja J, Lærke PE, Maljanen M, Mander Ü, Mäkiranta P, Minkkinen K, Ojanen P, Regina K, Strömgren M (2014) Nitrous oxide emission budgets and land-use-driven hotspots for organic soils in Europe. *Biogeosciences* 11:6595–6612
- Leturcq P (2020). GHG displacement factors of harvested wood products: the myth of substitution. *Scientific reports* 10:20752
- Lindroth A, Lund M, Nilsson M, Aurela M, Christensen TR, Laurila T, Rinne J, Riutta T, Sagerfors J, Ström L, Tuovinen J-P, Vesala T (2007) Environmental controls on the CO₂ exchange in north European mires. *Tellus B: Chemical and Physical Meteorology* 59:812–825
- Liu H, Zak D, Zableckis N, Cossmer A, Langhammer N, Meermann B, Lennartz B (2023) Water pollution risks by smoldering fires in degraded peatlands. *Science of the Total Environment* 871:161979
- Lohila A, Minkkinen K, Laine J, Savolainen I, Tuovinen J-P, Korhonen L, Laurila T, Tietäväinen H, Laaksonen A (2010) Forestation of boreal peatlands — impacts of changing albedo and greenhouse gas fluxes on radiative forcing. *Journal of Geophysical Research* 115:G04011
- Mäkiranta P, Minkkinen K, Hytönen J, Laine J (2008). Factors causing temporal and spatial variation in heterotrophic and rhizospheric components of soil respiration in afforested organic soil croplands in Finland. *Soil Biology and Biochemistry* 40:1592–1600
- Meyer A, Tarvainen L, Noursratpour A, Björk RG, Ernfors M, Grelle A, Kasimir Klemetsson Å, Lindroth A, Rantfors M, Rütting T, Wallin G, Weslien P, Klemetsson L (2013) A fertile peatland forest does not constitute a major greenhouse gas sink. *Biogeosciences* 10:7739–7758
- Minkkinen K, Laine J, Nykänen H, Martikainen P (1997) Importance of drainage ditches in emissions of methane from mires drained for forestry. *Canadian Journal of Forest Research* 27:949–952
- Minkkinen K, Ojanen P, Koskinen M, Penttilä T (2020) Nitrous oxide emissions of undrained, forestry-drained, and rewetted boreal peatlands. *Forest Ecology and Management* 478:118494
- Minkkinen K, Ojanen P, Penttilä T, Aurela M, Laurila T, Tuovinen JP, Lohila A (2018) Persistent carbon sink at a boreal drained bog forest. *Biogeosciences* 15:3603–3624
- Minkkinen K, Vasander H, Jauhiainen S, Karsisto M, Laine J (1999) Post-drainage changes in vegetation composition and carbon balance in Lakkasuo mire, Central Finland. *Plant and Soil* 207:107–120
- Ojanen P, Minkkinen K (2020) Rewetting offers rapid climate benefits for tropical and agricultural peatlands but not for forestry-drained peatlands. *Global Biogeochemical Cycles* 34:e2019GB006503
- Ojanen P, Minkkinen K, Alm J, Penttilä T (2010) Soil–atmosphere CO₂, CH₄ and N₂O fluxes in boreal forestry-drained peatlands. *Forest Ecology and Management* 260:411–421
- Ojanen P, Minkkinen K, Lohila A, Badorek T, Penttilä T (2012) Chamber measured soil respiration: a useful tool for estimating the carbon balance of peatland forest soils? *Forest Ecology and Management* 277:132–140
- Ojanen P, Minkkinen K, Penttilä T (2013). The current greenhouse gas impact of forestry-drained boreal peatlands. *Forest Ecology and Management* 289:201–208
- Poyda A, Reinsch T, Kluß C, Loges R, Taube F (2016) Greenhouse gas emissions from fen soils used for forage production in northern Germany. *Biogeosciences* 13:5221–5244
- Renou-Wilson F, Farrell EP (2007) Phosphorus in surface runoff and soil water following fertilization of afforested cutaway peatlands. *Boreal Environment Research* 12:693–709
- Rissanen AJ, Ojanen P, Stenberg L, Larmola T, Anttila J, Tuominen S, Minkkinen K, Koskinen M, Mäkipää R (2023) Vegetation impacts ditch methane emissions from boreal forestry-drained peatlands—Moss-free ditches have an order-of-magnitude higher emissions than moss-covered ditches. *Frontiers in Environmental Science* 11:1121969
- Rocheffort L, Quinty F, Campeau S, Johnson K, Malterer T (2003) North American approach to the restoration of Sphagnum dominated peatlands. *Wetlands Ecology and Management* 11:3–20

- Samariks V, Lazdiņš A, Bārdule A, Kalēja S, Butlers A, Spalva G, Jansons Ā (2023). Impact of Former Peat Extraction Field Afforestation on Soil Greenhouse Gas Emissions in Hemiboreal Region. *Forests* 14:184
- Santoro M, Cartus O, Carvalhais N, Rozendaal D, Avitabile V, Araza A, de Bruin S, Herold M, Quegan S, Rodríguez-Veiga P, Balzter H, Carreiras J, Schepaschenko D, Korets M, Shimada M, Itoh T, Martínez AM, Cavlovic J, Gatti RC, da Conceição Bispo P, Dewnath N, Labrière N, Liang N, Lindsell J, Mitchard ETA, Morel A, Pacheco Pascagaza AM, Ryan CM, Slik F, Vaglio Laurin G, Verbeeck H, Wijaya A, Willcock S (2021) The global forest above-ground biomass pool for 2010 estimated from high-resolution satellite observations. *Earth System Science Data* 13:3927–3950
- Siljander, R., Cederlöf, M, Skoglund K (2022) Climate annual report 2022. Ministry of the Environment, Helsinki, Finland.
- Simola H, Pitkänen A, Turunen J (2012) Carbon loss in drained forestry peatlands in Finland, estimated by re-sampling peatlands surveyed in the 1980s. *European Journal of Soil Science* 63:798–807
- Sloan TJ, Payne RJ, Anderson AR, Gilbert P, Mauquoy D, Newton AJ, Andersen R (2019) Ground surface subsidence in an afforested peatland fifty years after drainage and planting. *Mires and Peat* 23:6
- Soimakallio S, Böttcher H, Niemi J, Mosley F, Turunen S, Hennenberg KJ, Reise J, Fehrenbach H (2022) Closing an open balance: The impact of increased tree harvest on forest carbon. *Global Change Biology Bioenergy* 14:989–1000
- Soimakallio S, Saikku L, Valsta L, Pingoud K (2016) Climate change mitigation challenge for wood utilization—the case of Finland. *Environmental Science & Technology* 50:5127–5134
- Stachowicz M, Manton M, Abramchuk M, Banaszuk P, Jarašius L, Kamocki A, Povilaitis A, Samerkhanova A, Schäfer A, Sendžikaitė J, Wichtmann W, Zableckis N, Grygoruk M (2022) To store or to drain — To lose or to gain? Rewetting drained peatlands as a measure for increasing water storage in the transboundary Neman River Basin. *Science of The Total Environment* 829:154560
- Statistics Finland (2022) National Inventory Report under the UNFCCC and the Kyoto Protocol. 581 p.
- Statistics Finland (2022b) Greenhouse gas emissions in 2021 became revised – the land use sector was confirmed a source of emissions. Environment and Nature. Statistics Finland, Helsinki, Finland.
- Tanneberger F, Appulo L, Ewert S, Lakner S, Ó Brolcháin N, Peters J, Wichtmann W (2021). The Power of Nature-Based Solutions: How Peatlands Can Help Us to Achieve Key EU Sustainability Objectives. *Advanced Sustainable Systems* 5:2000146
- Tunved P, Hansson H-C, Kerminen V-M, Ström J, Dal Maso M, Lihavainen H, Viisanen Y, Aalto PP, Komppula M, Kulmala M (2006) High natural aerosol loading over boreal forests. *Science* 312:261–263
- Uri V, Kukumägi M, Aosaar J, Varik M, Becker H, Morozov G, Karoles K (2017) Ecosystems carbon budgets of differently aged downy birch stands growing on well-drained peatlands. *Forest Ecology and Management* 399:82–93
- Vanguelova E, Chapman S, Perks M, Yamulki S, Randle T, Ashwood F, Morison J (2018) Afforestation and restocking on peaty soils—new evidence assessment. *ClimateXChange*, Scotland
- Walker XJ, Baltzer JL, Cumming SG, Day NJ, Ebert C, Goetz S, Johnstone JF, Potter S, Rogers BM, Schuur EAG, Turetsky MR, Mack MC (2019) Increasing wildfires threaten historic carbon sink of boreal forest soils. *Nature* 572:520–523
- Wilson D, Blain D, Couwenberg J, Evans CD, Murdiyarso D, Page SE, Renou-Wilson F, Rieley JO, Sirin A, Strack M, Tuittila E-S (2016) Greenhouse gas emission factors associated with rewetting of organic soils. *Mires and Peat* 17:04
- Wilson D, Mackin F, Tuovinen JP, Moser G, Farrell C, Renou-Wilson F (2022) Carbon and climate implications of rewetting a raised bog in Ireland. *Global Change Biology* 28:6349–6365
- Yu Z, Loisel J, Brosseau DP, Beilman DW, Hunt SJ (2010). Global peatland dynamics since the Last Glacial Maximum. *Geophysical Research Letters* 37:L13402

- Zak D, McInnes R (2022) A call for refining the peatland restoration strategy in Europe. *Journal of Applied Ecology* 59:2698–2704
- Zhao B, Zhuang Q, Shurpali N, Köster K, Berninger F, Pumpanen J (2021) North American boreal forests are a large carbon source due to wildfires from 1986 to 2016. *Scientific reports* 11:7723
- Zheng B, Ciais P, Chevallier P, Yang H, Canadell JG, Chen Y, van der Felde IR, Aben I, Chuvieco E, Davis SJ, Deeter M, Hong C, Kong Y, Li H, Li H, Lin X, He, K, Zhang Q (2023) Record-high CO₂ emissions from boreal fires in 2021. *Science* 379:912–917