

Silver Diamine Fluoride in Pediatric Dentistry: SDF Review article

Dr. Bibhav Dubey

Post graduate student
Department of pediatric and preventive Dentistry
Babu Banarasi Das University, Lucknow

Dr. Neerja Singh

Professor
Department of pediatric and preventive Dentistry
Babu Banarasi Das University, Lucknow

Dr. Monika Rathore

Professor and Head
Department of pediatric and preventive Dentistry
Babu Banarasi Das University, Lucknow

Dr. Subash Singh

MDS, Associate Professor
Department of pediatric and preventive Dentistry
Babu Banarasi Das University, Lucknow

Dr. Malika Agarwal

MDS, Consultant pedodontist
Lucknow

Keywords: SDF, Silver Diamine Fluoride, magic bullet, nanobullet, miracle antidote

Author Declaration

Conflict of interest: The author declares that there no conflict of interest regarding the publication of this paper.

Funding statement: Nil

Abstract: In young children, untreated tooth decay can cause discomfort, low self-esteem, weight loss, sleeping difficulties, and loss of space, resulting in severe crowding and misalignment of succedaneous teeth. Apart from these concerns, treating young children is a difficult chore for dentists because of behavioral issues that make treatment more difficult. Considering these facts, an innovative anti-caries agent known as silver diamine fluoride has gained popularity among dentists. The aforementioned agent has antibacterial properties that perform both to prevent and arrest tooth decay. The noninvasive approach is the treatment of choice among dentists for dealing with uncooperative children since it is simple and quick to use. It can be used as preferred therapeutic agent for those patients where aerosol production is contraindicated.


Introduction:

Early childhood caries (ECC) is one of the most common diseases affecting deciduous dentition, despite continual advances in disease genesis and prevention efforts, it continues to have a negative influence on the health of young children, with social and economic consequences.

ECC occurs in young children as a result of a number of risk factors, including malnutrition, low socioeconomic status, prolonged bottle feeding, frequent snacking, iron and vitamin D deficiency, and untreated ECC has additional consequences, including an increased risk of caries, loss of succedaneous teeth, pain, weight loss, low self-esteem, missed school hours, costly emergency treatment, and limited growth and development of jaw bone. As a result, untreated carious lesions and underutilization of dental treatments have emerged as two of the most urgent public health issues confronting children in developing nations. Further, traditional restoration techniques are time-consuming and difficult, requiring child cooperation for a successful outcome. Pre-cooperative children are not ideal candidates for the same situation. Understanding the complexities of the condition and circumstances, a unique miracle panacea with anti-cariogenic properties has been developed which has proven to be a boon for both pediatric and general dentists.

Various fluoride-based prevention strategies and remineralizing agents are advocated to prevent dental caries in children. Among these caries arresting methods, topical Silver Diamine Fluoride (SDF) has recently gained enormous popularity among dentists worldwide due to its non-invasive, low-cost, easy-to-apply success in stopping and preventing the progression of dental caries.

History:



“Figure: 1. Summary of history “

Composition:

The fluoride content of SDF solutions vary based on the manufacturer and brand. The most usually utilized SDF concentration is 38%.^[8]

Silver Diamine Fluoride (At 38% concentration)					
S.No.	Components	ppm	W/v (%)	Properties	pH
1	Silver	255,000	24.4-28.8	Antimicrobial	10
2	Ammonia		7.5-11	Stabilizes high concentration of solution	
3	Fluoride	44,800	5.0-5.9	Antimicrobial & Remineralization	

“Table: 1. Composition of SDF most commonly used in concentration 38%”

Commercially Available Brand Names:

SDF (%) Product brand Manufacturer Country Product image

12 &30 Cariestop Biodinamica Brazil



Figure: 2. Cariestop

38 Saforide Toyo SeiyakuKasei Japan



Figure: 3. Saforide

38 Advantage Arrest Elevate Oral Care United States



Figure 4: Advantage Arrest

38 e-SDF Kids-e-dental,mumbai India



Figure 5: e-SDF

38 FAgamine Tedequim SRL Argentina



Figure 6: FAgamine

30-35 Riva Star SDI Dental Ltd Australia




Figure 7. Riva Star

“Table: 2.Commercially available brand names of SDF in different countries with concentration in percentage”

Physical properties:

- It is a colorless, odorless aqueous solution containing silver ions, ammonium and fluoride ions. [9]
- It is highly alkaline (pH = 10).
- IUPAC name Diamminesilver (I) fluoride
- Formula - AgF (NH₃)₂
- Molar mass - 160.927643g/mol [10]
- 3D model (JSmol)



“Figure: 8. 3D model of Silver diammine fluoride.”

Chemical misnomer:

Since 1969, the SDF compound has been incorrectly spelled or mispronounced as "Ammoniacal silver fluoride," "Silver fluoride diamine," and so on, even though the fact that the correct nomenclature term is silver diammine fluoride, which contains "ammine" groups (-NH3) rather than "amine" groups (NH2). "Ammine" refers to a chemical species in which one or more ammonia molecules (NH3) are bonded in a coordination complex with a metal ion." The nomenclature term "Diamine" has achieved universal approval and is now used in both scientific and promotional contexts.^[11]


Indications:

- Patients at a high risk of dental caries, For example, Xerostomia, salivary dysfunction, cancer therapy, and dental anxiety.^[12]
- Pre-cooperative children whose behavior limits invasive restorative treatment to avoid restorative care under general anesthesia or sedation.^[13]
- When aerosol generating procedures are unable to be performed. For example, COVID_19, Asthma, etc.
- Patient without access to dental care.^[12]
- Teeth that has no sign and symptoms of pulpitis.
- Active carious lesion on the root surface.^[14]
- Carious primary teeth that exhibit signs of exfoliation on radiographs.^[15]
- MIH (Molar incisor hypomineralization) to relieve the symptoms of dentin hypersensitivity.^[16]

Contraindications:

- Patient allergic to silver.^[17]
- Patient with oral soft tissue ulcers, for examples, Desquamative gingivitis, Stomatitis.^[12]
- Patient with thyroid medication, pregnancy, known allergy to potassium or iodide.^[13]
- If parents or guardians refuse to use SDF because they are concerned about color changes.
- It is not advisable to use where isolation of tooth or oral tissue is not possible.
- Clinical sign and symptoms of irreversible pulpitis.^[18, 19]
- Radiographic sign of pulp involvement or peri-apical pathology.^[13]

Mechanism of action of Silver Diamine Fluoride^{[20-23]:}



“Figure: 9. Flowchart representation of mechanism action of SDF.”

USE OF SDF IN CLINICAL PRACTICE:**SDF in Caries Arrest of Primary Dentition:**

SDF is known for its anti-cariogenic properties, but it has been employed in a variety of applications in the literature. The table 3 below presents summary of published studies on caries arrest in children by SDF.

Author	Study period months	Study design	Dentition studied	Study groups (sample size)	CA (%)	Application	Follow-up visits months
Chu et al. 2002 ^[24]	30	PCCT	Primary anterior	38% SDF (641) 5% NaF (576) No treatment (273)	65 41 34	Annual	6, 12, 18, 24, 30
Llodra et al. 2005 ^[25]	36	RCT	Primary canine, molars, & PFMs	38% SDF (675) No treatment (658)	85 62	Semi-annual	6, 12, 18, 24, 30
Braga et al (2009)	30	pilot RCT	first molar	cross-toothbrushing technique(22) 10% SDF(22) GIC(22)	No significant in all groups	Semi -annual	3, 6, 12, 18, and 30
Yee et al. 2009 ^[26]	24	RCT	Primary	38% SDF (3,396) 12% SDF (1,652) No treatment (1,590)	31 22 15	Single application	6, 12, 24
Zhi et al. 2012 ^[27]	24	RCT	Primary	38% SDF (218) 38% SDF (239) GIC (262)	91 79 82	Annual Semi-annual Annual	6, 12,
dos Santos et al. 2012 ^[28]	12	RCT	Primary	30% SDF (183) GIC (162)	67 39	Single application	12
Duangthip et al. 2016 ^[29]	18	RCT	Primary	30% SDF (458) 30% SDF (426) 5% NaF (523)	40 35 27	Annual Single application Single application	6, 12,18
Duangthip et al. 2018 ^[30]	30	RCT	Primary	30% SDF (377) 30% SDF (367) 5% NaF (484)	48 33 34	Annual Once a week for 3 weeks Once a week for 3 weeks	5, 12, 18, 30
Fung et al. 2018 ^[31]	30	RCT	Primary	12% SDF (927) 12% SDF (987) 38% SDF (971) 38% SDF (905)	55.2 58.6 66.9 75.7	Annual Semi-annual Annual Semi-annual	6, 12, 18, 24, 30

PCCT, prospective controlled clinical trial; RCT, randomized clinical trial; PFMs, permanent first molars; NaF, 5% sodium fluoride varnish; GIC, glass ionomer cement; CA, caries arrest; M, month

“Table: 3. Summary of published studies on caries arrest in children

Root canal irrigant:

Author	Year	Study	SDF (%)	Conclusion
Hiraishi et al. ^[32]	2010	In vitro	3.8%	“They reported that 3.8% SDF showed 100% efficiency against E. faecalis after a direct 60-min exposure.”
Mathew et al. ^[33]	2012	Ex vivo	3.8%	“Both 3.8% silver diamine fluoride and 2% chlorhexidine showed a superior capacity to sterilize the root canals than control groups.”
Ebtissam & their colleagues. ^[34]	2019	In vitro	3.8%	“SDF possessed higher antimicrobial activity than 2% CHX against E. faecalis biofilms.”
Minavi et al. ^[35]	2021	In vitro	3.8%	“They demonstrated that SDF possesses antimicrobial properties against the opportunistic pathogen E. faecalis. Moreover, using a dentin model the substantivity of 3.8% SDF is significantly greater than 6.25% NaOCl, but is comparable to 2% CHX.”
Maru et al. ^[36]	2022	RCT	3.8%	“A randomized, controlled clinical trial was performed that included primary molars with pulp necrosis. After analyzing samples before and after irrigation in the control group (NaOCl), they found a strong significant decrease of bacterial load. The same occurred in the 3.8% SDF group samples. SDF (Experimental) group was superior to control group.”

“Table: 4. Summary of published studies utilizing SDF as root can irrigant”

Dentine hypersensitivity:


Kiesow A. and colleagues (2022) ^[37] “conducted an in vitro study in which they compared 38% SDF gel to non-viscous commercially available SDF. Human root surface dentin specimens were treated with gelled or conventional 38% SDF or negative control. Penetration depths of up to 500 m were found for both SDF formulations. Both formulations occluded dentinal tubules in the same way. Precipitates were observed on the dentin surface and within dentinal tubules for both SDF formulations, with the experimental gel SDF product being slightly more abundant than the commercially available one. In terms of dentinal tubule penetration and occlusion, the 38% SDF gel formulation was indistinguishable from the commercial 38% SDF product.”

Indirect pulp capping agent:

S.No.	Studies (Authors, Years)	Study Type	Results /Conclusion
1	L de F(2011) ^[38]	In vivo	“Both glass ionomer and SDF can be potential IPC material.”
2	Gupta et al (2011) ^[39]	Ex vivo	“The calcium hydroxide group had the highest increase in calcium and phosphate ion levels. Fluoride ion levels increased significantly in the SDF and GC VII groups. The samples treated with GC VII showed the highest increase in microhardness. The SDF group had the highest zone of bacterial inhibition. Both SDF and GC VII can serve as excellent IPC materials.”
3.	Sinha et al (2011) ^[40]	In vivo	“The percentage of calcium levels increased about equally in the GC VII and Ca (OH) 2 groups, followed by the SDF group. The GC VII group had the highest percentage rise in phosphate ions, followed by the SDF group and the Ca (OH) 2 group. The GC VII group had the highest percentage of fluoride increase, followed by the SDF group and the Ca (OH) 2 group. Both SDF and GC VII can serve as excellent IPC materials.”
4	Shah A. & their co-worker (2020) ^[41]	In vivo	“SDF was evaluated as an indirect pulp capping agent in primary teeth, and the result showed that clinical and radiographic success at one month was 100% in SDF and 93.75% in calcium hydroxide, with no significant difference between the two groups. The study found that SDF can be used as a viable alternative for calcium hydroxide in primary teeth for IPC.”
5	Shafi N.& Colleagues (2022) ^[42]	RCT	“The clinical and radiographic effects of diluted silver diamine fluoride (1:10) and light cure calcium hydroxide as indirect pulp capping agents in primary molars were evaluated. At the end of 12 months, the overall clinical and radiographic success rate of indirect pulp therapy with SDF was 96% and 91.6% with light cure calcium hydroxide, respectively, although the difference was not statistically significant. In primary molars with deep carious lesions, dilute silver diamine fluoride (1:10) can be recommended as a potential indirect pulp capping agent.”

“Table: 5. Summary of SDF as Indirect pulp capping agent.”

SIDE EFFECTS OF SILVER DIAMINE FLUORIDE [9, 21]:



“Figure: 10. Flowchart representation of side effect of silver diammine fluoride”

FLOWCHART OF CLINICAL SUMMARY OF SDF:




Figure: 11. Caries Arrest




Figure: 12. Root Canal Irrigant




Figure: 13. Desensitizing Agent



Figure: 14. Indirect Pulp Agent

“Figure: 15. Flowchart of clinical summary of silver diammine fluoride”

PATIENT INFORMATION SHEET ^[43]:**Silver diammine fluoride information sheet**

Silver diammine fluoride (SDF) is an antibiotic liquid. It is used on decayed baby teeth to treat tooth sensitivity and it can also help to stop tooth decay. It is most effective when applied twice yearly. However, treatment with SDF does not remove the need for regular dental checks, fillings or crowns to repair function or aesthetics.

The procedure: 1) Drying the affected area, 2) Placing a small amount of SDF on the affected area, 3) Allowing SDF to dry for one minute, 4) Rinsing with water.

Your child's teeth should not be treated with SDF if they:


- 1) are allergic to silver
- 2) have painful gums or sores in their mouth.

Advantages of SDF

- ✓ Relieves tooth sensitivity.
- ✓ Prevents abscesses by slowing down or stopping tooth decay.
- ✓ Can buy time for children who are too young or fearful, or who have special needs, until they can manage.
- ✓ Avoids fillings or extractions by stopping decay.

Disadvantages of SDF

- * The painted area will stain black permanently. Healthy parts of teeth will not stain. However, stained tooth may be covered with a filling or a crown to make it look white again.
- * Tooth-coloured fillings may discolour if SDF is applied to them but this can usually be polished off to make them white again.
- * If accidentally applied to the skin or gum, a brown stain may appear. Although this cannot be washed off, it causes no harm and will disappear in 1-3 weeks.
- * If accidentally spilled on clothes, it can leave a stain that does not come out.
- * A metallic taste may be noticed during the application. This will go away rapidly.

Before SDF**After SDF****Alternatives to SDF to discuss with your child's dentist (not limited to the following):**

- * No treatment, which may lead to continued deterioration of tooth structures and cosmetic appearance. Symptoms may increase in severity.
- * Depending on the location, extent of the tooth decay and your child's ability to cooperate, other treatment options may include a filling, a silver crown, or an extraction.

If SDF is being used to stop tooth decay, sometimes the decay will still progress. If this happens, the tooth may require further treatment, such as reapplication of SDF, placement of a filling or a crown or extraction.

Reproduced with permission: Professor Innes, University of Dundee. Dundee Dental Hospital & School, UK

CONCLUSION:

SDF is an antibacterial agent that is simple, safe, non-invasive (painless), and cost-effective for the treatment of carious lesions across the age spectrum. It is advised or can be the material of choice for children or individuals who are unable to tolerate conventional restorative treatment modalities, those with special health care needs, and populations with limited access to a dental care. SDF is not only the best possible way to arrest and prevent dental caries in individuals at the dental clinic but also at the community level.

REFERENCE:

1. Rosenblatt A, Stamford TC, Niederman R (February 2009). "Silver diamine fluoride: a caries "silver fluoride bullet"". *Journal of Dental Research*. 88 (2): 116–25.
2. Stebbins EA. What value has argentinitras as a therapeutic agent in dentistry?. *Int Dent J*. 1891;12:661-70.
3. Howe PR. A method of sterilizing and at the same time impregnating with a metal affected dentinal tissue. *Dent Cosmos*. 1917; 59:891-904.
4. Nishino M, Yoshida S, Sobue S, Kato J, Nishida M. Effect of topically applied ammoniacal silver fluoride on dental caries in children. *The Journal of Osaka University Dental School*. 1969 Sep;9:149-55.
5. Yamaga R. Diamine silver fluoride and its clinical application. *J Osaka Univ Dent Sch*. 1972;12:1-20.
6. Crystal YO, Marghalani AA, Ureles SD, Wright JT, Sulyanto R, Divaris K, Fontana M, Graham L. Use of silver diamine fluoride for dental caries management in children and adolescents, including those with special health care needs. *Pediatric dentistry*. 2017 Sep 15;39(5):135E-45E.
7. American Dental Association. CDT 2017 Dental Procedures Codes. Chicago: American Dental Association Publishing; 2017.
8. Mei ML, Chu CH, Lo EC, Samaranyake LP. Fluoride and silver concentrations of silver diammine fluoride solutions for dental use. *International journal of paediatric dentistry*. 2013 Jul;23(4):279-85.
9. Shah S , Bhaskar V , Venkatraghavan K , Choudhary P , M.Ganesh , Trivedi K.Silver Diamine Fluoride: A Review and Current Applications. *Journal of Advanced Oral Research / Jan-Apr 2014 / Vol. 5 No.1*
10. "SILVERDIAMINEFLUORIDE (CAS No. 34445-07-3)Retrieved 14 July 2016.
11. Peng JY, Botelho MG, Matinlinna JP. Silver compounds used in dentistry for caries management: a review. *Journal of dentistry*. 2012 Jul 1;40(7):531-41.
12. Horst JA, Ellenikiotis H, Milgrom PL (January 2016). "UCSF Protocol for Caries Arrest Using Silver Diamine Fluoride: Rationale, Indications and Consent". *Journal of the California Dental Association*. **44** (1): 16–28.
13. Seifo N, Robertson M , MacLean J , Blain K , Grosse S , Milne R , Seeballuck C and Innes N_The use of silver diamine fluoride (SDF) in dental practice *BRITISH DENTAL JOURNAL | VOL 228 NO. 2 | January 24*
14. Oliveira BH, Cunha-Cruz J, Rajendra A, Niederman R (August 2018). "Controlling caries in exposed root surfaces with silver diamine fluoride: A systematic review with meta-analysis". *Journal of the American Dental Association*. **149** (8): 671–679.e1.
15. Horst JA, Ellenikiotis H, Milgrom PL. UCSF protocol for caries arrest using silver diamine fluoride: rationale, indications, and consent. *J Calif Dent Assoc* 2016;44(1):16–28.
16. "Minimally Invasive Treatment for Molar Incisor Hypomineralization". *Decisions in Dentistry*. Retrieved 2020-04-09
17. Mei ML, Lo EC, Chu CH (February 2016). "Clinical Use of Silver Diamine Fluoride in Dental Treatment". *Compendium of Continuing Education in Dentistry*. **37** (2): 93–8, quiz100. PMID 26905088.
18. Crystal YO, Marghalani AA, Ureles SD, Wright JT, Sulyanto R, Divaris K, Fontana M, Graham L (September 2017). "Use of Silver Diamine Fluoride for Dental Caries Management in Children and Adolescents, Including Those with Special Health Care Needs". *Pediatric Dentistry*. **39** (5): 135–145. PMID 29070149
19. Crystal YO, Niederman R (January 2019). "Evidence-Based Dentistry Update on Silver Diamine Fluoride". *Dental Clinics of North America*. **63** (1): 45–68.
20. Zhao IS, Gao SS, Hiraishi N, Burrow MF, Duangthip D, Mei ML, et al. (April 2018). "Mechanisms of silver diamine fluoride on arresting caries: a literature review". review. *International Dental Journal*. **68** (2): 67–76.
21. Mei ML, Ito L, Cao Y, Li QL, Chu CH, Lo EC (March 2014). "The inhibitory effects of silver diamine fluorides on cysteine cathepsins". primary. *Journal of Dentistry*. **42** (3): 329–35.
22. Suzuki T, Tsutsumi N, Sobue S, Suginaka H (1976). "Effect of diammine silver fluoride on plaque formation by *Streptococcus mutans*". primary. *Japanese Journal of Oral Biology*. **18** (3): 268–278. .
23. Yamaga R. Mechanisms of action of diammine silver fluoride and its use. *Nippon Dent. Rev* 1970 328:180-187. (In Japanese)
24. Chu CH, Lo EC, Lin HC. Effectiveness of silver diamine fluoride and sodium fluoride varnish in arresting dentin caries in Chinese preschool children. *J Dent Res* 2002;81(11):767–770
25. Llodra JC, Rodriguez A, Ferrer B, et al. Efficacy of silver diamine fluoride for caries reduction in primary teeth and first permanent molars of schoolchildren: 36-month clinical trial. *J Dent Res* 2005;84(8):721–724.
26. Yee R, Holmgren C, Mulder J, et al. Efficacy of silver diamine fluoride for arresting caries treatment. *J Dent Res* 2009;88(7):644–647.
27. Zhi QH, Lo EC, Lin HC. Randomized clinical trial on effectiveness of silver diamine fluoride and glass ionomer in arresting dentine caries in preschool children. *J Dent* 2012;40(11):962–967.

28. dos Santos Jr VE, de Vasconcelos FM, Ribeiro AG, et al. Paradigm shift in the effective treatment of caries in schoolchildren at risk. *Int Dent J* 2012;62(1):47–51.
29. Duangthip D, Chu CH, Lo ECM. A randomized clinical trial on arresting dentine caries in preschool children by topical fluorides—18 month results. *J Dent* 2016;44:57–63.
30. Duangthip D, Wong MCM, Chu CH, et al. Caries arrest by topical fluorides in preschool children: 30-month results. *J Dent* 2018;70: 74–79.
31. Fung MHT, Duangthip D, Wong MCM, et al. Randomized clinical trial of 12% and 38% silver diamine fluoride treatment. *J Dent Res* 2018;97(2):171–178.
32. Hiraishi N, Yiu C K, King N M, Tagami J, Tay F R. Antimicrobial efficacy of 3.8% silver diamine fluoride and its effect on root dentin. *J Endod* 2010; 36: 1026–1029.
33. Mathew VB, Madhusudhana K, Sivakumar N, Venugopal T, Reddy RK. Anti-microbial efficiency of silver diamine fluoride as an endodontic medicament - An ex vivo study. *Contemp Clin Dent*. 2012 Jul;3(3):262-4.
34. Ebtissam M. Al-Madi ,Manar A. Al-Jamie,Noura M. Al-Owaid,Amal A. Almohaimede, Albandary M. Al-Owid . Antibacterial efficacy of silver diamine fluoride as a root canal irrigant. Received: 9 April 2019 Revised: 22 June 2019 Accepted: 25 June 2019
35. Minavi B, Youssefi A, Quock R, et al. Evaluating the substantivity of silver diamine fluoride in a dentin model. *Clin Exp Dent Res*. 2021;7:628–633.
36. Maru V, Padawe D, Naik S, et al. Assessment of Bacterial Load Using 3.8% SDF as an Irrigant in Pulpectomized Primary Molars: A Randomized Controlled Trial. *Int J Clin Pediatr Dent* 2022;15(S-1):S47–S51.
37. Kiesow A, Menzel M , Lippert F , Tanzer J M and Milgrom P. Dentin tubule occlusion by a 38% silver diamine fluoride gel: an in vitro investigation, *BDJ Open* (2022) 8:1 ;
38. Monse B, Heinrich-Weltzien R, Mulder J, Holmgren C, van Palenstein Helderma WH. Caries preventive efficacy of silver diamine fluoride (SDF) and ART sealants in a school-based daily fluoride toothbrushing program in the Philippines. *BMC Oral Health*. 2012 Nov 21;12: 52.
39. Gupta A, Sinha N, Logani A, Shah N. An ex vivo study to evaluate the remineralizing and antimicrobial efficacy of silver diamine fluoride and glass ionomer cement type VII for their proposed use as indirect pulp capping materials - Part I. *J Conserv Dent*. 2011 Apr 14:113-116
40. Sinha N, Gupta A, Logani A, Shah N. Remineralizing efficacy of silver diamine fluoride and glass ionomer type VII for their proposed use as indirect pulp capping materials - Part II (A clinical study). *J Conserv Dent*. 2011 Jul 14 :233-236.
41. Shah A , Ganesh M, Kaur M . Evaluation of Silver Diamine Fluoride as Indirect Pulp Capping Agent in Primary Teeth: An in vivo Study, Year:2020, Volume: 10, Issue: 2, Page no. 102-107.
42. Shafi N , Kaur H, Choudhary R ,Yeluri R. Dilute Silver Diamine Fluoride (1:10) Versus Light Cure Calcium Hydroxide as Indirect Pulp Capping Agents in Primary Molars – A Randomized Clinical Trial . *J Clin Pediatr Dent* (2022) 46 (4): 273–279.
43. https://en.wikipedia.org/wiki/File:SDF_patient_info_leaflet.pdf