

A new species of *Argyreia* (Convolvulaceae) from Yunnan, China

Mao-Lin Zhang¹, Yi He¹, Quan-Ru Liu¹

¹ Key Laboratory of Biodiversity Science and Ecological Engineering, Ministry of Education, College of Life Sciences, Beijing Normal University, Beijing 100875, China

Corresponding author: Quan-Ru Liu (liuquanru@bnu.edu.cn)

Academic editor: L. Giacomini | Received 16 January 2023 | Accepted 17 April 2023 | Published 2 May 2023

Citation: Zhang M-L, He Y, Liu Q-R (2023) A new species of *Argyreia* (Convolvulaceae) from Yunnan, China. *PhytoKeys* 225: 199–209. <https://doi.org/10.3897/phytokeys.225.100646>

Abstract

Argyreia subrotunda, a new species from Yunnan Province, China, is described and illustrated. The new species resembles *A. fulvocymosa* and *A. wallichii*, but differs from these in the flowers with an entire or shallowly lobed corolla, as well as smaller elliptic bracts, lax flat-topped cymes and shorter corolla tubes. An updated key to the species of *Argyreia* from Yunnan province is also provided.

Keywords

Argyreia subrotunda, flora of Yunnan, morphology, new taxon, taxonomy

Introduction

Argyreia Lour., a genus comprising scandent shrubs or lianas, is mainly distributed throughout tropical Asia (Staples and Traiperm 2017). *Argyreia* species mainly inhabit open and sunny places such as roadsides, thickets, and edges of mingled forest (Fang and Huang 1979; Fang and Staples 1995). The number of *Argyreia* species has been increasing and is now up to 143 species following the discovery of new species (Traiperm et al. 2019; Traiperm and Suddee 2020) and the establishment of new combinations (Shalini et al. 2017; Staples and Traiperm 2017; Rattanakrajang et al. 2022). There are about 25 species in China (14 of which are endemic) and 92% of the species found in

China in Yunnan Province (23 recorded species). The province of Yunnan is, therefore, the main center of diversity in China (Fang and Huang 1979; Fang and Staples 1995; Yang et al. 2015).

Loureiro (1790) published *Argyreia* as a genus within Convolvulaceae. The genus is mainly characterized by indehiscent fleshy or mealy berries (Staples and Traiperm 2017). The various types of indumentum, inflorescence architecture, depth of corolla lobes, and number of seeds in individual berries are the main taxonomically informative characters for the delimitation of species in *Argyreia*.

Argyreia seems to be non-monophyletic in recent works, because it includes at least one of the moth-pollinated species of *Rivea* Choisy (Manos et al. 2001; Stefanovic et al. 2003). Furthermore, there is evidence that *Blinkworthia* Choisy should be subsumed under *Argyreia* (Rattanakrajang et al. 2022). We support this conclusion and although the *Argyreia* alliance clade as recovered is paraphyletic, only one *Rivea* species was used and the inclusion of other species might lead to different conclusions in the future (Rattanakrajang et al. 2022). Therefore, we think that *Rivea* and *Argyreia* are two independent genera and both supposedly monophyletic, but their limits should be revised under a phylogenetic perspective with a comprehensive sampling.

Although recent studies have shown that *Argyreia* should be treated as part of *Ipomoea* L. (Muñoz-Rodríguez et al. 2019) and *Argyreia* is merged into *Ipomoea* by Wood et al. (2020), we chose not to follow the proposed classification in the present work, as further study of Old World taxa is still required (Traiperm and Suddee 2020). As it concerns *Ipomoea*, the possibility to keep the several established smaller genera has the potential to maintain nomenclatural stability (Eserman et al. 2020). So we do not subsume *Argyreia* into *Ipomoea* at this time and accordingly maintain the well-established generic concepts (Rattanakrajang et al. 2022).

During recent field surveys in Yunnan Province, an interesting population of *Argyreia* with an entire or shallowly lobed corolla was found. After reviewing literature and comparing specimens, especially native species in Yunnan Province and adjacent countries (Vietnam, Laos, Myanmar, Cambodia and Thailand), we found that the taxon was not completely similar to any species known worldwide. Therefore, a new *Argyreia* species from China is described and illustrated here.

Material and method

Plant material was collected during field surveys in Yunnan Province from 2020 to 2021. The type specimens have been stored in the herbarium of Beijing Normal University (BNU). Morphological measurements were made from dried specimens of herbarium by Nikon digital camera, Stereoscope (ZEISS V8) and software ImageJ (Abràmoff et al. 2004). Materials for observation of pollen morphology were obtained from herbarium, picking mature pollen from the dried specimens, sticking it on the sample stages with conductive adhesive, spraying gold and photographing by Scanning Electron Microscope. The collected specimens were compared with the type specimens

of morphologically similar species at main herbariums in China (BNU, HITBC, IBSC, KUN, PE, WUK, YUKU), as well as digital images available online provided by JSTOR and herbaria abroad that are relevant for the group (A, BM, E, G, K, P). Fresh plant materials of the similar species (*A. wallichii*) were also collected for further careful comparison. All type specimens (or photos of type specimens) of accepted names and their synonyms in *Argyreia* known around the world were examined, which refer to the voucher information provided by Staples and Traiperm (2017).

Taxonomic treatment

Argyreia subrotunda Q.R.Liu & M.L.Zhang, sp. nov.

urn:lsid:ipni.org:names:77318528-1

Figs 1, 2

Type. CHINA. Yunnan Province: Malipo County, Xinzhai Village, 22°57'48.01"N, 104°46'31.11"E, along roadside, 1300 m elev., 27 Aug 2021, fl. M. L. Zhang BNU2021YN074 (holotype: BNU0053319!; isotypes: BNU!).

Diagnosis. *A. subrotunda* is unique, a small-flowered type with an entire or shallowly lobed corolla as well as exerted stamens and pistils (included in dry specimens), smaller elliptic bracts, and outer sepals ovate-circular. It is similar to *A. wallichii* in indumentum features (whitish tomentose) and fruit types (red globose berry), but differs by its smaller elliptic bracts (vs. ovate-oblong), lax flat-topped cymes (vs. compact capitate) and shorter corolla tubes (2–2.5 cm vs. 4–5 cm). Additionally, *A. subrotunda* is similar to *A. fulvocymosa* in leaf shape (broadly ovate-circular to nearly circular) and inflorescence (flat-topped cymes), but the latter is covered with densely yellowish villus and has a distinctly 5-lobed corolla, which is very easy to distinguish (Table 1).

Description. Climbing lianas; stem woody at base, herbaceous above, the former puberulent, the latter covered with whitish trichomes. Leaves simple, alternate; petiole 6–10 cm long, tomentose; leaf blades broadly ovate to rounded, 13–16 × 12–15 cm; base truncate or slightly cordate, occasionally oblique, margins entire, apex acute or obtuse, sometimes slightly emarginate; adaxially green, sparsely whitish velutinous only along leaf veins, abaxially paler, densely shining tomentose; secondary veins 13–15 on either side, curved to edge, veins slightly raised adaxially, more prominently raised abaxially. Inflorescences flat-topped cymes, axillary or terminal; peduncle 2–5 cm long, tomentose, angulate, secondary and tertiary peduncle 6–12 mm long; bracts small, elliptic, 8–10 × 4–8 mm, obtuse, hairy outside, veined; pedicels 5–7 mm long, up to 10 mm in fruit. Flowers diurnal; sepals unequal, 2 outer ovate-circular, 8–9 × 6–7 mm, 3 inner elliptic, 6–7 × 3–5 mm, apex obtuse, abaxially whitish tomentose, adaxially glabrous, veined, enlarged in fruit, rose-red, shiny. Corolla tubular-funnelform, 2–2.5 cm long, pink, densely whitish villous outside on mid-petaline bands, otherwise glabrous, limb entire or shallowly lobed. Stamens exerted; filaments filiform, 14–15 mm long, attaching to the site of ca. 5 mm from stamens base, expanded at attachment points




Figure 1. *Argyreia subrotunda* Q.R.Liu & M.L.Zhang, sp. nov. **A** stem with leaves and inflorescences **B** bract (outside) **C** bract (inside) **D** sepals from outer (left) to innermost (right) **E** opened corolla showing mid-petaline bands **F** opened corolla with stamens **G** pistil **H** fruit with persistent sepals **I** seed (adaxial surface) **J** seed (lateral surface). All drawn by Quan-Ru Liu from voucher specimens *M. L. Zhang* BNU2021YN074 (BNU!) (**A–G**), *X. B. Guo* BNU2021YN081 (BNU!) (**H–J**).




Figure 2. *Argyreia subrotunda* Q.R.Liu & M.L.Zhang, sp. nov. **A** plant habit **B** inflorescence **C** flower in frontal view **D** fruit with persistent sepals **E** seeds: adaxial surface (left); lateral surface (right). Scale bar: 5 mm. Photographs **A–C, E** by Mao-Lin Zhang, **D** by Xi-Bing Guo.

and densely whitish hairy there; anthers oblong, 3–4 mm long; pollen globose, pantoporate, with spines, 90–101 μm in diameter. Pistil exserted; disc ringlike, glabrous, ca. 1 mm high; ovary glabrous, ovoid, 2–3 mm high; style filiform, glabrous, 20–22 mm

1000 m elev., 2 Feb. 1940, fr. *C. W. Wang et al.* 86509 (PE!); Malipo County, Wentian Road beside National Highway G246, 650 m elev., 23 Nov. 2021, fr. *X. B. Guo* BNU2021YN081 (BNU!); Malipo County, Xinzhai Village, 1300 m elev., 23 Nov. 2021, fl. *X. B. Guo* BNU2021YN082 (BNU!).

Pollen morphology

The observed pollen grains of *A. subrotunda* were monad, spheroidal to subspheroidal and radially symmetrical, with polyantoporate and echinate ornamentation (Fig. 4). It was possible to divide into two types based on the pollen morphology as follows: the diameter of the pollen grain was less than 100 μm with shorter bottle-like spines (5–7 μm), such as *A. wallichii* and the new species *A. subrotunda*; the diameter of pollen grains was over 100 μm with longer cone-shaped spines ($\geq 10 \mu\text{m}$), such as *A. marlipoensis*, which is endemic to Yunnan province, and the flower of which is first seen in this study.

Discussion

Morphologically, this species is most similar to *A. wallichii* and *A. fulvocymosa*, and it can be easily distinguished by the characters summarized in Table 1. The new species was similar to *A. wallichii*, both having similar indumentum features and fruit types as well as being almost sympatric. However, based on 17 specimens from two populations of *A. subrotunda*, we found the length of corolla was a very stable feature, about 2–2.5 cm, which was significantly shorter than *A. wallichii* (4–5 cm). The latter could also be easily distinguished from the new species by its compact capitulate cymes and ovate-oblong bracts instead of flat-topped cymes and elliptic bracts. Additionally, *A. subrotunda* was similar to *A. fulvocymosa* in leaf shape and inflorescence, but the latter had a distinctly 5-lobed corolla instead of an entire or shallowly lobed corolla. Morphological comparisons of fresh plants between *A. subrotunda* and *A. wallichii* were provided in Figure 5. Furthermore, detailed comparisons of *A. fulvocymosa*, *A. subrotunda* and *A. wallichii* were provided in Table 1.




Figure 4. Comparison of pollen morphology **A** *Argyrea subrotunda* **B** *A. wallichii* **C** *A. marlipoensis*. Scale bars: 20 μm .

Table 1. Comparisons of *A. fulvocymosa*, *A. subrotunda* and *A. wallichii*.

Character		<i>A. fulvocymosa</i>	<i>A. subrotunda</i>	<i>A. wallichii</i>
Inflorescence		flat-topped cymes	flat-topped cymes	capitate cymes
Bract	Shape	unknown	elliptic	ovate-oblong
	Size	unknown	0.8–1 cm × 0.4–0.8 cm	2.5–3.5 cm × 1.5–2.5 cm
Outer sepal	Shape	broadly ovate-circular	ovate-circular	elliptic-oblong
Corolla	Length	ca. 2 cm	2–2.5 cm	4–5 cm
	Mid-petaline bands indumentum	yellowish hirsute	whitish villous	whitish villous
	Limb	distinctly 5-lobed	entire or shallowly lobed	entire or shallowly lobed
Stamen and pistil		exserted	exserted	included




Figure 5. *Argyreia subrotunda* **A** opened corolla with 5 stamens **B** bract **E** inflorescence. *A. wallichii* **C** opened corolla with 5 stamens **D** bract **F** inflorescence.

The discovery of the new species has important value for further understanding of the morphological patterns of *Argyreia* in China. The new species is endemic to southwest China and compared to other species with an entire or shallowly lobed corolla in China, it seems to have comparative smaller corollas, which reinforces its recognition as a new taxa.

Key to the species of *Argyreia* from Yunnan, China

- 1 Corolla entire or shallowly lobed.....2
- Corolla deeply 5-lobed.....17

- 2 Climbing herb, stems slender, roots thick 3
- Climbing shrubs or lianas, stem woody at base 4
- 3 Leaf blade linear; sepals ovate; corolla pale purple, 4–4.5 cm long
..... *A. lineariloba*
- Leaf blade ovate to ovate-deltate; sepals linear-lanceolate; corolla red, ca. 7 cm
long *A. baoshanensis*
- 4 Bracts soon deciduous 5
- Bracts persistent 7
- 5 Leaf blade densely yellowish sericeous-velutinous abaxially *A. velutina*
- Leaf blade sparsely strigose or hispid abaxially 6
- 6 Leaf blade lanceolate or ovate to ovate-elliptic *A. henryi*
- Leaf blade broadly ovate to nearly circular, truncate or slightly cordate
..... *A. strigillosa*
- 7 Bracts with more than 15 mm in length 8
- Bracts 2–13 mm in length 13
- 8 Inflorescence paniculate-umbelliform, lax 9
- Inflorescence capitate, condensed 10
- 9 Leaf blade sparsely hispid abaxially, somewhat reddish purple colored; bracts
narrowly lanceolate; corolla campanulate-funnelform, 5–7 cm long
..... *A. marlipoensis*
- Leaf blade densely silvery sericeous; bracts ovate-circular; corolla urceolate-
funneliform, 2.5–3.5 cm long *A. monosperma*
- 10 Indumentum brown or dull yellow, hirsute *A. capitiformis*
- Indumentum whitish or pale yellow, villous or pubescent 11
- 11 Bracts ligulate, petiolate, apex acuminate *A. roxburghii* var. *ampla*
- Bracts ovate-circular or broadly ovate, apex acute or obtuse 12
- 12 Bracts densely curly sericeous villous abaxially, ovate-circular; sepals brown,
narrow-oblong *A. eriocephala*
- Bracts densely pubescent abaxially, broadly ovate; sepals rose purplish, ovate-
oblong *A. wallichii*
- 13 Leaf blade broadly ovate-circular, base truncate or slightly cordate, densely
yellowish villous or whitish tomentose abaxially 14
- Leaf blade narrowly oblong or ovate-oblong, to elliptic, base rounded or
broadly cuneate, densely silvery sericeous-pilose abaxially 15
- 14 Leaf blade densely dull yellowish sericeous villous abaxially; corolla broadly
funneliform, ca. 4.5 cm long, purple *A. fulvovillosa*
- Leaf blade densely whitish tomentose abaxially; corolla tubular-funneliform,
2–2.5 cm long, pink *A. subrotunda*
- 15 Peduncle 10.5–13.5 cm long *A. splendens*
- Peduncle less than 10 cm long 16
- 16 Sepals subequal, ovate-oblong; corolla white *A. cheliensis*
- Sepals unequal, elliptic or oblong; corolla purple *A. monglaensis*

- 17 Inflorescence capitate; bracts persistent; peduncle short to none, 0–0.3 cm long *A. osyrensis*
- Inflorescence paniculate-umbelliform; bracts tiny or caducous; peduncle elongated, ca. 0.5 cm long **18**
- 18 Young stems, leaves abaxially, inflorescence all densely silvery sericeous-pilose; cymes axillary or terminal, 3–10-flowered..... *A. obtusifolia*
- Young stems, leaves abaxially, inflorescence all densely yellowish tomentose; cymes axillary, 9–40-flowered *A. fulvocymosa*

Acknowledgements

We would like to express gratitude to the curators of the herbaria A, BM, E, G, HITBC, IBSC, K, KUN, P, PE, WUK, YUKU for assistance and access to specimens; Lai Wei (BNU) for providing valuable comments and suggestions; Xi-Bing Guo for taking part in field collection work and providing specimens. This research was supported by the National Natural Science Foundation of China (no. 31770213).

References

- Abràmoff MD, Magalhães PJ, Ram SJ (2004) Image processing with ImageJ. *Biophotonics International* 11(7): 36–42.
- Eserman LA, Sosef MSM, Simão-Bianchini R, Utteridge TMA, Barbosa JCJ, Buril MT, Chatrou LW, Clay K, Delgado G, Desquilbet TE, Ferreira PPA, Grande AJR, Hernández AL, Huerta-Ramos G, Jarret RL, Kojima RK, Landrein S, Lourenço JAAM, Man ID, Miller RE, More S, Moreira ALC, Mwanga-Mwanga I, Nhanala S, Pastore M, Petrongari FS, Pisuttimarn P, Pornpongrungrueng P, Rifkin J, Santos FDS, Shimpale VB, Silva SS, Stinchcombe JR, Traiperm P, Vasconcelos LV, Wang ML, Villordon A, Yang J, Yenchu GC, Heider B, Simões ARG (2020) Proposal to change the conserved type of *Ipomoea*, nom. cons. (Convolvulaceae). *Taxon* 69(6): 1369–1371. <https://doi.org/10.1002/tax.12400>
- Fang RC, Huang SH (1979) *Flora Reipublicae Popularis Sinicae*. Science Press, Beijing, 64(1): 118–120. [http://www.iplant.cn/frps/pdf/64\(1\)/118y.pdf](http://www.iplant.cn/frps/pdf/64(1)/118y.pdf)
- Fang RC, Staples GW (1995) Convolvulaceae. In: Wu CY, Raven PH (Eds) *Flora of China*. Science Press, Beijing, 313–321. <http://www.iplant.cn/foc/pdf/Argyreia.pdf>
- IUCN (2019) *Guidelines for Using the IUCN Red List Categories and Criteria*. Version 14. Prepared by the Standards and Petitions Committee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [Accessed 5 January 2020]
- Loureiro J (1790) *Flora Cochinchinensis* (Vol. 1). Typis, et expensis academicis, Ulyssipone [Lisboa], 134–135. <https://doi.org/10.5962/bhl.title.560>
- Manos PS, Miller RE, Wilkin P (2001) Phylogenetic analysis of *Ipomoea*, *Argyreia*, *Stictocardia*, and *Turbina* suggests a generalized model of morphological evolution in morning glories. *Systematic Botany* 26: 585–602.

- Muñoz-Rodríguez P, Carruthers T, Wood JR, Williams BR, Weitemier K, Kronmiller B, Goodwin Z, Sumadijaya A, Anglin NL, Filer D, Harris D, Rausher MD, Kelly S, Liston A, Scotland RW (2019) A taxonomic monograph of *Ipomoea* integrated across phylogenetic scales. *Nature Plants* 5(11): 1136–1144. <https://doi.org/10.1038/s41477-019-0535-4>
- Rattanakrajang P, Sumanon P, Traiperm P, Staples G, Utteridge T (2022) Reduction of *Blinkworthia* (Convolvulaceae) based on multilocus phylogenetic reconstruction and resurrection of a species from synonymy revealed by phenetic analyses. *Kew Bulletin* 77(4): 859–883. <https://doi.org/10.1007/s12225-022-10052-1>
- Shalini S, Lakshminarasimhan P, Maity D (2017) A new combination and lectotypification in *Argyreia* (Convolvulaceae). *Nelumbo* 59(2): 145–148. <https://doi.org/10.20324/nelumbo/v59/2017/120448>
- Staples GW, Traiperm P (2017) A nomenclatural review of *Argyreia* (Convolvulaceae). *Taxon* 66(2): 445–477. <https://doi.org/10.12705/662.12>
- Stefanovic S, Austin DF, Olmstead RG (2003) Classification of Convolvulaceae: A Phylogenetic Approach. *Systematic Botany* 28(4): 791–806.
- Traiperm P, Suddee S (2020) A new species of *Argyreia* (Convolvulaceae) from Thailand. *PhytoKeys* 149: 109–115. <https://doi.org/10.3897/phytokeys.149.50379>
- Traiperm P, Fujikawa K, Chitchak N, Srisanga P, Maknoi K, Staples GW (2019) A new species of *Argyreia* (Convolvulaceae) from Myanmar. *Willdenowia* 49(1): 65–70. <https://doi.org/10.3372/wi.49.49108>
- Wood JRI, Muñoz-Rodríguez P, Williams BRMBRM, Scotland RW (2020) A foundation monograph of *Ipomoea* (Convolvulaceae) in the New World. *PhytoKeys* 143: 1–823. <https://doi.org/10.3897/phytokeys.143.32821>
- Yang SZ, Chen PH, Staples GW (2015) *Argyreia akoensis* (Convolvulaceae), a new species from southern Taiwan. *Taiwania* 60(3): 117–122. <https://doi.org/10.6165/tai.2015.60.117>