



CODEN [USA]: IAJ PBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Research Article

INCIDENCE OF MYOCARDIAL INFARCTION AMONG KNOWN PATIENTS OF ISCHEMIC STROKE

Pooja Bai¹, Manisha Lohana¹, Munesh Kumar¹, FNU Saveeta², Anand Kumar Lohana¹,
Ahmed Qudoos¹

¹Liaquat University of Medical & Health Sciences, Jamshoro., ²Peoples University of Medical & Health Sciences, for Women, Shaheed Benazirabad.

Article Received: March 2023

Accepted: April 2023

Published: May 2023

Abstract:

Background Acute myocardial infarction in individuals who have had a cerebrovascular accident or transient ischemic attack (CVA-TIA) is a medical emergency, which must be examined and treated as soon as possible. Physicians face a significant problem in managing this scenario because early treatment of one ailment would surely postpone treatment of the other. Early detection and treatment will have an impact on the patient's morbidity and mortality in the future, as well as aid in the patient's rehabilitation.

Objective: To determine the incidence of myocardial infarction among known patients of Ischaemic stroke.

Methodology: This prospective observational study was conducted on 103 patients who presented with stroke at the medicine outpatient department of Liaquat University Hospital (Hyderabad). On the basis of ECG alterations and cardiac biomarkers, Myocardial Infarction was diagnosed among patients who presented with altered sensorium. The diagnosis of stroke/TIA was based on the onset of a localized neurological loss that lasted for at least 24 hours (stroke) or recovering earlier (TIA). A CT/MRI scan of the brain was performed for a definitive diagnosis to identify ischemia/hemorrhage as the cause of ischaemic stroke.

Results: According to the age-based distribution, the highest rate of myocardial infarction (8%, 8) was observed in those aged 51-60 years. The male-to-female ratio is 1.86:1. Thirty-two patients had diabetes, among them 75% had only elevated creatine kinase MB (CKMB) with no myocardial infarction (MI), whereas 59 patients had hypertension of which 70% had only elevated CK-MB with no MI. ST-elevation myocardial infarction (STEMI) with high CKMB accounted for 14.78% (15) of the cases, but the majority (71%, 73) of the cases had elevated CKMB with no MI, and the rest presented with normal CKMB.

Conclusion: The study reported that the incidence of MI among patients of Ischaemic Stroke was found to be higher than that of the general population. This finding highlights the need for increased vigilance and monitoring for MI in this patient population. **Keywords:** Acute Myocardial Infarction, ST-Elevation Myocardial Infarction (STEMI), CK-MB, Transient Ischemic Attacks, Cerebro-Vascular Accident

Corresponding author:

Dr. Pooja Bai,

Liaquat University of Medical & Health Sciences, Jamshoro.

poojashewkani7@gmail.com

+923353560306 | +17182008744

QR code



*Please cite this article in press Pooja Bai et al, **Incidence of Myocardial Infarction among known patients of Ischemic Stroke,** Indo Am. J. P. Sci, 2023; 10 (05).*

INTRODUCTION:

Acute coronary syndrome and cerebrovascular accident (CVA) are both medical emergency that needs to be diagnosed and treated side by side. Acute ischemic stroke has been shown to increase the incidence of acute coronary syndrome and vice versa [1-3]. Patients with transient ischemic attack have a higher relative risk of myocardial infarction (MI) than the general population [4]. According to a study, 13.7% of individuals with acute ischemic stroke exhibited an increased cardiac biomarker [5]. In the absence of primary cardiac issues, myocardial damage was found following ischemic stroke, the pathogenesis of which is unknown. More than 50% of coronary stenosis was found in up to one-third of patients with ischemic stroke who had no previous history of cardiac symptoms.

Following an acute ischemic stroke, there is a substantial chance of recurrence; nevertheless, myocardial infarction is the major cause of death in these patients [4, 6, 7]. Treating such patients is extremely difficult due to the lack of a clear guideline in them. Patients presenting with acute MI with concomitant CVA-TIA (cerebrovascular accidents or transient ischemic attacks) exhibited a complicated hospital course, with a mortality rate of 41%-61% [8,9]. The goal of this study was to determine the incidence of myocardial infarction in patients who had a cerebrovascular accident.

METHODOLOGY:

This prospective observational study was conducted on 103 patients who presented with stroke at the medicine outpatient department of Liaquat University Hospital (Hyderabad). On the basis of ECG alterations and cardiac biomarkers, Myocardial Infarction was diagnosed among patients who presented with altered sensorium. The diagnosis of stroke/TIA was based on the onset of a localized neurological loss that lasted for at least 24 hours (stroke) or recovering earlier (TIA). A CT/MRI scan of the brain was performed for a definitive diagnosis to identify ischemia/hemorrhage as the cause of ischemic stroke. Myocardial infarction was defined on the basis of ECG findings: ST-segment elevation of at least 2 mm in two contiguous precordial leads and 1 mm in two adjacent limb leads or new left bundle branch block (LBBB) or development of pathological Q waves in the ECG was suggestive of ST-elevation myocardial infarction


(STEMI) whereas for unstable angina or non-ST-segment elevation MI criteria were transient ST elevation, ST depression, or new T wave inversion. Chronic kidney disease patients with baseline elevated CK-MB enzymes were excluded from the study. For all patients with a confirmed CVA, demographic data such as age and sex, as well as historical data such as diabetes mellitus, hypertension, and coronary artery disease, were obtained during the trial.

RESULTS:

The study included 103 patients who had suffered a cerebro-vascular event. According to their CK-MB values, patients were divided into three subgroups: normal (15), elevated with no MI (73) and ST-elevation myocardial infarction (STEMI) with elevated CK-MB (15).

The age of the patients ranged from 11 to 90 years, with a mean of 59.84 ± 14.65 years. There were 67 men and 36 women among the 103 patients. Gender (male/female) distribution among subgroups was as follows: STEMI with high CK-MB was 11/4, high CK-MB without myocardial infarction was 45/28, and normal CK-MB was 11/4. However, the association was not found to be statistically significant ($p > 0.05$).

FIGURE NO. 1: INCIDENCE OF MYOCARDIAL INFARCTION AMONG DIFFERENT SEX



Seventy percent (71) patients had only elevated CK-MB without myocardial infarction, while 16% (17) had STEMI with increased CK-MB, and 14% (15) had normal CK-MB ($p > 0.05$). Only 22 of the patients had a history of coronary artery disease. Fifteen of them had only elevated CKMB at the time of presentation and four had normal CK-MB and STEMI each.

TABLE NO. 1: INCIDENCE OF MYOCARDIAL INFARCTION AMONG DIABETES MELLITUS & HYPERTENSIVE PATIENTS

	NORMAL CKMB	ELEVATED CKMB WITHOUT MI	STEMI WITH INCREASED CKMB	TOTAL	P-VALUE
DIABETES MELLITUS					
No	12(80)	49(67.12)	10(66.67)	71(68.93)	
Yes	3(20)	24(32.88)	5(33.33)	32(31.07)	0.64
	15(100)	73(100)	15(100)	103(100)	
HYPERTENSION					
No	7(46.67)	32(43.84)	5(33.33)	44(42.72)	
Yes	8(53.33)	41(56.16)	10(66.67)	59(57.28)	0.714
	15(100)	73(100)	15(100)	103(100)	

A total of 32 patients had diabetes, among them 75% (24) had only elevated CKMB with no MI, 16% (5) had STEMI, and 9% (3) had normal CK-MB. A total of 59 patients had hypertension, among them 70% (41) had only elevated CK-MB with no MI, 17% (10) had STEMI, and 13% (8) had normal CK-MB ($p > 0.05$). At the end of the study, total mortality was 18% (18): 16% (16) in the group with raised CKMB without MI and 2% (2) in patients with STEMI ($p > 0.05$).

DISCUSSION:

The purpose of this study was to determine the rate of acute myocardial infarction in patients who presented with altered sensorium due to cerebrovascular accidents in a tertiary care hospital in Pakistan. Several studies have found an association between myocardial infarction and CVA. As both are emergency medical situations, they must be handled together and without delay in view of increased morbidity and mortality. Atherosclerosis is the core pathophysiology in stroke, coronary artery disease, and myocardial infarction. Any atherosclerotic event, whether in the same or a different arterial bed, increases the chance of another [10].

According to this study, the total incidence of myocardial infarction is 15%, with a maximum incidence of eight cases in the 51-60 year age group, which is consistent with the study, which found 12.7% incidence of acute myocardial infarction in cerebrovascular accident patients [11]. Gunnoo et al. found that more than half of individuals with CVA exhibit signs of asymptomatic coronary plaque and one-third have severe stenosis, defined as $>50\%$ blockage. As a result, many people without a history

of ischemic heart disease (IHD) may be at risk for myocardial infarction. According to the MASS (Multiple Atherosclerosis Site in Stroke) study, 51% of coronary plaque was demonstrated in patients of stroke with no atherosclerotic plaque in cerebral arteries [2,12].

The male-to-female ratio is 1.86:1, which is similar to other studies [13, 14]. Hormonal effects (estrogen) on cerebral circulation, optimal blood pressure, and genetic factors contribute to a lower incidence of acute cerebrovascular accidents among the female population [13, 14].

In our study, only ischemic stroke patients were studied. The majority has increased CK-MB without any heart ailments. In patients without MI, CK-MB was found to be elevated in 24% of people in 51-60 years followed by 17% of people in 61-70 years who had CVA. Isolated elevations of CKMB were found more in CVA patients with both diabetes and hypertension. These findings were similar to a study by Mochmann et al., who reported that the majority of acute ischemic stroke patients with elevated cardiac biomarkers do not have concomitant acute coronary syndrome, and that nearly half of the patients had no

signs of obstructive CAD. Also while there was angiographic evidence of a coronary culprit lesion and high troponin in around one-fourth of all patients with acute ischemic stroke, there was no underlying culprit lesion in the majority of acute ischemic stroke patients [15].

The increased levels of cardiac biomarkers in patients with acute stroke could be explained by cardiac myocyte rupture caused by sympathetic nervous system activation and injury to the insular cortex, which causes a rise in catecholamine such as adrenaline and noradrenaline. Due to disruption in capillary perfusion induced by increased platelet aggregation, high catecholamine levels in the myocardium promote calcium influx in cardiac myocytes, which compromises cardiac contractility and function. The higher level of CK-MB in brain injury is also explained by the B subunit of CK-BB. Increased CPK-MB without changes in troponin I could be the result of neurogenic effects [16-18].

There was 18% mortality in cases presenting with high CK-MB (>35 ng/ml) although the association was not found to be significant. James et al. have also reported 17% mortality in individuals with acute ischemic stroke and elevated cardiac biomarker. The increased mortality could be attributed to severe stroke in these patients. Raised Troponin T was found to have a non-significant trend with the severity of ischemic stroke. Diabetes, peripheral vascular disease, and cardiac arrhythmia are all risk factors for early MI following a stroke [19].

Routine cardiac biomarker measurement in patients with CVA is not routinely done. In this study, increased cardiac biomarkers had a non-significant trend with a poor prognosis for short-term survival and hence can be used as a prognosticating factor. As per a study done by Etgen et al., the American Stroke Association, the Scottish Intercollegiate Guidelines Network, and the National Institute of Clinical Excellence-based UK acute stroke guidelines do not recommend routine measurement of biomarkers, but it may be helpful for patients who require early intervention for secondary prevention of coronary artery disease [20-23].

The limitation of the study is the small sample size, so a larger cohort is needed to ascertain the association of mortality and cardiac biomarkers in patients with CVA.

CONCLUSION:

The study reported that the incidence of MI among patients of Ischaemic Stroke was found to be higher than that of the general population. This finding highlights the need for increased vigilance and monitoring for MI in this patient population.

REFERENCES:

1. Hyperacute simultaneous cardiocerebral infarction: rescuing the brain or the heart first? Kijpaisalratana N, Chutinet A, Suwanwela NC. *Front Neurol*. 2017;8:664
2. Quantifying the risk of heart disease following acute ischaemic stroke: a meta-analysis of over 50,000 participants. Gunnoo T, Hasan N, Khan MS, Slark J, Bentley P, Sharma P. *BMJ Open*. 2016;6:0.
3. The incidence of stroke after myocardial infarction: a meta-analysis. Witt BJ, Ballman KV, Brown RD Jr, Meverden RA, Jacobsen SJ, Roger VL. *Am J Med*. 2006;119:354.
4. Incidence and predictors of myocardial infarction after transient ischemic attack: a population-based study. Burns JD, Rabinstein AA, Roger VL, Stead LG, Christianson TJ, Killian JM, Brown RD Jr. *Stroke*. 2011;42:935–940.
5. Myocardial infarction as a complication in acute stroke: results from the Austrian stroke unit registry. Gatteringer T, Niederkorn K, Seyfang L, et al. *Cerebrovasc Dis*. 2014;37:147–152.
6. Coronary risk evaluation in patients with transient ischemic attack and ischemic stroke: a scientific statement for healthcare professionals from the Stroke Council and the Council on Clinical Cardiology of the American Heart Association/American Stroke Association. Adams RJ, Chimowitz MI, Alpert JS, Awad IA, Cerqueria MD, Fayad P, Taubert KA. *Circulation*. 2003;108:1278–1290.
7. Inclusion of stroke as an outcome and risk equivalent in risk scores for primary and secondary prevention of vascular disease. Dhamoon MS, Elkind MS. *Circulation*. 2010;121:2071–2078.
8. Myocardial infarction and stroke. Komrad MS, Coffey CE, Coffey KS, McKinnis R, Massey EW, Califf RM. *Neurology*. 1984;34:1403–1409.
9. Cerebrovascular accident complicating acute myocardial infarction: incidence, clinical significance, and short-long-term mortality rates. Behar S, Tanne D, Abinader E, et al. *Am J Med*. 1991;91:45–50.
10. Prevalence of asymptomatic coronary artery disease in ischemic stroke patients: the

- PRECORIS study. Calvet D, Touzé E, Varenne O, Sablayrolles JL, Weber S, Mas JL. *Circulation*. 2010;121:1623–1629.
11. Myocardial infarction coincident with cerebrovascular accidents in the elderly. Chin PL, Kaminski J, Rout M. *Age Ageing*. 1977;6:29–37.
 12. Autopsy prevalence of coronary atherosclerosis in patients with fatal stroke. Gongora-Rivera F, Labreuche J, Jaramillo A, Steg PG, Hauw JJ, Amarenco P. *Stroke*. 2007;38:1203–1210.
 13. Feasibility study of stroke surveillance: data from Bangalore, India. Nagaraja D, Gururaj G, Girish N, Panda S, Roy AK, Sarma GRK, Srinivasa R. <https://pubmed.ncbi.nlm.nih.gov/19942742/> *Indian J Med Res*. 2009;130:396–403.
 14. Study of ECG changes and its relation to mortality in cases of cerebrovascular accidents. Purushothaman S, Salmani D, Prarthana KG, Bandelkar SM, Varghese S. *J Nat Sci Biol Med*. 2014;5:434–436.
 15. Coronary angiographic findings in acute ischemic stroke patients with elevated cardiac troponin: the troponin elevation in acute ischemic stroke (TRELAS) study. Mochmann HC, Scheitz JF, Petzold GC, et al. *Circulation*. 2016;133:1264–1271.
 16. The insular cortex and cardiovascular system: a new insight into the brain-heart axis. Nagai M, Hoshida S, Kario K. *J Am Soc Hypertens*. 2010;4:174–182.
 17. Neuroanatomic correlates of stroke-related myocardial injury. Ay H, Koroshetz WJ, Benner T, et al. *Neurology*. 2006;66:1325–1329.
 18. The insular cortex and cardiac response to stroke. Cheshire WP Jr, Saper CB. *Neurology*. 2006;66:1296–1297.
 19. Relation between troponin T concentration and mortality in patients presenting with an acute stroke: observational study. James P, Ellis CJ, Whitlock RM, McNeil AR, Henley J, Anderson NE. *BMJ*. 2000;320:1502–1504.
 20. Cardiac troponins and N-terminal pro-brain natriuretic peptide in acute ischemic stroke do not relate to clinical prognosis. Etgen T, Baum H, Sander K, Sander D. <https://pubmed.ncbi.nlm.nih.gov/15604421/> *Stroke*. 2005;36:270–275.
 21. Guidelines for the early management of adults with ischemic stroke: a guideline from the American Heart Association/American Stroke Association Stroke Council, Clinical Cardiology Council, Cardiovascular Radiology and Intervention Council, and the Atherosclerotic Peripheral Vascular Disease and Quality of Care Outcomes in Research Interdisciplinary Working Groups: the American Academy of Neurology affirms the value of this guideline as an educational tool for neurologists. Adams HP Jr, del Zoppo G, Alberts MJ, et al. *Stroke*. 2007;38:1655–1711.
 22. Elevated troponin after stroke: a systematic review. Kerr G, Ray G, Wu O, Stott DJ, Langhorne P. *Cerebrovasc Dis*. 2009;28:220–226.
 23. National Collaborating Centre for Chronic Conditions. London: Royal College of Physicians; 2008. *Stroke: National Clinical Guideline for Diagnosis and Initial Management of Acute Stroke and Transient Ischaemic Attack (TIA)*.