TWO SMARANDACHE SERIES

Maohua Le

Abstract. In this paper we consider the convergence for two Smarandache series.

Key words. Smarandache reciprocal series, convergence.

Let $A = \{a(n)\}_{n=1}^{\infty}$ and $B = \{b(n)\}_{n=1}^{\infty}$ be two Smarandache sequences. Then the series

$$S(A,B) = \sum_{n=1}^{\infty} \frac{a(n)}{b(n)}$$

is called the Smarandache series of A and B. Recently, Castillo [1] proposed the following two open problems.

Problem 1. Is the series

(1)
$$S_1 = \frac{1}{1} + \frac{1}{12} + \frac{1}{123} + \frac{1}{1234} + \cdots$$

convergent?

Problem 2. Is the series

(2)
$$S_2 = \frac{1}{1} + \frac{12}{21} + \frac{123}{321} + \frac{1234}{4321} + \cdots$$

Convergent?

In this paper we completely solve the mentioned problems as follows.

Theorem. The series S_1 is convergent and the series S_2 is divergent.

Proof. Let $r(n)=1/12\cdots n$ for any positive integer n. Since

(3)
$$\lim_{n\to\infty} \frac{r(n+1)}{r(n)} = \frac{12\cdots n}{12\cdots n(n+1)} < 1,$$

by D'Alembert's criterion, we see from (3) that S_1 is convergent.

Let $s(n)=12\cdots(n-1)n$ / $n(n-1)\cdots 21$ for any positive integer n. If $n=10^t+1$, where t is a positive integer, then we hare

(4)
$$S(n) = \frac{12 \cdots (10 \cdots 01)}{(10 \cdots 01) \cdots 21} > 1.$$

Therefore, by (4), we get from (2) that
$$\sum_{\infty}^{\infty} S(n) > \sum_{t=1}^{\infty} s(10^{t}+1) > \sum_{t=1}^{\infty} 1 = \infty$$
.

Thus, the series S_2 is divergent. The theorem is proved.

Reference

J. Castillo, Smarandache series, Smarandache Notions J., http://www.gallup.unm.edu/~smarandache/SERIES.TXT.

Department of Mathematics Zhanjiang Normal College Zhanjiang, Guangdong P.R. CHINA