

Two new species of *Brachytrycherus* Arrow, 1920 from China (Coleoptera, Endomychidae)

Ling-Xiao Chang¹, Wen-Xuan Bi², Guo-Dong Ren¹

1 College of Life Sciences, Hebei University, Baodin 071002, China **2** Room 401, No. 2, Lane 155, Lianhua South Road, Shanghai, 201100 China

Corresponding author: Guo-Dong Ren (gdren@hbu.edu.cn)

Academic editor: M. Thomas | Received 21 December 2015 | Accepted 30 May 2016 | Published 2 June 2016

<http://zoobank.org/B8182BB2-4A5D-49F9-BDB0-FEC5791F136F>

Citation: Chang L-X, Bi W-X, Ren G-D (2016) Two new species of *Brachytrycherus* Arrow, 1920 from China (Coleoptera, Endomychidae). ZooKeys 595: 137–146. doi: 10.3897/zookeys.595.7569

Abstract

Two new species of *Brachytrycherus* from China, *B. conaensis* **sp. n.** and *B. curviantennae* **sp. n.** are described and illustrated. *Brachytrycherus conaensis* **sp. n.** is the first species of the Handsome Fungus Beetles recorded feeding on Ascomycetes. A key to the species of *Brachytrycherus* known in China is provided.

Keywords

Coleoptera, Endomychidae, new species, taxonomy, China

Introduction

The genus *Brachytrycherus* was established by Arrow (1920) with *B. perotteti* as the type species. It is a member of the largest endomychid subfamily Lycoperdininae, the monophyly of which was confirmed by the phylogenetic studies of Tomaszewska (2000, 2005).

In 2015, a large-scale phylogenetic study was presented for Cucujoidea by Robertson et al., using molecular evidence to rebuild the relationship tree of this superfamily, and established one new superfamily, Coccinelloidea Robertson et al., 2015. The Endomychidae was included in it. Through this significant study, the monophyly of subfamily Lycoperdininae is more clear, and with subfamily Epipocinae forms the sister group to Endomychinae+Stenotarsinae (Robertson et al. 2015).

Tomaszewska (2005) placed *Brachytrycherus* with another seven genera in the Amphisternus-group: *Amphisternus* Germar, 1843, *Amphistethus* Strohecker, 1964, *Cacodaemon* Thomson, 1857, *Gerstaeckerus* Tomaszewska, 2005, *Ohtaius* Chùjò, 1938, *Spathomeles* Gerstaecker, 1857 and *Stictomela* Gorham, 1886. The monophyly of this group is well supported based on the following synapomorphies: mesoventrite with intercoxal process widened laterally towards apex, overlapping parts of coxae; elytra with basal margins thickened and raised, mandible with apical tooth widely chisel-shaped, male genital segment with additional internal V- or U-shaped sclerite (Tomaszewska 2005). Since then, two new genera of the Amphisternus-group were described, *Stroheckeria* Tomaszewska, 2006 from Vietnam, and *Humerus* Chang & Ren, 2013a from China.

Strohecker (1964) in his synopsis of the tribe Amphisternini (=Amphisternus-group of Tomaszewska (2005)) listed four species of *Brachytrycherus*, (*B. convexus* Strohecker, 1964, *B. gemmatus* (Arrow, 1928), *B. madurensis* Arrow, 1920 and *B. perotetti* Arrow, 1920) of which *B. convexus* as a new species and *B. gemmatus* as a new combination moved from *Engonius*. He also provided a key to the species of *Brachytrycherus* known at that time. Prior to the present study, *Brachytrycherus* included six species (Shockley et al. 2009a): *B. concolor* Arrow, 1937 (Borneo), *B. convexus* Strohecker, 1964 (India), *B. femoralis* (Arrow), 1928 (Laos, Vietnam), *B. gemmatus* (Arrow), 1928 (Laos, Myanmar and Thailand), *B. madurensis* Arrow, 1920 (India, Taiwan) and *B. perotteti* Arrow, 1920 (India). Only one of which was previously known from China: *B. madurensis*.

During the examination of the Endomychidae collected in China, two new species were recognized and are described here.

Material and methods

Type specimens of the new species described here are deposited in the following institutions or private collections:

- MHBU** Museum of Heibei University, Baoding, China;
CBWX Collection of Wenxuan Bi, Shanghai, China;
SHEM Shanghai Entomology Museum, Chinese Academy of Sciences, Shanghai, China.
MIZ Museum and Institute of Zoology, Polish Academy of Sciences, Warszawa, Poland.

The specimens were examined and described using a Nikon® SMZ800 dissecting microscope. The following measurements were made using a Leica® M205 A dissecting microscope: body length from apical margin of clypeus to apex of elytra; width across both elytra (at widest part); elytral length along suture, including scutellum. The aedeagus was boiled in 10% NaOH solution, cleaned, and finally dissected in distilled water. Habitus photos were taken using a Canon® Eos 5D III SLR camera and Canon® MP-E 65mm macro lens. All photographs were modified in Adobe Photoshop® CC 2015.

Taxonomy

Brachytrycherus Arrow, 1920

Brachytrycherus Arrow, 1920: 12.

Type species. *Brachytrycherus perotteti* Arrow, 1920.

Diagnosis. The species of *Brachytrycherus* resemble those of *Ohtaius* and *Gerstaeckerus* in having the body black or blackish-brown, elytral maculae transverse, most often orange or yellow. These genera share the feature of having the mandibles chisel-shaped apically. However, *Brachytrycherus* can be distinguished from these other genera by the following combination of characters: 1) body less elongate; 2) head with well-developed gular sutures; 3) mesoventral process with sides parallel; 4) maxillary laciniae with tuft of S-like setae apically (Tomaszewska 2005).

Brachytrycherus conaensis sp. n.

<http://zoobank.org/278F3113-648B-4DFF-BBFB-D2AB2B47177A>

Figs 1–2, 5

Type material. Holotype, male, Xizang, Cona, Lexiang, 2500–2600 m, 20–30.VI.2013, Wen-Xuan Bi leg. (MHBU); Paratypes, 1 female, same data as holotype. 2 females, Xizang, Medog, Beibeng, Gelincun, 1700 m, 3.VIII.2014, Wen-Xuan Bi leg. (CBWX); 3 males, 7 females, Xizang, Cuona, Lexiang, 2500 m, 6.VIII.2010, Wen-Xuan Bi leg. (CBWX); 5 males, 6 females, ditto except 15.VII.2011 (CBWX); 26 males, 11 females, ditto except 29–30.VI.2013 (CBWX); 1 male, 1 female, ditto except (MZPW); 18 males, 1 female, ditto except 2500–2600 m, 20–30.VI.2013 (CBWX); 1 female, ditto except 2700 m, 18.VI.2013 (CBWX).


Etymology. The specific name is derived from the type locality.

Diagnosis. *Brachytrycherus conaensis* is similar to *B. madurensis* in appearance, but can be differentiated by each elytron with three maculae, anterior two maculae nearly rhomboid in shape, sometimes connected to each other, and the anterior and posterior elytral maculae without dentition.

Description. Length 8.2–8.3 mm. Body oval, about 1.8–1.9 times as long as wide; rather convex; shiny. Colour black with three red maculae on elytra.

Head. Antenna 11-segmented, long and rather slender, nearly 1/2 body length, with antennomeres 1–8 distinctly longer than wide; scape approximately 4.5 times as long as pedicel; antennomere 3 slightly shorter than 4 and 5 combined; antennomeres 4 nearly as long as 5, antennomeres 5–8 gradually shorter; club composed of 3 antennomeres, moderately broad, flat, loose. Maxilla with terminal palpomere elongated, almost 2.0 times as long as palpomere 3, tapering anteriorly, truncate apically.

Thorax. Pronotum 2.0–2.3 mm long, 3.2–3.3 mm wide; widest near 1/2 of pronotal length; coarsely and densely punctate; lateral margins rather narrowly bordered,



Figures 1–2. Dorsal and ventral habitus of *B. conaensis* sp. n. **1** male **2** female. a = dorsal view, b = ventral view. Scale bar 1 mm.

sides nearly parallel; front angles produced anteriorly, rather acute; disc weakly convex, two small round raised area laterally; transverse wrinkle laterally; median furrow absent; lateral sulci linear, deep, extending to basal 1/3 length of pronotum; basal sulcus nearly straight, deep. Prosternal process rather narrowly separates procoxae; not extending beyond coxae; sides in male weakly curved outwardly, rounded apically; in female sides nearly straight, weakly truncate apically. Mesoventral process transverse, lateral margins widening apically and overlapping part of mesocoxae; posterior margin nearly straight. Elytra 5.9–6.1 mm long, 4.5–4.7 mm wide; 2.7–3.0 times as long as pronotum and 1.4 times as wide as pronotum, sides curved, widest near 1/2 length of elytron; densely and coarsely punctate; humeri rather prominent. Each elytron with three irregular red maculae. Anterior 2 elytral maculae nearly rhombus, located near apical 1/4, medial macula larger than lateral one, sometimes narrowly connected. Posterior macula transverse, anterior margin shallowly emarginate or nearly straight, posterior margin U-shaped, widely emarginate.. All tibiae with sexual characters; protibiae in male with concavity on inner edge of apical 1/4, in female without concavity; mesotibiae abruptly curved from near apical 1/3 to apex, in female gently curved; metatibiae in male abruptly widened from near 1/3 length to apex, in female gently widened.

Abdomen with five ventrites in both sexes. Ventrite 5 with lateral margins gently converging posteriorly, three pairs of longitudinal short wrinkles laterally; posterior margin weakly curved medially in male; in female ventrite 5 lateral margins abruptly converging posteriorly, without longitudinal wrinkles; posterior margin truncate, nearly straight medially. Aedeagus (Fig. 5) rather long, heavily sclerotized, straight. Median lobe branched apically; branch long and rather straight, abruptly raised near basal 1/3 length, gently converging apically, flat, acute and weakly reflexed apically. Tegmen basal, comparatively large, ring-shaped.

Biology and ecology. Almost all individuals were found active on fence, woodpile or timber piles within the village and its surrounding area at night (Figs 9–11). Some larvae and adults were found (sometimes at the same time) feeding on the surface of the perithecia or spores of *Daldinia concentrica* (Xylariaceae) (Fig. 10), seeming to prefer the asexual phase; however, individuals were also found on mature ascocarps.


Based on the study of the natural history of the handsome fungus beetles (Shockley et al. 2009b), this report is possibly the first record of the handsome fungus beetles feeding on ascomycetes. In addition, some individuals were found active on the wood without fungus, and may be feeding on lichen growing on the wood (Fig. 11). This species association may not be host-specific.

***Brachytrycherus curviantennae* sp. n.**

<http://zoobank.org/CA118D1D-C4CD-4DB6-BEFB-22D840A76F1A>

Figs 3–4, 6

Type material. Holotype, male, Xizang, Medog, 1500 m, 20.VIII.2013, Wen-Xuan Bi leg. (SHEM); Paratypes, 1 female, Xizang, Medgo, Beibeng, Gelincun, 3.VIII.2014,



Figures 3–4. Dorsal and ventral habitus of *B. curviantennae* sp. n. **3** male **4** female. **a** = dorsal view, **b** = ventral view. Scale bar 1 mm.



Figures 5–6. Aedeagus **5** *B. conaensis* sp. n.; **6** *B. curviantennae* sp. n. **a** = lateral view, **b** = apical view. Scale bars 1 mm.

Wen-Xuan Bi leg. (MHBV); 1 female, Xizang, Medgo, Beibeng, Gelincun, 3.VIII.2014, Wen-Xuan Bi leg. (CBWX).

Etymology. The name refers to the antennomere 3 distinctly outwardly curved.

Diagnosis. *Brachytricherus curviantennae* is similar to *B. madurensis* in appearance, but can be differentiated by antennomere 3 distinctly curved outwards, pronotum sides strongly curved, elytral maculae with front and hind margins emarginate.

Description. Length 8.5–9.4 mm. Body broadly oval, approximately 1.6–1.8 times as long as wide; convex; shiny. Colour black with four red maculae on elytra.

Head. Antenna 11-segmented, long and rather slender, nearly 1/2 body length, with antennomeres 1–8 distinctly longer than wide; scape approximately 4.5 times as long as pedicel; antennomere 3 distinctly curved, and nearly as long as 4 and 5 combined; antennomere 4 as long as 5, antennomeres 5–8 gradually shorter; club composed of 3 antennomeres, broad and flat. Maxilla with terminal palpomere elongate, almost 2.0 times as long as palpomere 3, tapering anteriorly, truncate apically.

Thorax. Pronotum 2.0–2.4 mm long, 4.1–4.2 mm wide; widest near 1/2 of pronotal length; coarsely and densely punctate; lateral margins narrowly bordered, sides in male wavy and strongly curved; in female sides smooth and strongly curved; front angles produced anteriorly, blunt; disc weakly convex, with two large round raised areas laterally; transverse wrinkle laterally; median furrow absent; lateral sulci linear, deep, extending to basal 1/3 of pronotal length; basal sulcus nearly straight, deep. Prosternal process moderately separates the procoxae; sides weakly curved outwardly, weakly truncate apically. Mesoventral process transverse, lateral margins widening apically and overlapping part of mesocoxae; posterior margin nearly straight. Elytra 6.0–7.1 mm long, 5.1–5.2 mm wide; 3.0 times as long as



Figures 7–11. Habitats of *Brachytrycherus* species. **7** large clump of Fagaceae plants of collecting site in Xizang, China **8** male of *B. curviantennae* sp. n. (arranged) **9** village of collecting site in Xizang, China **10** male of *B. conaensis* sp. n. and larva on the wood pile **11** female of *B. conaensis* sp. n. feeding on the lichen growing on wood.

pronotum and 1.2–1.3 times as wide as pronotum, sides curved, widest near 1/2 length of elytron; densely and coarsely punctate; humeri rather prominent. Each elytron with two transverse, irregular in shape red maculae. Anterior elytral macula

nearly cymbiform, anterior margin widely U-shaped and deeply emarginate, posterior margin weakly wavy. Posterior macula transverse, inversely cymbiform, anterior margin weakly wavy, posterior margin widely U-shaped and deeply emarginate. Protibiae in male with concavity on inner edge of apical 1/3, in female without concavity; mesotibiae abruptly curved from near apical 1/3 to apex, in female gently curved; metatibiae in male abruptly widened from near 1/3 length to apical 1/4, in female gently widened.

Abdomen with 5 ventrites in both sexes. Ventrite 5 with lateral margins gently converging posteriorly, posterior margin widely rounded medially in male; in female ventrite 5 lateral margins abruptly converging posteriorly, posterior margin truncate, nearly straight medially. Aedeagus (Figs 6) rather long, heavily sclerotized, straight. Median lobe branched apically; branch long and rather straight, gently rising from about basal 1/3 to apical 1/3, flat, acute and weakly reflexed apically. Tegmen placed basally, comparatively large, ring-shaped.

Biology and ecology. The male was hand collected by simple searching, as it is active on branches at night (Fig. 7). Two females were collected by shaking the tree from a large clump of dead wood of Fagaceae plants (Fig. 8).

Key to the species of *Brachytrycherus* known in China

- 1 Antennomere 3 distinctly outwardly curved; pronotum sides strongly curved..... *B. curviantennae* sp. n.
- Antennomere 3 straight; sides of pronotum nearly parallel..... **2**
- 2 Each elytron bearing 2 transverse, strongly dentate maculae.... *B. madurensis*
- Each elytron bearing 3 maculae, anterior 2 maculae nearly rhomboid, sometimes connected to each other; elytral maculae not dentate *B. conaensis* sp. n.

Conclusions

Prior to this study, only *B. madurensis* was recorded from China (Taiwan) (Shockley et al. 2009a). Two new species comprise the first record of *Brachytrycherus* from mainland China.

Acknowledgements

We are indebted to Dr. W. Tomaszewska (Museum and Institute of Zoology, Warsaw, Poland) and Dr. F.W. Shockley (University of Georgia, Athens) for providing useful comments to improve this manuscript. We thank very much Mr. Michael Thomas (Subject editor of ZooKeys) for made some corrections about the English usage of this manuscript.

References

- Arrow GJ (1920) A contribution to the classification of the Coleopterous family Endomychidae. Transactions of the Entomological Society of London, 1–83. doi: 10.1111/j.1365-2311.1920.tb00205.x
- Chang L-X, Ren G-D (2013a) *Humerus medogicus*, new genus and new species from Xizang, China (Coleoptera: Endomychidae). Annales Zoologici 63(1): 95–100. doi: 10.3161/000345413X666138
- Gorham HS (1886) On new Genera and Species of Endomychidae. Proceedings of the Zoological Society, 154–163.
- Shockley FW, Tomaszewska KW, McHugh JV (2009a) An annotated checklist of the handsome fungus beetles of the world (Coleoptera: Cucujoidea: Endomychidae). Zootaxa 1999: 1–113.
- Shockley FW, Tomaszewska KW, Mchugh JV (2009b) Review of the Natural History of the Handsome Fungus Beetles (Coleoptera: Cucujoidea: Endomychidae). Insecta Mundi 0072: 1–24.
- Robertson et al. (2015) Phylogeny and classification of Cucujoidea and the recognition of a new superfamily Coccinelloidea (Coleoptera: Cucujiformia). Systematic Entomology 40: 745–778. doi: 10.1111/syen.12138
- Strohecker HF (1964) A synopsis of the Amphisternini (Coleoptera: Endomychidae). Pacific Insects 6(2): 319–357.
- Tomaszewska KW (2000) Morphology, phylogeny and classification of adult Endomychidae (Coleoptera: Cucujoidea). Annales Zoologici 50(4): 449–558.
- Tomaszewska KW (2005) Phylogeny and generic classification of the subfamily Lycoperdinae with a re-analysis of the family Endomychidae (Coleoptera: Cucujoidea). Annales Zoologici 55 (suppl. 1): 1–172.
- Tomaszewska KW (2006) *Stroheckeria quadrimaculata*, new genus and new species of Lycoperdinae from Vietnam (Coleoptera: Endomychidae). Annales Zoologici 56: 465–470.