

A checklist of the mammals (Mammalia) from Durango, western Mexico

Diego F. García-Mendoza* and Celia López-González

CIIDIR Unidad Durango, Instituto Politécnico Nacional, Calle Sigma 119, Fraccionamiento 20 de Noviembre II, Durango, Durango, México 34220.
* Corresponding author. E-mail: diegofgarciam@gmail.com

ABSTRACT: An updated list of the mammals of Durango state, Mexico was built from literature records and Museum specimens. A total of 139 species have been recorded, representing 28.3 % of the Mexican terrestrial mammals, and 25.1 % species more compared to the previous account. Two species have been extirpated from the state, 23 are endemic to Mexico. Four major ecoregions have been previously defined for the state, Arid, Valleys, Sierra, and Quebradas. Species richness is highest at the Quebradas, a tropical ecoregion, whereas the aridlands are the least species-rich. The Sierra has the highest number of endemic species (11) followed by Quebradas (7), Valleys and Arid (3). Despite the fact that Durango harbors one of the largest diversities of mammals of the country, conservation efforts are minimal, and the current protected areas do not include the most species-rich regions. The current rate of anthropogenic modification in the state makes urgent to put in practice already existing plans to protect Durango's unique biodiversity.

INTRODUCTION

Mexico harbors one of the most diverse mammalian fauna (527 species, 488 terrestrial), surpassed only by Brazil and Indonesia. It also has the second largest number of endemic species worldwide (Bradley *et al.* 2004; Carraway 2007; Ceballos *et al.* 2005, Romo-Vázquez 2005; Tejedor 2006). As with the rest of the world, in Mexico Mammalia is one of the best known zoological groups. Few new species have been described from the country recently, and those new taxa are mostly cryptic species discovered through analysis of DNA sequences. Nonetheless, some areas are better known than others: most mammalian research and inventories have been historically focused on the central and southeastern portions of the country, while less attention has been paid to the northern areas. Durango is the fourth largest state in Mexico (123,314 km² INEGI 2005), a landlocked area located in the north central portion of the country (Figure 1). Durango is traversed by the tropic of Cancer from east to west, and by the Sierra Madre Occidental from Northwest to Southeast, both factors resulting in a complex landscape. Moreover, in the Sierra Madre of Durango, as in other parts of Mexico, the Nearctic and Neotropical biogeographic regions meet, thus potentially a larger number of species than expected otherwise may occur within its borders (López-González and García-Mendoza 2007).

Because historically most of the current territory of Durango was not explored by naturalists, prior to the 1880s very few records of its mammalian fauna are available, and usually they are part of historical writings, rather than biological accounts (e.g. de Morfi 1777-1778). The oldest account of the local mammalian fauna we could find is by O. Thomas, who reported a small list of mammals collected in southwestern Durango (Thomas 1882). The mammal list for the state increased slowly in the following century with works by Allen (1903; 1904), Goldman (1951) and Baker (1960). The first state-wide account was published in 1962 (Baker and Greer 1962), and remains to date

the only treatise on the mammalian fauna of the state. Following this work, a number of papers on the taxonomy and distribution of Durango mammals have been published (Webb and Baker 1962; Jones 1964; Gardner 1965; Crossin *et al.* 1973; Baker and Baker 1975; Petersen 1976; 1979; Álvarez and Polaco 1984; Houseal *et al.* 1987; Servín *et al.* 1995; Hernández Huerta *in litt*; Muñiz-Martínez and Arroyo-Cabral 1996; Muñiz-Martínez and Polaco 1996; Servín *et al.* 1996; Servín 1997; Aragón-Piña and Garza-Herrera 1999; Bell *et al.* 2001; López-González 2003; Muñiz-Martínez *et al.* 2003; Servín *et al.* 2003a; Bradley *et al.* 2004; Gómez-Ruiz *et al.* 2006; Carraway 2007; Torres-Morales *et al.* 2010; Charre-Medellín *et al.* 2011). During the last quarter of the 20th century, a group of researchers associated to the only protected areas of the state at that time (La Michilía and Mapimí Biosphere Reserves)




FIGURE 1. Geogaphic location of Durango State, Mexico, and main ecological regions (after González Elizondo *et al.* 2007). See text for detailed description of regions.

published the results of numerous research projects, mostly focused on the ecology of the species (Gallina *et al.* 1978; Gallina 1981; Grenot 1981; Gallina y Folliott 1983; Servín 1991; Gallina and Excurra 1992; Servín and Huxley 1991a; 1991b; Servín 1997; Hernández *et al.* 1987; Alvarez *et al.* 1988; Alvarez and Arroyo-Cabral 1990; Rogovin *et al.* 1991; Servín and Huxley 1993; Hernández and Delibes 1994; López Vidal and Alvarez 1997; Servín, 2000; Servín *et al.* 2003b; González Romero *et al.* 2005; Hernández *et al.* 2005; Servín 2005). Further research has been conducted in other areas of the state, on ecology (Drake 1958; Petersen 1973; 1975; 1978; Petersen y Petersen 1979; Petersen 1980a; 1980b; 1985; Servín 1991; Servín and Huxley 1993a; 1993b; Petersen 1993; Muñiz-Martínez 1997; Servín 1997; Servín *et al.* 2003b; López-González and Torres-Morales 2004; Portales *et al.* 2004; Lopez-González 2005; López-González *et al.* 2010), ectoparasites (Montiel-Parra *et al.* 2001; Villegas-Guzmán *et al.* 2005), and more recently ethnozoology (Jacobo-Salcedo *et al.* 2011). These papers have added a significant number of species to Baker and Greer's (1962) inventory, and have increased our knowledge on the mammalian fauna of Durango. Nonetheless, no updated checklist on the mammals for the state has been published since that account; thus the objective of this work is to present an updated checklist of the mammals of Durango state, with comments on their general distribution.

MATERIALS AND METHODS

The updated list was built up from Baker and Greer (1962) and Ramírez Pulido *et al.* (2000). We conducted a search in the major internet journal databases for all literature published up to November 2012, using "mammals" and "Durango" as keywords. Additional, previously unpublished records were scored from specimens deposited at the Mammal Collection, CIIDIR-IPN Unidad Durango (CRD); Colección Nacional de Mamíferos, Universidad Nacional Autónoma de México (CNMA); Colección de Mamíferos, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, México (ENCB); the Museum of Michigan State University mammal (MSU); Natural Science Research Laboratory, The Museum, Texas Tech University (TTU); and the Mammal Collection, Department of Ecology and Evolutionary Biology, University of Kansas (KU). Thus, all records included in this account are based either on specimens examined by us or by published records based on museum specimens. Nomenclature was standardized following Ceballos *et al.* (2005), except for the order Soricomorpha, for which we follow Carraway (2007), the genera *Natalus*, for which we follow Tejedor (2005, 2006) and *Spermophilus*, for which we follow Helgen *et al.* (2009). Zoogeographic affinities follow Álvarez and Lachica (1974). Protection status and endemisms are indicated when appropriate following Ceballos *et al.* (2005), current Mexican regulations (SEMARNAT 2010) and IUCN red list (IUCN 2012). We did not include in the checklist introduced or domestic species.

To give general information on the habitat of species, we assigned each species to one or more ecological regions as defined by González Elizondo *et al.* (2007). These authors divided the state into four ecological zones that

encompass the major biomes of the state (Figure 1), from east to west they were:

1) Arid and semiarid region - Lowland plains and mountain ranges from the Bolson de Mapimí in the north to the border with Zacatecas in the south. This area is part of the Chihuahuan desert and constitutes the western portion of the Mexican Plateau. Elevations range from 1070 to 2200 m above sea level. Climates vary from dry, to very dry and warm (BWh, BSh, BWh' – Köppen modified by García 1988). Vegetation includes a variety of thorn scrubs, as well as halophytic and gypsophile associations in a few isolated areas.

2) Valleys region - A wide strip of land extending from northwest to southeast between the Chihuahuan desert and the Sierra Madre Occidental Mountains. It includes mostly grasslands, as well as mesas and a few small mountain ranges. Elevation ranges from 1600 to 2400 m, descending northeastwards to the Mapimi desert. Climate is warm and dry, with extreme temperatures throughout the day (BS₁K - González-Elizondo *et al.* 2007) in most of the area. Although originally it was covered with grasslands and mesquite grassland, currently most of this area has been modified for agriculture and cattle ranching.

3) Sierra Region - It includes the Sierra Madre Occidental mountains, from 2400 m above sea level on the eastern slope, and 2000 m on the western slope, to 3340 m on the summits. Climates are temperate, dry on the eastern slope (C(w)), and more mesic and cooler (C(E)(w)) on the higher portions, González Elizondo *et al.* 2007). Vegetation includes pine, pine and oak forests, open oak forests, as well as secondary chaparral produced by deforestation. Small patches of cloud forest also occur on the highlands of the western slope.

4) Quebradas Region - Lowlands of the western slope of the Sierra Madre Occidental. Elevation ranges from 2000 to 130 m above sea level. The region is rugged, with steep slopes and deep canyons. The climate is warm, with a long dry season and high precipitation (700-1500 mm) in the rain season (AC(w), González-Elizondo *et al.* 2007). Accordingly, the local vegetation consists of tropical associations, such as dry, deciduous, or semideciduous tropical forests, mixed with secondary vegetation resulting from agricultural and cattle ranching activities.

We constructed a checklist including all species and ecorregions for which records were available. Regional faunas were compared via Cluster analysis (UPGMA) of a resemblance matrix based on presence-absence data. Analysis was performed using the SAHN module of NTSYSpc Ver. 2.02 (Applied Biostatistics Inc. 1998). The Jaccard index (Brower and Zar 1977) was used as a distance measure.

RESULTS AND DISCUSSION

Species richness

Eight orders, 23 families 77 genera, and 139 species of mammals were recorded from Durango (Table 1), representing 28.30% of the terrestrial mammalian fauna of Mexico. Our list increases the number of species listed by Baker and Greer (1962) in 35 (25.1%).

Species richness in Durango, by order, is what would be expected proportionally (i. e., 28.3% of the Mammals of Mexico) for Didelphimorphia, Cingulata,

and Lagomorpha (Figure 3). Artiodactyla, Rodentia and Soricomorpha are proportionally underepresented; whereas Chiroptera and Carnivora are overrepresented. Neither Primates or Perissodactyla occur in the state, as they are restricted to more humid tropical areas of Mexico. Seventy-four percent of the species have Nearctic affinities (Álvarez and Lachica, 1974 and Figure 2). Artiodactyla and Carnivora are orders of holarctic affinities, which explains their abundance in a mostly subtropical area. Bats, on the other hand, are a mixture of nearctic elements from the highlands and Mexican plateau and tropical species from the Pacific coastal plain that penetrate into the Sierra Madre Occidental through the canyons on its western slope. Rodentia might be underepresented due to the absence of tropical species that do not penetrate into the Sierra Madre Canyons, but it is also possible that the area is still not fully explored and species remain undescribed.




FIGURE 2. Percentage of neotropical and nearctic species, by order, for the mammals of Durango, Mexico. Affinities after Alvarez and Lachica, (1974).




FIGURE 3. Mammal species richness by order for Mexico (red bars) and Durango (blue bars). Data for Mexico from Bradley *et al.* 2004; Carraway 2007; Ceballos *et al.* 2005, Romo-Vázquez 2005 and Tejedor 2006.

Conservation status

Of the 139 species recorded, 23 are endemic of Mexico; most endemics are rodents (16 species), followed by bats (5, Table 1). Twenty are species of concern according to Mexican laws (SEMARNAT 2010): 7 are of special concern for being endemic or migratory, 8 are considered threatened, and 3 are endangered. Under this regulations, the Mexican wolf (*Canis lupus baileyi* Nelson and Goldman, 1929) is considered extinct, but Ceballos *et al.* (2005) considered wolves as extirpated rather than extinct because of successful reintroduction efforts in Arizona, USA (Hedrick and Fredrickson 2008). For Durango, Allen (1942) and Leopold (1959) reported *Canis lupus*, but they

did not provide specific localities or specimens to back up their claims. Baker and Greer (1962) on the other hand, collected and examined specimens from Rancho Las Margaritas, near what is today la Michilí Reserve. The last published record we could find for wolves in Durango is by McBride (1980). The grizzly bear (*Ursus arctos horribilis* Ord, 1815) is considered extinct for Mexico by Mexican law (SEMARNAT 2010). As with the Mexican wolf, because wild populations of *U. arctos* still occur in the United States (Wilson and Ruff 1999), it is more appropriate to consider grizzly bears as extirpated from Mexico rather than extinct (Ceballos and Navarro 1991). According to Allen (1942), grizzly bears (under the name *Ursus nelsoni*) were distributed across the Sierra Madre Occidental from the northwestern state of Chihuahua and northeastern Sonora to southern Durango. Subsequently, Hall (1981, 1984) and Moctezuma Orozco (2005) proposed the same distribution in their accounts on grizzly bears. Nonetheless, as far as we could investigate, there are no documented records for this species of bear in Durango, and none of these authors give evidence of their presence or refer to data gathered by other author. The southernmost record for *U. arctos* in México appears to be from Sierra del Nido, in north-central Chihuahua (Leopold 1959). Although it is possible that grizzly bears occurred in Durango, there is no conclusive evidence in historical times that document their presence.

The black bear (*Ursus americanus* Pallas, 1780) did occur in the highlands of the Sierra Madre Occidental in Durango, the last documented record we could find is a specimen killed in 1956 at Rancho Las Margaritas (Baker and Greer 1962). Delfín-Alonso *et al.* (2011) add four more records, the last one as recent as 2005. Unfortunately, they are personal communications with no evidence available to back them up. It is possible that the species still occurs in Durango, but a systematic effort would be necessary to verify its presence.

The pronghorn *Antilocapra americana* (Ord, 1815) was reported for the last time in the wild by Nelson (1925), from an area between Mapimí and Tlahualilo, in the northeastern portion of the state. Subsequent authors did not include it in the fauna of Durango (Ceballos and Navarro 1991), and we could not find recent records of pronghorns within the state; thus, we consider them as extirpated from Durango. The American bison *Bison bison* (Linnaeus, 1758) was mentioned by Allen (1876), who asserted that it was distributed in Durango in the XVI century. Nonetheless Pacheco (2005) considered that its historical distribution did not include Durango. We could find no evidence for the presence of *Bison bison* in Durango in historical times. Ceballos *et al.* (2005) reported american bisons as “critically endangered” in Mexico, because in the last 30 years the only confirmed record came from a small population observed near Janos, Chihuahua.

Additional Records

Baker and Greer (1962) suggested that the bighorn sheep (*Ovis canadensis* Shaw, 1804) occurred at the Bolsón de Mapimí, northeastern Durango, based on a record by Baker (1956) from Sierra del Diablo, 20 miles (32 km) north of the border between the states of Chihuahua and Durango. Notwithstanding, we found no record of occurrence for that area or any other in Durango. Aragón

Piña and Garza Herrera (1999) reported the presence of *Myotis occultus* Hollister, 1909 (as *Myotis lucifugus* (Le Conte, 1831)) and *Erethizon dorsatum* (Linnaeus, 1758) at Mapimí Biosphere Reserve, but they did not provide specific localities. Because the reserve lays across three different states (Durango, Chihuahua, and Coahuila), it is not possible to determine with certainty the occurrence of these species in Durango. Aragón *et al.* (2009) reported *Sciurus niger* Linnaeus, 1758 and *Neotoma lepida* Thomas, 1893 for the municipality of Canatlán, central Durango. These reports need to be reevaluated, given that *N. lepida* is a woodrat known only from Baja California and northern Sonora, and *S. niger* is a species of squirrel that in Mexico is distributed only in northern Coahuila (over 500 km northeast from Canatlán). Furthermore, externally *S. niger* resembles *S. nayaritensis* J.A. Allen, 1890, a common squirrel in their study area (Hall 1981; Ceballos and Oliva 2005), and with which it could have been confused. Muñiz-Martínez *et al.* (2003) reported *Nyctinomops laticaudatus* (E. Geoffroy, 1805) for Durango, but a recent review of the specimens on which the report is based (CRD2497-2500), revealed that they are actually *N. femorosaccus* (Merriam, 1889). Finally, *Panthera onca* (Linnaeus, 1758) (Navarro-Serment *et al.* 2005), *Sigmodon mascotensis* J.A. Allen, 1897 (CRD774-776), and *Myotis fortidens* Miller and G.M. Allen, 1928 (Jones *et al.* 1972) have been recorded in localities less than 15 km from the border with Durango, and therefore are likely to occur within the state.

Distribution by ecological regions

Species richness (136, not including the 2 extirpated species and the black bear) was higher at the Quebradas (65), followed by the Valleys (62), the Sierra (57) and the Arid region (48). The cluster analysis (Figure 4) grouped Valleys and Sierra, which share almost half the species (43.37%), most of them of Nearctic affinities (mostly Vespertilionidae bats and Muridae rodents). This group is joined to the Arid region, with which it shares 30.41% of the species, mostly vespertilionids and some carnivores (Figure 4, Table 1). The Quebradas region is the most distinct, sharing only 17.87% of the species with the rest. This region includes mostly tropical elements that are lacking in the rest of the state, but are widely distributed in the Pacific coastal plain (Figure 4). From the 65 species occurring in this area, 7 are endemic to México (*Tlacuatzin canescens* (J.A. Allen, 1893), *Artibeus hirsutus* Andersen, 1906, *Myotis carteri* La Val, 1973, *Rhogeessa parvula* H. Allen, 1866, *Cynomops mexicanus* (Jones and Genoways, 1967), *Sciurus colliae* Richardson, 1839, *Chaetodipus artus* (Osgood, 1900). The Quebradas and Arid regions share 16 species, even though they are separated by a 250 km-wide mountain range. Eight are widely distributed generalist species (e.g. *Eptesicus fuscus* (Beauvois, 1796), *Lasiurus cinereus* (Beauvois, 1796)), the rest are species that are distributed in arid and dry tropical environments (e.g. *Mormoops megalophylla* (Peters, 1864), *Choreonycteris mexicana* Tschudi, 1844).

The Sierra and Quebradas are contiguous areas that share mostly generalist species (13). *Dermanura azteca* (Andersen, 1906) and *Anoura geoffroyi* Gray, 1838, tropical species, have been captured in the highlands as well (Hernández Huerta 1996; CRD). They likely

represent vagrants from the lowlands, but also it is possible that elevational, seasonal migrations may occur between the tropical and temperate areas of the Sierra Madre Occidental. This is a subject that remains to be investigated. We recorded 15 species only at the highlands of the Sierra Madre Occidental, 5 are endemic to the Sierra (*Callospermophilus madrensis* (Merriam, 1901), *Tamias bulleri* J.A. Allen, 1889, *T. durangae* (J.A. Allen, 1903), *Peromyscus schmidlyi* Bradley *et al.*, 2004 and *Nelsonia neotomodon* Merriam, 1897) and four are endemic to México (*Corynorhinus mexicanus* G.M. Allen 1916, *Peromyscus difficilis* J.A. Allen, 1871, *Reithrodontomys zacatecae* Merriam, 1901 and *Sigmodon leucotis* Bailey, 1902).

The Arid zone (Chihuahuan Desert) is the least diverse, only 10 species occurred exclusively there, two of them endemic to Mexico (*Dipodomys nelsoni* Merriam, 1907 and *Neotoma goldmani* Merriam, 1903). This region has a long history of agricultural and livestock activity, which has resulted in considerable fragmentation and habitat loss. The effects of these changes in the mammalian fauna have not been quantified.

The Valleys region includes mostly species of Nearctic affinities. Because of its being rather a transitional area between the aridlands to the east and the Sierra to the west, only one exclusive species occurs (*Dipodomys phillipsii* Gray, 1841). It presents, instead, a mixture of elements from the desert and the Sierra. Three species occurring in this region are endemic to Mexico (*Chaetodipus nelsoni* (Merriam, 1894), *Dipodomys phillipsii*, and *Peromyscus melanophrys* (Coues, 1874)). As with the aridlands, the valleys have been strongly modified by agricultural and cattle ranching practices. Most of the original grasslands were replaced by pastures, and the area devoted to livestock has tripled in the last 30 years. In general, the landscape of the valleys is considered to have suffered from medium to high degree of modification (Márquez-Linares 2007). In the same way as the aridlands, the effect of these transformations in the mammalian fauna is unknown.

Final Remarks

Durango state is home to almost 30 % of the terrestrial mammal species known for Mexico. It is likely that the proportion will be similar for other zoological groups, and we know it is so at least for vascular plants (González-Elizondo *et al.* 1991). Durango is one of the most species rich states in Mexico, and includes both Nearctic and Neotropical species (López-González and García-Mendoza




FIGURE 4. Cluster analysis (UPGMA) of mammalian species assemblages for four ecological regions in Durango, Mexico. The Jaccard index was used as a measure of similarity. Numbers on nodes indicate percentage of species shared by branches.

2007). Nonetheless, our knowledge on their local distribution, ecology, evolution, and response to anthropic modification is scant or nonexistent. This situation is worrying, especially because even though Durango is one of the largest states in Mexico, the proportion of its territory that is protected is minimal (2.04 %, Márquez-Linares 2007), and it does not include all the major ecoregions of the state. No protected area exists for the Quebradas region, although it is the most species-rich (Table 1) and the Mexican National Comission for

Biodiversity (CONABIO) has designated as priorities for conservation five areas that include, at least partially, the Quebradas region (Arriaga *et al.* 2000). This is also one of the least studied areas, mainly due to its inaccesibility. A formal proposal for protecting these and other areas in the state already exists (Márquez-Linares 2007), and suggests a protected area that will cover up to 10.8 % of the state territory. At the current rate of modification, it is urgent to take this proposal to reality in order to ensure the persistence of Durango's unique diversity.

TABLE 1. Mammals recorded for the state of Durango, Mexico to 2012. Ar = Arid, Va = Valleys, Si = Sierra, Qu = Quebradas. * Species extirpated from the state. Conservation status as to NOM_ECOL 059-2001 (SEMARNAT 2001)/IUCN: A= threatened, Pr= Special protection, E= Extinct, P= Endangered, VU= Vulnerable, EN= Endangered. Endemic species according to Ceballos and Oliva (2005). Regions as defined by González Elizondo *et al.* (2007); see text for details.

TAXON	REGION				MEXICO ENDEMIC	STATUS	SOURCE OF RECORDS			
	Ar	Va	Si	Qu						
Order Didelphimorphia										
Family Didelphidae										
Subfamily Didelphinae										
<i>Tlacuatzin canescens</i> (J.A. Allen, 1893)			X		X		Baker and Greer (1962), CRD			
<i>Didelphis virginiana</i> Kerr, 1792		X	X	X			Baker and Greer (1962), MSU, CRD, CNMA			
Order Cingulata										
Family Dasypodidae										
Subfamily Dasypodinae										
<i>Dasypus novemcinctus</i> Linnaeus, 1758				X			Baker and Greer (1962), MSU			
Order Soricomorpha										
Family Soricidae										
Subfamily Soricinae										
<i>Sorex emarginatus</i> Jackson, 1825			X		X		Baker and Greer (1962), KU, MSU, CRD, ENCB			
<i>Sorex monticolus</i> Merriam, 1890			X				Baker and Greer, (1962), MSU			
<i>Sorex veraecrucis</i> Carraway, 2007				X			Carraway (2007)			
<i>Notiosorex crawfordi</i> (Coues, 1877)	X	X			A		Baker (1965), MSU, CRD			
Order Chiroptera										
Family Emballonuridae										
Subfamily Emballonurinae										
<i>Balantiopteryx plicata</i> Peters, 1867				X			Torres-Morales <i>et al.</i> (2010), CRD			
Family Mormoopidae										
<i>Mormoops megalophylla</i> (Peters, 1864)	X	X		X			Baker and Greer, (1962), MSU, CRD			
<i>Pteronotus parnellii</i> (Gray, 1843)				X			Baker and Greer (1962), KU, CRD			
Family Phyllostomidae										
Subfamily Macrotinae										
<i>Macrotus californicus</i> Baird, 1858				X			Baker and Greer (1962), KU, CRD			
Subfamily Micronycterinae										
<i>Glyonycteris syvestris</i> Thomas, 1896 ^a				X			CRD			
Subfamily Desmodontinae										
<i>Desmodus rotundus</i> (É.Geooffroy St.-Hilaire, 1810)				X			Gardner (1963), KU, CRD, TTU			
Subfamily Glossophaginae										
<i>Anoura geoffroyi</i> Gray, 1838			X	X			Baker and Greer (1962), MSU, CRD, CNMA			
<i>Choeronycteris mexicana</i> Tschudi, 1844	X	X		X	A		Baker and Greer (1962), MSU, CRD, CNMA			
<i>Glossophaga commissarisi</i> Gardner, 1962				X			Baker and Greer (1962), MSU, CRD			
<i>Glossophaga leachii</i> Gray, 1844 ^a				X			MSU, CRD			
<i>Glossophaga soricina</i> (Pallas, 1766)				X			Baker and Greer (1962), KU, MSU, CRD			
<i>Leptonycteris yerbabuenae</i> Martinez and Villa-Ramírez, 1940	X			X	A/VU		Crossin <i>et al.</i> (1973), MSU, KU, CRD			
<i>Leptonycteris nivalis</i> (de Saussure, 1860)	X			X	A/EN		Arita and Humprey (1988), MSU, CRD, CNMA			
Subfamily Stenodermatinae										
<i>Artibeus hirsutus</i> Andersen, 1906			X		X		Torres-Morales <i>et al.</i> (2010), CRD			
<i>Artibeus jamaicensis</i> Leach, 1821			X				Baker and Greer (1962), MSU, CRD, TTU			
<i>Artibeus intermedius</i> J.A. Allen, 1897			X				Torres-Morales <i>et al.</i> (2010), MSU, KU, CRD, CNMA TTU			
<i>Centurio senex</i> Gray, 1842			X				Crossin <i>et al.</i> (1973), MSU			
<i>Chiroderma salvini</i> Dobson, 1878			X				Crossin <i>et al.</i> (1973), MSU, CRD			
<i>Dermanura azteca</i> (Andersen, 1906)			X	X			Baker and Greer (1962), MSU, CRD, CNMA			

TABLE 1. CONTINUED.

TAXON	REGION				MEXICO ENDEMIC	STATUS	SOURCE OF RECORDS
	Ar	Va	Si	Qu			
<i>Dermanura tolteca</i> (de Saussure, 1860)			X				Baker and Greer (1962), MSU, CRD, CNMA
<i>Dermanura phaeotis</i> (Miller, 1902)			X				CNMA
<i>Sturnira lilium</i> (É. Geoffroy St.-Hilaire, 1810)			X				Baker and Greer (1962), MSU, CRD, TTU
<i>Sturnira ludovici</i> Anthony, 1824			X				Baker and Greer (1962), MSU
Family Natalidae							
<i>Natalus lanatus</i> Tejedor, 2005			X				Tejedor (2005), CRD
<i>Natalus mexicanus</i> Miller, 1902			X				Baker and Greer (1962)
Family Vespertilionidae							
Subfamily Vespertilioninae							
<i>Corynorhinus mexicanus</i> G. M. Allen, 1916			X		X		Muñiz-Martínez y Polaco (1996), CRD, CNMA
<i>Corynorhinus townsendii</i> (Cooper, 1837)	X	X	X				Baker and Greer (1962), MSU, CRD, CNMA
<i>Eptesicus fuscus</i> (Palisot de Beauvois, 1796)	X	X	X	X			Baker and Greer (1962), MSU, KU, CRD, CNMA, ENCB, TTU
<i>Euderma maculatum</i> (J.A. Allen, 1891)			X	X		Pr	Gardner (1965), ENCB, CNMA
<i>Idionycteris phyllotis</i> (G.M. Allen, 1916)			X	X			Gardner (1965), CRD, ENCB, CNMA
<i>Lasiurus blossevillii</i> (Lesson and Garnot, 1826)			X	X	X		Urbano-Vidales <i>et al.</i> (1987), CRD, CNMA
<i>Lasiurus cinereus</i> (Palisot de Beauvois, 1796)	X	X	X	X			Gardner (1965), MSU, CRD, CNMA, ENCB, TTU
<i>Lasiurus xanthinus</i> (Thomas, 1897)	X	X	X				Baker and Greer (1962), MSU, CRD, CNMA
<i>Myotis auriculus</i> Baker and Stains, 1955				X			Gardner (1965), CRD, ENCB, CNMA
<i>Myotis californicus</i> (Audubon and Bachman, 1842)	X	X	X	X			Gardner (1965), CRD, ENCB, CNMA
<i>Myotis carteri</i> LaVal, 1973				X	X	Pr	Torres-Morales <i>et al.</i> (2010), CRD
<i>Myotis melanorhinus</i> (Merriam, 1886)			X	X			Baker and Greer (1962), KU, CRD, CNMA, TTU
<i>Myotis thysanodes</i> Miller, 1897			X	X			Baker and Greer (1962), MSU, CRD, ENCB, CNMA
<i>Myotis velifer</i> (J.A. Allen, 1890)	X	X	X	X			Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA
<i>Myotis volans</i> (H.A. Allen, 1866)			X	X			Gardner (1965), MSU, CRD, ENCB, CNMA, TTU
<i>Myotis yumanensis</i> (H.A. Allen, 1864)			X	X	X		Baker and Greer (1962), MSU, CRD, ENCB, CNMA, TTU
<i>Parastrellus hesperus</i> (H.A. Allen, 1864)	X	X		X			Baker and Greer (1962), MSU, CRD, CNMA
<i>Rhogeessa parvula</i> H. Allen, 1866			X		X		Gardner (1963), KU, CRD, CNMA
Subfamily Antrozoinae							
<i>Antrozous pallidus</i> (Le Conte, 1856)	X	X	X				Baker and Greer (1962), MSU, KU, CRD, CNMA
Family Molossidae							
Subfamily Molossinae							
<i>Cynomops mexicanus</i> (Jones and Genoways, 1967)			X		X	Pr	Muñiz-Martínez <i>et al.</i> (2003), CRD
<i>Eumops perotis</i> (Schinz, 1821)	X			X			Aragón-Piña and Garza Herrera (1999), CRD, CNMA
<i>Molossus molossus</i> (Pallas, 1766)				X			Muñiz-Martínez <i>et al.</i> (2003), CRD
<i>Molossus rufus</i> É. Geoffroy St.-Hilaire, 1805				X			Gardner (1963), KU, CRD
<i>Nyctinomops femorosaccus</i> (Merriam, 1889)				X			Gardner (1963), KU, CRD
<i>Nyctinomops macrotis</i> (Gray, 1839)				X			Gardner (1965), CRD
Subfamily Tadarinae							
<i>Tadarida brasiliensis</i> (I. Geoffroy Saint-Hilaire, 1824)	X	X	X	X			Baker and Greer (1962), MSU, KU, CRD, CNMA, TTU
Order Carnivora							
Family Canidae							
Subfamily Caninae							
<i>Canis latrans</i> Say, 1823	X	X	X				Baker and Greer, (1962), MSU, KU, CRD, ENCB, CNMA
<i>Canis lupus</i> Linnaeus, 1758*				+		E	Baker and Greer, (1962), MSU
<i>Urocyon cinereoargenteus</i> (Schreber, 1775)	X	X	X	X			Baker and Greer, (1962), MSU, CNMA
<i>Vulpes macrotis</i> Merriam, 1888	X					A	Baker and Greer (1962), MSU, KU, ENCB, CNMA
Family Felidae							
Subfamily Felinae							
<i>Leopardus pardalis</i> (Linnaeus, 1758)				X		P	Servín, <i>et al.</i> (2003a)
<i>Leopardus wiedii</i> (Schinz, 1821)				X		p	Servín, <i>et al.</i> (2003a), MSU, CNMA
<i>Lynx rufus</i> (Schreber, 1777)	X	X	X				Baker and Greer (1962), MSU, CRD, ENCB, CNMA
<i>Puma concolor</i> (Linnaeus, 1771)	X	X	X				Baker and Greer (1962), MSU, CNMA
<i>Puma yagouaroundi</i> (Lacépède, 1809)				X		A	Servín <i>et al.</i> (2003a), CNMA
Family Mustelidae							
Subfamily Lutriniae							
<i>Lontra longicaudis</i> (Olfers, 1818)	X	X	X			A	Baker and Greer (1962), CNMA
Subfamily Mustelinae							
<i>Mustela frenata</i> Lichtenstein, 1831			X				Baker and Greer (1962), MSU, CRD

TABLE 1. CONTINUED.

TAXON	REGION				MEXICO ENDEMIC	STATUS	SOURCE OF RECORDS
	Ar	Va	Si	Qu			
Subfamily Taxidiinae							
<i>Taxidea taxus</i> (Schreber, 1778)	X	X			A	Baker and Greer (1962), MSU, CRD, CNMA	
Family Mephitidae							
<i>Conepatus leuconotus</i> (Lichtenstein, 1832)	X				Baker and Greer (1962), MSU, CRD		
<i>Mephitis macroura</i> Lichtenstein, 1832	X	X			Baker and Greer (1962), MSU, CRD, ENCB, CNMA		
<i>Mephitis mephitis</i> (Schreber, 1776)	X	X			Baker and Greer (1962), CRD		
<i>Spilogale gracilis</i> Merriam, 1890		X	X		Baker and Baker (1975), MSU, CRD, CNMA		
Family Procyonidae							
Subfamily Procyoninae							
<i>Bassaris astutus</i> (Lichtenstein, 1830)	X	X		X	Baker and Greer (1962), MSU, CRD, CNMA		
<i>Nasua narica</i> (Linnaeus, 1766)				X	MSU, CRD, CNMA		
<i>Procyon lotor</i> (Linnaeus, 1758)	X	X	X		Baker and Greer (1962), MSU, CRD, ENCB, CNMA		
Family Ursidae							
Subfamily Ursinae							
<i>Ursus americanus</i> Pallas, 1780			?		P	Baker and Greer (1962), MSU	
Order Artiodactyla							
Family Antilocapridae							
Subfamily Antilocaprinae							
<i>Antilocapra americana</i> (Ord, 1815)*	+	+			P	Nelson (1925)	
Family Cervidae							
Subfamily Odocoileinae							
<i>Odocoileus hemionus</i> (Rafinesque, 1817)	X				Baker and Greer (1962), MSU, ENCB		
<i>Odocoileus virginianus</i> (Zimmermann, 1780)	X	X	X		Baker and Greer (1962), MSU, CRD, CNMA		
Family Tayassuidae							
Subfamily Tayassuinae							
<i>Pecari tajacu</i> (Linnaeus, 1758)	X	X	X		Baker and Greer (1962), MSU, CRD, CNMA		
Order Rodentia							
Family Sciuridae							
Subfamily Sciurinae							
<i>Ammospermophilus interpres</i> (Merriam, 1890)	X				Baker and Greer (1962), MSU, KU		
<i>Callospermophilus madrensis</i> (Merriam, 1901)		X		X	Pr	Servín <i>et al.</i> (1996), CRD, CNMA	
<i>Ictyomys mexicanus</i> (Erxleben, 1777)	X					Aragón <i>et al.</i> (1993), CNMA	
<i>Otospermophilus variegatus</i> (Erxleben, 1777)	X	X	X	X		Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA	
<i>Sciurus aberti</i> Woodhouse, 1853			X		Pr	Baker and Greer (1962), MSU, KU, CRD, CNMA	
<i>Sciurus colliae</i> Richardson, 1839			X	X		Baker and Greer (1962), KU, CNMA	
<i>Sciurus nayaritensis</i> J.A. Allen, 1890		X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA	
<i>Tamias bulleri</i> J.A. Allen, 1889 ^b		X		X	VU	Baker and Greer (1962), MSU, CRD, ENCB, CNMA	
<i>Tamias dorsalis</i> (Baird, 1855) ^b		X				Baker and Greer (1962), MSU, CRD, ENCB, CNMA	
<i>Tamias durangae</i> (J. A. Allen, 1903) ^b		X		X		Baker and Greer (1962), MSU, KU, CRD, CNMA, TTU	
<i>Xerospermophilus spilosoma</i> Bennett, 1833	X	X				Baker and Greer (1962), MSU, KU, CRD, CNMA	
Family Geomyidae							
Subfamily Geomyinae							
<i>Cratogeomys castanops</i> (Baird, 1852)	X				Baker and Greer (1962), CNMA		
<i>Thomomys umbrinus</i> (Richardson, 1829)	X	X	X	X		Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU	
Family Heteromyidae							
Subfamily Heteromyinae							
<i>Liomys irroratus</i> (Gray, 1868)	X	X				Baker and Greer (1962), MSU, KU, CRD, CNMA, TTU	
<i>Liomys pictus</i> (Thomas, 1893)			X			Baker and Greer (1962), MSU, CRD, CNMA, TTU	
Subfamily Perognathinae							
<i>Chaetodipus artus</i> (Osgood, 1900)			X	X		Baker and Greer (1962), KU	
<i>Chaetodipus eremicus</i> (Mearns, 1898)	X	X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA	
<i>Chaetodipus hispidus</i> (Baird, 1858)	X					Baker (1965), MSU, CRD, CNMA	
<i>Chaetodipus nelsoni</i> (Merriam, 1894)	X	X			X	Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU	
<i>Chaetodipus pernix</i> (Woodhouse, 1852)			X			KU, CRD	
<i>Perognathus flavus</i> Baird, 1855	X	X				Baker and Greer (1962), MSU, KU, CRD, TTU	
Subfamily Dipodomysinae							
<i>Dipodomys merriami</i> Mearns, 1890	X					Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA	
<i>Dipodomys nelsoni</i> Merriam, 1907	X			X		Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA	

TABLE 1. CONTINUED.

TAXON	REGION				MEXICO ENDEMIC	STATUS	SOURCE OF RECORDS
	Ar	Va	Si	Qu			
<i>Dipodomys ordii</i> Woodhouse, 1853	X						Baker and Greer (1962), MSU, KU, CRD
<i>Dipodomys phillipsii</i> Gray, 1841		X			X	Pr	Baker and Greer (1962), MSU, KU, CRD, TTU
<i>Dipodomys spectabilis</i> Merriam, 1890	X						Servín <i>et al.</i> (2003a), CRD
Family Muridae							
Subfamily Arvicolinae							
<i>Microtus mexicanus</i> (de Saussure, 1861)			X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
Subfamily Neotominae							
<i>Baiomys taylori</i> (Thomas, 1887)		X					Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Nelsonia neotomodon</i> Merriam, 1897			X		X	Pr	Baker and Greer (1962), MSU, CRD, ENCB, CNMA
<i>Neotoma goldmani</i> Merriam, 1903	X				X		Baker and Greer (1962), MSU, KU, ENCB
<i>Neotoma leucodon</i> Merriam, 1894	X	X					Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Neotoma mexicana</i> Baird, 1855		X	X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Onychomys arenicola</i> Mearns, 1896	X	X					Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Peromyscus boylii</i> (Baird, 1855)		X	X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA
<i>Peromyscus difficilis</i> (J.A. Allen, 1891)			X		X		Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Peromyscus eremicus</i> (Baird, 1858)	X	X		X			Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Peromyscus gratus</i> Merriam, 1898		X	X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Peromyscus leucopus</i> (Rafinesque, 1818)		X					Baker and Greer (1962), MSU, KU, CRD, CNMA, TTU
<i>Peromyscus maniculatus</i> (Wagner, 1845)	X	X	X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Peromyscus melanophrys</i> (Coues, 1874)		X			X		Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Peromyscus melanotis</i> J.A. Allen and Chapman, 1897			X				Baker and Greer (1962), MSU, KU, CRD, CNMA, TTU
<i>Peromyscus pectoralis</i> Osgood, 1904	X	X					Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Peromyscus schmidlyi</i> Bradley <i>et al.</i> , 2004			X		X		Bradley <i>et al.</i> (2004), CRD, ENCB, CNMA, TTU
<i>Peromyscus spicilegus</i> J.A. Allen, 1897				X	X		Baker and Greer (1962), MSU, CRD, CNMA, TTU
<i>Reithrodontomys fulvescens</i> J.A. Allen, 1894	X		X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
<i>Reithrodontomys megalotis</i> (Baird, 1858)		X					Baker and Greer (1962), MSU, CRD, CNMA, TTU
<i>Reithrodontomys montanus</i> (Baird, 1855)	X						Baker and Greer (1962)
<i>Reithrodontomys zacatecae</i> Merriam, 1901		X			X		Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA, TTU
Subfamily Sigmodontinae							
<i>Sigmodon arizonae</i> Mearns, 1890			X				Baker and Greer (1962), MSU, KU, CRD, CNMA, TTU
<i>Sigmodon fulviventer</i> J.A. Allen, 1889		X	X				Baker and Greer (1962), MSU, CRD, CNMA, TTU
<i>Sigmodon hispidus</i> Say and Ord, 1825	X	X					Baker and Greer (1962), MSU, KU, CRD
<i>Sigmodon leucotis</i> Bailey, 1902			X		X		Baker and Greer (1962), MSU, CRD, ENCB, CNMA, TTU
<i>Sigmodon ochrognathus</i> Bailey, 1902		X					Baker and Greer (1962), MSU, CRD, TTU
Order Lagomorpha							
Family Leporidae							
<i>Lepus californicus</i> Gray, 1837	X	X	X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA
<i>Lepus callotis</i> Wagler, 1830		X	X				Baker and Greer (1962), MSU, KU, CRD
<i>Sylvilagus audubonii</i> (Baird, 1858)	X	X					Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA
<i>Sylvilagus floridanus</i> (J.A. Allen, 1890)		X	X				Baker and Greer (1962), MSU, KU, CRD, ENCB, CNMA
TOTAL	48	62	57	65	23	20	

a - Previously unpublished records from museum specimens.

b - Though recently pooled in one single taxon (Piaggio and Spicer 2001) morphological evidence (CRD unpublished data) does not support this decision, and therefore we keep them as separate species.

ACKNOWLEDGMENTS: Funding for this research was provided by CONABIO project (GT015) and Instituto Politécnico Nacional (SIP 2010-0434, 2011-0349 to CLG and SIP 2010-1385 to DFGM). We thank curators and collection managers at the University of Kansas (R. Timm); Michigan State University (B. Lundrigan and L. Abraczinskas); The Museum, Texas Tech University (R. J. Baker and H. Gardner); National Mammal Collection, Instituto de Biología, UNAM, Mexico City (F. A. Cervantes and Y. Hortelano); and Mammal Collection, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Mexico City (J. C. López-Vidal and C. Elizalde-Arellano) for allowing access to the specimens under their care. Guilherme Garbino and one anonymous reviewers provided advice that helped improve the manuscript.

LITERATURE CITED

Allen, G.M. 1942. *Extinct and Vanishing Mammals of the Western Hemisphere, with the marine species of all the oceans*. Lancaster: American Committee for International Wildlife Protection. 620 p.

Allen, J.A. 1876. The American Bisons, living and extinct. *Memoirs of the Museum of Comparative Zoology, at Harvard College* 4:520 p.

Allen, J.A. 1903. List of mammals collected by Mr. J. H. Batty in New Mexico and Durango with descriptions of new species and subspecies. *Bulletin of the American Museum of Natural History* 19: 587-612.

Allen, J.A. 1904. Further notes on mammals from northwestern Durango. *Bulletin of the American Museum of Natural History* 20: 205-210.

Álvarez, T. and F. de Lachica. 1974. Zoogeografía de los vertebrados de México; p. 221-296 In. A. Flores Díaz, L. González Quintero, T. Álvarez and F. de Lachica (ed.). *El escenario geográfico*. México: Instituto Nacional de Antropología e Historia.

Álvarez, T. and J. Arroyo-Cabral. 1990. Cálculos de captura, densidad y ámbito hogareño de tres especies de roedores en un área de influencia humana en el sureste de Durango, México. *Anales de la Escuela Nacional de Ciencias Biológicas*, México 33(2): 185-210.

Álvarez, T. and O. Polaco. 1984. Estudio de los mamíferos capturados en La Michilí, sureste de Durango, México. *Anales de la Escuela Nacional*

- de Ciencias Biológicas, México* 28(1): 99-148.
- Álvarez, T., J.C. López and O.J. Polaco. 1988. Estudio de las madrigueras de la rata magueyera, *Neotoma mexicana* (Rodentia), en la Reserva de la Biosfera La Michilí, Durango, México. *Anales de la Escuela Nacional de Ciencias Biológicas, México* 32(1): 131-154.
- Applied Biostatistics, Inc. 1998. NTSYpc Ver. 2.02. New Cork: Exeter Software.
- Aragón Piña, E.E. and A. Garza Herrera. 1999. Actualización del inventario de los mamíferos silvestres de la Reserva de la Biosfera de Mapimí. *Acta Científica Potosina* 14(1): 7-25.
- Aragón, E.E., A. Garza and F.A. Cervantes. 2009. Estructura y organización de los ensambles de roedores de un bosque de la Sierra Madre Occidental, Durango, México. *Revista Chilena de Historia Natural* 82(4): 523-542.
- Arriaga, L., J.M. Espinoza, C. Aguilar, E. Martínez, L. Gómez and E. Loa (coord.). 2000. *Regiones terrestres prioritarias de México*. Escala de trabajo 1:1 000 000. México D. F.: Comisión Nacional para el Conocimiento y uso de la Biodiversidad. 609 p.
- Baker, R.H. 1956. Mammals of Coahuila, México. *University of Kansas Publications, Museum of Natural History* 9(7): 125-335.
- Baker, R.H. 1960. Mammals of the Guadiana lava field, Durango, Mexico. *Publications of the Museum, Michigan State University* 1: 303-328.
- Baker, R.H. and M.W. Baker. 1975. Montane Habitat Used by the Spotted Skunk (*Spilogale putorius*) in Mexico. *Journal of Mammalogy* 56(3): 671-673.
- Baker, R.H. and J.K. Greer. 1962. Mammals of the Mexican state of Durango. *Publications of the Museum, Michigan State University, Biological Series* 35: 25-154.
- Bradley, R.D., D.S. Carroll, M.L. Haynie, R. Muñiz-Martínez, M.J. Hamilton and C.W. Kilpatrick. 2004. A new species of *Peromyscus* from western Mexico. *Journal of Mammalogy* 85(4): 1184-1193.
- Brower, J.E. and J.H. Zar. 1977. *Field and Laboratory Methods for General Ecology*. Iowa: Wm C Brown Publishers. 226 p.
- Bell, D.M., M.J. Hamilton, C.W. Edwards, L.E. Wiggins, R. Muñiz-Martínez, R.E. Strauss, R.D. Bradley and R.J. Baker. 2001. Patterns of Karyotypic megaevolution in *Reithrodontomys*: Evidence from a Cytochrome-b phylogenetic hypothesis. *Journal of Mammalogy* 82(1): 81-91.
- Carraway, L.N. 2007. Shrews (Eulypotyphla: Soricidae) of Mexico. *Monographs of the Western North American Naturalist* 3: 1-91.
- Ceballos, G. and D. Navarro. 1991. Diversity and Conservation of Mexican Mammals. p. 167-198. In: M.A. Mares and D.J. Schmidly (ed.). *Latin American Mammalogy, History, Biodiversity and Conservation*. Norman: University of Oklahoma Press.
- Ceballos, G., J. Arroyo-Cabrales, R.A. Medellín, L. Medrano González and G. Oliva. 2005. Diversidad y Conservación de los Mamíferos de México p. 21-66 In: G. Ceballos and G. Oliva (ed.). *Los mamíferos silvestres de México*. México D.F.: CONABIO-Fondo de Cultura Económica.
- Charre-Medellín, J.F., C. López-González, A. Lozano and A.F. Guzmán. 2011. Estado actual del conocimiento sobre la nutria neotropical (*Lontra longicaudis annectens Major, 1897*) en el estado de Durango, México. *Revista Mexicana de Biodiversidad* 82(4): 1343-1347.
- Crossin, R.S., O.H. Soulé, R.G. Webb and R.H. Baker. 1973. Biotic relationships in the canon del Rio Mezquital, Durango, Mexico. *The Southwestern Naturalist* 18(2): 187-200.
- Delfín-Alfonso, C.A., C.A. López-González and N.E. Lara-Díaz. 2011. El oso negro Americano en el noroeste de México: Recuperación de registros de ocurrencia. *Acta Zoológica Mexicana (nueva serie)* 27(3): 777-801.
- De Morfi, J.A. 1777-1778. *Viaje de Indios y Diario del Nuevo México*. Reprinted 1980. México D.F.: Manuel Porrúa S.A. 446 p.
- Drake, J.J. 1958. The brush mouse *Peromyscus boylii* in southern Durango. *Publications of the Museum, Michigan State University, Biological Series* 1: 97-132.
- Hedrick, P.W. and R.J. Fredrickson. 2008. Captive breeding and the reintroduction of Mexican and red wolves. *Molecular Ecology* 17(1): 344-350.
- Helgen K.M., F.R. Cole, L.E. Helgen and D.E. Wilson. 2009. Generic revision in the Holarctic ground squirrel genus *Spermophilus*. *Journal of Mammalogy* 90(2): 270-305.
- Instituto Nacional de Estadística, Geografía e Informática (INEGI). 2005. Marco Geoestadístico 2005 versión 5.0 Electronic Database accesible at http://inegi.org.mx/geo/contenidos/geoestadistica/M_Geoestadistico.aspx. Captured on 17 December 2012.
- Gallina, S., E. Maury and V. Serrano. 1978. Hábitos alimenticios del venado cola blanca en la Reserva La Michilí, estado de Durango. p. 57-108 In: G. Halfpter (ed.). *Reservas de la Biosfera en el Estado de Durango*. México D.F.: Instituto de Ecología, A.C.
- Gallina, S. 1981. Forest ecosystems in northwestern Mexico. p. 26-56 In: Ffolliott, P. and S. Gallina (ed.). Deer biology, habitat requirements and management in western North America. México D.F.: Instituto de Ecología, A.C.
- Gallina, S. and E. Ezcurra. 1992. Deer densities in La Michilia: a reply to Galindo. *The Southwestern Naturalist* 37(4): 422-424.
- Gallina, S. and P. Folliott. 1983. Overstory-understory relationships: oak-pine forest of Sierra Madre Occidental, México. p. 19-20 In: E. Bartlett and D. R. Betters (ed.). *Overstory-understory relationships in western forests*. Western Regional Research Publication 1. Fort Collins, Colorado.
- Gardner, A.L. 1965. New bat records from the Mexican state of Durango. *Proceedings of the Western Foundation of Vertebrate Zoology* 1: 101-106.
- Goldman, E.A. 1951. *Biological investigations in Mexico*. Smithsonian Miscellaneous Collection 115: 1-476.
- Gómez Ruiz E.P., C. López-González and D.F. García-Mendoza. 2006. *Corynorhinus mexicanus y C. townsendii* (Chiroptera: Vespertilionidae) en la Sierra Madre Occidental. *Vertebrata Mexicana* 19: 7-12.
- González-Elizondo, S. 1991. Flora de Durango. Listados florísticos de México. Instituto de Biología UNAM : 1-167.
- González-Elizondo, M.S., M. González Elizondo and M.A. Márquez Linares. 2007. *Vegetación y Ecorregiones de Durango*. México D.F.: Instituto Politécnico Nacional and Plaza y Valdez Editores. 219 p.
- González-Romero, A., L. Hernández, J.W. Laundré, E. Aragón and J. López-Portillo. 2005. Monitoreo de dos comunidades de roedores en la reserva de la biosfera Mapimí, Durango, México. p. 15-26 In: V. Sánchez-Cordero and R.A. Medellín (ed.). *Contribuciones Mastozoológicas en homenaje a Bernardo Villa*. México D.F.: Instituto de Biología y Instituto de Ecología, Universidad Nacional Autónoma de México; Comisión Nacional para el Estudio de la Biodiversidad.
- Grenot, C. and V. Serrano. 1981. Ecological organization of small mammal communities at the bolsón de Mapimí (México). p. 89-100 In: R. Barbault and G. Halfter (ed.). *Ecology of the Chihuahuan Desert*. México D.F.: Instituto de Ecología, A.C.
- Hall, E.R. 1981. *The mammals of North America*. New York: John Wiley & Sons. Vol. 1: 1-600, Vol. 2: 601-1181.
- Hall, E.R. 1984. Geographic variation among brown and grizzly bears (*Ursus arctos*) in North America. *Special Publication of the Museum of Natural History, University of Kansas* 13: 1-16.
- Hernández, L. and M. Delibes. 1994. Seasonal food habits of coyotes, *Canis latrans*, in the Bolsón de Mapimí, southern Chihuahuan Desert, Mexico. *Zeitschrift für Saugetierkunde* 59(1): 82-86.
- Hernández-Huerta, A. *in litt.* Los pequeños mamíferos como indicadores de la heterogeneidad ambiental en la Reserva de la Biosfera La Michilí, México. Final report, MAB/UNESCO. 11 p.
- Hernández, L., A. González-Romero, J.W. Laundré, D. Lightfoot, E. Aragón and J. López-Portillo. 2005. Changes in rodent community structure in the Chihuahuan Desert México: Comparisons between two habitats. *Journal of Arid Environments* 60(2): 239-257.
- Houseal, T.W., I.F. Greenbaum, D.J. Schmidly, S.A. Smith and K.M. Davis. 1987. Karyotypic variation in *Peromyscus boylii* from Mexico. *Journal of Mammalogy* 68(2): 281-296.
- IUCN. 2012. IUCN Red List of Threatened Species. Version 2012.2 accesible at www.iucnredlist.org. Captured on 19 December 2012.
- Jacobo-Salcedo, M.R., A.J. Alonso-Castro and A. Zárate-Martínez. 2011. Folk medicinal use of fauna in Mapimí, Durango, México. *Journal of Ethnopharmacology* 133(2): 902-906.
- Jones, J.K., Jr. 1964. Additional records of mammals from Durango, Mexico. *Transactions of the Kansas Academy of Science* 66: 750-753.
- Jones, J.K., Jr., J.R. Choate and A. Cadena. 1972. Mammals from the Mexican state of Sinaloa II Chiroptera. *Occasional Papers Museum of Natural History, University of Kansas* 6: 1-29.
- Leopold, A. 1959. *Fauna Silvestre de México Aves y Mamíferos de Caza*. México D.F.: Pax-INIREB. 608 p.
- López-González, C. 2003. Murciélagos (Chiroptera) del estado de Durango, México: composición, distribución y estado de conservación. *Vertebrata Mexicana* 13: 15-23.
- López-González, C. 2005. Use of abandoned mines by a community of temperate bats in Durango, Mexico. *Acta Chiropterologica* 7(2): 285-292.
- López-González, C. and D.F. García-Mendoza. 2007. Mamíferos del Estado de Durango en la Colección del CIIDIR Durango: Las colecciones Biológicas frente a la pérdida de Biodiversidad. p. 174-182 In: M.S. González Elizondo, M. González Elizondo and M.A. Márquez Linares. *Vegetación y Ecorregiones de Durango*. México D.F.: Instituto Politécnico Nacional and Plaza y Valdez Editores.
- López-González, C. and L. Torres-Morales. 2004. Use of abandoned mines by two species of long eared bats, genus *Corynorhinus* (Chiroptera: Vespertilionidae) in Durango, Mexico. *Journal of Mammalogy* 85(5): 989-994.
- López-González, C., J. Rascón and F.D. Hernández-Velazquez. 2010. Population structure of migratory Mexican free-tailed bats *Tadarida brasiliensis mexicana* (Chiroptera) in a Chihuahuan Desert roost. *Chiroptera Neotropical* 16(1): 557-566.
- López-Vidal, J.C. and T. Álvarez. 1993. Biología de la rata montera *Neotoma mexicana* en La Michilí, Durango, México. p. 185-195 In: R.A. Medellín and G. Ceballos (eds.). *Avances en el estudio de*

- los mamíferos de México.* México D.F.: Publicaciones Especiales, Asociación Mexicana de Mastozoología A.C.
- Márquez-Linares, M.A. (ed.). 2007. *Ordenamiento ecológico del Estado de Durango.* Mexico: Secretaría de Medio Ambiente y Recursos Naturales-Gobierno del Estado de Durango. 194 p.
- McBride, R. 1980. The Mexican wolf (*Canis lupus baileyi*): a historical review and observations on its status and distribution. *U.S. Fish and Wildlife Service Endangered Species Report* 8: 1-38.
- Moctezuma Orozco, O. 2005. Oso plateado, p. 422-425 In: G. Ceballos y G. Oliva (ed.). *Los mamíferos silvestres de México.* México D.F.: CONABIO-Fondo de Cultura Económica.
- Montiel-Parra, G., G.A.Villegas-Guzmán and O.J. Polaco. 2001. Pseudoscorpiones asociados a nidos de *Neotoma albigenula* (Rodentia: Muridae) de Durango, México, p. 93-99 In: M. Vargas, O.J. Polaco and G. Zúñiga (ed.). *Contribuciones Entomológicas, Homenaje a la Dra. Isabel Bassols Batalla.* México D.F.: Instituto Politécnico Nacional, Escuela Nacional de Ciencias Biológicas.
- Muñiz-Martínez, R. and J. Arroyo-Cabral. 1996. El registro más norteño de la rata enana *Nelsonia neotomodon* (Rodentia: Muridae). *Vertebrata Mexicana* 2: 13-16.
- Muñiz-Martínez, R. and O.J. Polaco. 1996. Nuevos registros de simpatría de dos especies del Género *Corynorhinus* (Chiroptera: Vespertilionidae) en México. *Vertebrata Mexicana* 1: 13-16.
- Muñiz-Martínez, R. 1997. Relación roedor-vegetación en el sureste del estado de Durango, México, p 223-237 In: J. Arroyo-Cabral and O.J. Polaco (ed.). *Homenaje al profesor Ticul Álvarez* México D.F.: Colección Científica, Instituto Nacional de Antropología e Historia.
- Muñiz-Martínez, R., C. López-González, J. Arroyo-Cabral and M. Ortiz-Gómez. 2003. Noteworthy records of free-tailed bats (Chiroptera: Molossidae) from Durango, Mexico. *Southwestern Naturalist* 48(1): 138-144.
- Navarro-Serment, C.L., C.A. López-González, and J.P. Gallo-Reynoso. 2005. Occurrence of jaguar (*Panthera onca*) in Sinaloa, Mexico. *Southwestern Naturalist* 50 (1): 102-106.
- Nelson, E.W. 1925. Status of the Pronghorned antelope, 1922-1924. *United States Department of Agriculture, Department Bulletin* 1346:1-64.
- Pacheco, R. J. Bisonte Americano, p. 505-506 In: G. Ceballos and G. Oliva (ed.). *Los mamíferos silvestres de México.* CONABIO-Fondo de Cultura Económica.
- Petersen, M.K. 1973. Interactions between the cotton rats, *Sigmodon fulviventer* and *S. hispidus*. *American Midland Naturalist* 90(2): 319-333.
- Petersen, M.K. 1975. An analysis of multiple captures in several rodents from Durango, Mexico. *Journal of Mammalogy* 56(3): 703-705.
- Petersen, M.K. 1976. Noteworthy range extensions of some mammals in Durango, Mexico. *The Southwestern Naturalist* 21(1): 139-142.
- Petersen, M.K. 1978. *Rodent ecology and natural history, observations on the mammals of Atotonilco de Campa, Durango.* Iowa: Carter Press. 129 p.
- Petersen, M.K. 1979. Some morphometric observations on *Notiosorex crawfordi* from Durango, Mexico. *Occasional Papers in Zoology of Michael K. Petersen* 1: 1-9.
- Petersen, M.K. 1980a. The taxonomic status of a dusky-colored population of *Peromyscus* at Atotonilco, Durango, Mexico. *Occasional Papers in Zoology of Michael K. Petersen* 5: 1-19.
- Petersen, M.K. 1980b. A comparison of small mammal populations sampled by pit-fall and live-traps in Durango, Mexico. *The Southwestern Naturalist* 25(1): 122-124.
- Petersen, M.K. 1993. Dietary overlap and livestock forage relationships in two species of *Sigmodon* from Durango, México. p. 289-300 In: R.A. Medellín and G. Ceballos (ed.). *Avances en el estudio de los mamíferos de México* México D.F.: Asociación Mexicana de Mastozoología.
- Petersen, M.K. and M.K. Petersen. 1979. A temporal comparison of owl pellet contents with small mammal population levels in Durango, Mexico. *Occasional Papers in Zoology of Michael K. Petersen* 4: 1-27.
- Piaggio A.J. and G.S. Spicer. 2001. Molecular Phylogeny of the Chipmunks inferred from mitochondrial Cytochrome b and Cytochrome Oxidase II Gene Sequences. *Molecular Phylogenetics and Evolution* 20(3): 335-350.
- Portales, G.L., L. Hernández, F.A. Cervantes and J.W. Laundré. 2004. Reproduction of Black-tailed jackrabbits (Lagomorpha: *Lepus californicus*) in relation to environmental factors in the Chihuahuan Desert, Mexico. *The Southwestern Naturalist* 49(3): 359-366.
- Ramírez-Pulido, J., A. Castro-Campillo, M.A. Armella and A. Salame-Méndez. 2000. *Bibliografía reciente de los mamíferos de México 1994-2000.* México: Universidad Autónoma Metropolitana, 280 p.
- Rogovin, K., G.Shenbrot and A. Surov. 1991. Analysis of spatial organization of a desert rodent community in the Bolson de Mapimí, Mexico. *Journal of Mammalogy* 72(2): 347-359.
- Romo-Vázquez, E., L. León-Paniagua and O. Sánchez. 2005. A new species of *Habromys* (Rodentia:Neotominae) from México. *Proceedings of the Biological Society of Washington* 118(3): 605-618.
- Secretaría del Medio Ambiente y Recursos Naturales (SEMARNAT). 2010. *Norma Oficial Mexicana NOM-059-ECOL-2010, Protección de especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en riesgo.* Diario Oficial de la Federación, México, December 30th: 1-78.
- Servín, J. 1991. Algunos aspectos de la conducta social del lobo mexicano (*Canis lupus baileyi*) en cautiverio. *Acta Zoológica Mexicana (nueva serie)* 45: 1-41.
- Servín, J. 1997. El periodo de apareamiento, nacimiento y crecimiento del lobo mexicano (*Canis lupus baileyi*). *Acta Zoológica Mexicana (nueva serie)* 71: 45-56.
- Servín, J. 2000. Duration and frequency of chorus howling of the Mexican wolf (*Canis lupus baileyi*). *Acta Zoológica Mexicana (nueva serie)* 80: 223-231.
- Servín, J. 2005. La cacería como estrategia conductual para determinar el tamaño de grupo del lobo mexicano. p. 461-474 In: V. Sánchez-Cordero and R.A. Medellín (ed.). *Contribuciones mastozoológicas en Homenaje a Bernardo Villa.* México D.F.: Instituto de Biología and Instituto de Ecología, Universidad Nacional Autónoma de México; Comisión Nacional para el Estudio de la Biodiversidad.
- Servín, J. and C. Huxley. 1991a. La dieta del coyote en un bosque de encino-pino de la Sierra Madre Occidental de Durango, México. *Acta Zoológica Mexicana (nueva serie)* 44: 1-26.
- Servín, J. and C. Huxley. 1991b. Algunos aspectos de la conducta social del lobo mexicano (*Canis lupus baileyi*) en cautiverio. *Acta Zoológica Mexicana (nueva serie)* 45: 1-43.
- Servín, J. and C. Huxley. 1993. Biología del coyote (*Canis latrans*) en la reserva de la biosfera "La Michilía" Durango, p. 197-204 In: R.A. Medellín and G. Ceballos (ed.). *Avances en el estudio de los mamíferos de México*, México D.F.: Publicaciones especiales, Asociación Mexicana de Mastozoología A.C.
- Servín, J., C. Huxley and E. Chacon. 1995. First record of the spotted skunk *Spilogale putorius* (Carnivora: Mustelidae) in the State of Durango, Mexico. *The Southwestern Naturalist* 40(2): 227-228.
- Servín, J., V. Sánchez-Cordero and F.A. Cervantes. 1996. First record of the Sierra Madre mantled ground squirrel (*Spermophilus madrensis*; Rodentia: Sciuridae) from Durango, Mexico. *The Southwestern Naturalist* 41(1): 189-190.
- Servín, J., E. Chacón, N. Alonso-Pérez and C. Huxley. 2003a. New records of mammals from Durango, Mexico. *The Southwestern Naturalist* 48(1): 136-138.
- Servín, J., V. Sánchez-Cordero and S. Gallina. 2003b. Distances traveled daily by coyotes, *Canis latrans* in a pine-oak forest in Durango, Mexico. *Journal of Mammalogy* 84(2): 547-552.
- Tejedor, A. 2005. A new species of funnel-eared bat (Natalidae: *Natalus*) from Mexico. *Journal of Mammalogy* 86(6): 1109-1120.
- Tejedor, A. 2006. The type locality of *Natalus stramineus* (Chiroptera:Natalidae): implications for the taxonomy and biogeography of the genus *Natalus*. *Acta Chiropterologica* 8(2): 361-380.
- Torres-Morales, L., D.F. García-Mendoza, C. López-González and R. Muñiz-Martínez. 2010. Bats of Northwestern Durango, Mexico: Species Richness at the Interface of Two Biogeographic Regions. *The Southwestern Naturalist* 55(3): 347-362.
- Thomas, O. 1882. On a small collection of mammals from central Mexico. *Proceedings of the Zoological Society of London* 50(2): 371-372.
- Villegas-Guzmán, G.A., C. López-González, and M. Vargas. 2005. Ectoparasites associated to two species of *Corynorhinus* (Chiroptera: Vespertilionidae) from the Guanaceví mining region, Durango, Mexico. *Journal of Medical Entomology* 42: 125-127.
- Webb, R.G. and R.H. Baker. 1962. Terrestrial Vertebrates of the Pueblo Nuevo Area of Southwestern Durango, Mexico, *American Midland Naturalist* 68(2): 325-333.
- Wilson, D. and S. Ruff. 1999. *The Smithsonian book of North American Mammals.* Washington: Smithsonian Institution Press. 750 p.

RECEIVED: April 2011

ACCEPTED: March 2013

PUBLISHED ONLINE: April 2013

EDITORIAL RESPONSIBILITY: Fabio Oliveira do Nascimento